

RESOLUTIONS WITH CONICAL SLICES AND DESCENT FOR THE BRAUER GROUP CLASSES OF CERTAIN CENTRAL REDUCTIONS OF DIFFERENTIAL OPERATORS IN CHARACTERISTIC p

DMITRY KUBRAK AND ROMAN TRAVKIN

*“Even more so is the word “crystalline”,
a glacial and impersonal concept of his
which disdains viewing existence from a
single portion of time and space”*

Eileen Myles, *“The Importance of Being
Iceland”*

ABSTRACT. For a smooth variety X over an algebraically closed field of characteristic p , to a differential 1-form α on the Frobenius twist $X^{(1)}$ one can associate an Azumaya algebra $\mathcal{D}_{X,\alpha}$, defined as a certain central reduction of the algebra $\mathcal{D}_X$ of “crystalline differential operators” on X . For a resolution of singularities $\pi : X \rightarrow Y$ of an affine variety Y , we study for which α does the class $[\mathcal{D}_{X,\alpha}]$ in the Brauer group $\mathrm{Br}(X^{(1)})$ descend to $Y^{(1)}$. In the case when X is symplectic, this question is related to Fedosov quantizations in characteristic p and the construction of non-commutative resolutions of Y . We prove that the classes $[\mathcal{D}_{X,\alpha}]$ descend étale locally for all α if $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^{1,2} \pi_* \mathcal{O}_X = 0$. We also define a certain class of resolutions which we call resolutions with conical slices, and prove that for a general reduction of a resolution with conical slices in characteristic 0 to an algebraically closed field of characteristic p classes $[\mathcal{D}_{X,\alpha}]$ descend to $Y^{(1)}$ globally for all α . Finally we give some examples, in particular we show that Slodowy slices, Nakajima quiver varieties and hypertoric varieties are resolutions with conical slices.

CONTENTS

1. Introduction	2
1.1. Motivation	3
1.2. Plan of the paper	5
1.3. Remaining questions	9
1.4. Acknowledgements	9
2. Frobenius twist and differential operators	9
2.1. Frobenius twist	9
2.2. Cartier operator	10
2.3. Crystalline differential operators	11
2.4. Azumaya algebras $\mathcal{D}_{X,\alpha}$	12
2.5. Azumaya algebras $\mathcal{D}_{X,\alpha}$ and p -curvature	12
2.6. The gerbes $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$	15
2.7. Additivity of the map c_X : cohomology	17
2.8. Additivity of the map c_X : Picard groupoids	18
3. Descent étale locally	20
3.1. Setup	20

3.2. Splitting étale locally on Y	20
4. Obstructions to global descent	23
4.1. Picard obstruction	23
4.2. Sheaves $\mathcal{Q}_{\pi,N}$	23
5. Resolutions with conical slices	26
5.1. Setup	26
5.2. Category of étale germs of resolutions	26
5.3. Separable $\mathbb{G}_m$ -action	30
5.4. Definition of a resolution with conical slices	31
5.5. Construction of a slice	32
5.6. Reductions to characteristic p	34
6. $\mathbb{G}_m$ -weights of 1-forms on a conical resolution	36
6.1. Totally positive forms	40
6.2. Application to resolutions with conical slices	44
7. Descent for resolutions with conical slices	44
7.1. Sheaf $\mathcal{Q}_{\pi,N}$ and étale equivalences	44
7.2. Descent for a resolution with conical slices	45
8. Examples of resolutions with conical slices	47
8.1. Background on symplectic resolutions	47
8.2. Marsden-Weinstein reductions for representations of quivers	47
8.3. Hamiltonian reductions of vector spaces	55
8.4. Slodowy slices	61
References	62

1. INTRODUCTION

¹ Let X be a smooth variety over an algebraically closed field $\mathbb{k}$ of characteristic p . To each differential 1-form α on the Frobenius twist $X^{(1)}$ one can associate an Azumaya algebra $\mathcal{D}_{X,\alpha}$ on $X^{(1)}$ in the following way. The sheaf of “crystalline differential operators” $\mathcal{D}_X$ on X defines an Azumaya algebra on the cotangent bundle $T^*X^{(1)}$. Given a differential 1-form α on $X^{(1)}$, restricting $\mathcal{D}_X$ to the graph $\Gamma_\alpha \subset T^*X^{(1)}$ defines an Azumaya algebra $\mathcal{D}_{X,\alpha}$ on $X^{(1)}$. In this paper we study the following question: given a resolution of singularities $\pi : X \rightarrow Y$ for which differential 1-forms α does the class $[\mathcal{D}_{X,\alpha}]$ in the Brauer group of $X^{(1)}$ descend to $Y^{(1)}$? We prove that $[\mathcal{D}_{X,\alpha}]$ descends étale locally on $Y^{(1)}$ for any α if $\mathcal{O}_Y \xrightarrow{\sim} \pi_*\mathcal{O}_X$ and $R^i\pi_*\mathcal{O}_X = 0$ for $i = 1, 2$. We also define a class of resolutions which we call resolutions with conical slices, such that given a resolution with conical slices over a field of characteristic 0, for a general enough reduction $\pi_s : X_s \rightarrow Y_s$ to an algebraically closed field $\mathbb{k}_s$ of characteristic p , the classes $[\mathcal{D}_{X_s,\alpha}]$ descend to $Y_s^{(1)}$ for all α . We also prove that quiver and hypertoric varieties are resolutions with conical slices.

In the rest of the introduction we try to provide a motivation for this question. If the reader feels motivated enough already they can pass directly to Sect. 1.2 where they will find a detailed plan of the paper.

¹This material is based upon work supported by The Wolfensohn Fund and the National Science Foundation under agreement No. DMS-1128155. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

1.1. Motivation. Given a singular variety Y over a field of characteristic 0 one wants to study its resolutions of singularities $\pi : X \rightarrow Y$. Usually there are too many of them, but one can impose certain additional conditions on the resolutions. In particular for a normal singular conical Poisson variety Y , one can study its symplectic resolutions $\pi : X \rightarrow Y$. One can then try to classify all of them, and in some form this was done recently by Namikawa [Na1]. Namely, provided there exists at least one symplectic resolution $\pi : X \rightarrow Y$, the vector space $V_{\mathbb{R}} = \text{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{R}$ can be partitioned into a union of rational cones and there is an action of a finite group W on $V_{\mathbb{R}}$ that maps cones to cones. The set of symplectic resolutions $\pi : X \rightarrow Y$ is then identified with the set of cones modulo the action of W .

One can ask given two symplectic resolutions $\pi : X \rightarrow Y$, $\pi' : X' \rightarrow Y$ if their derived categories of coherent sheaves are equivalent. This is a particular case of “ K -equivalence implies D -equivalence” conjecture due to Kawamata [Kaw] (see also Conjecture 5.1 in the ICM talk by Bondal and Orlov [BO]). Étale locally on the base this was proved first by Bezrukavnikov and Kaledin for quotient singularities in [BK2] and then separately by Kaledin for general symplectic resolution in [Kal3]. Both proofs are based on the notion of a Fedosov quantization in characteristic p , and this notion is what connects the algebras $\mathcal{D}_{X,\alpha}$ and symplectic resolutions. We will first sketch Kaledin’s proof and then will show how classes $[\mathcal{D}_{X,\alpha}]$ naturally pop up in the context and why it can be important to study their descent.

Results of [BK1] imply the existence of a canonical Frobenius-constant quantization $\mathcal{A}_0$ of X . By definition of being Frobenius-constant, $\mathcal{A}_0$ possesses an isomorphism $\mathcal{Z}(\mathcal{A}_0) = \mathcal{O}_X^p$ and $(\text{Fr}_X)_* \mathcal{A}_0$ is an Azumaya algebra on $X^{(1)}$. For a class $\lambda \in \text{Pic}(X)$ one can define $\mathcal{A}_\lambda$ as the twist of $\mathcal{A}_0$ by the corresponding linear bundle $\mathcal{O}(\lambda)$. It is known that the isomorphism class of $\mathcal{A}_\lambda$ depends only on $\bar{\lambda} \in \text{Pic}(X)/p \cdot \text{Pic}(X)$ and that all $\mathcal{A}_\lambda$ have the same class in the Brauer group. One can prove that $(\text{Fr}_X)_* \mathcal{A}_\lambda$ splits on some étale neighbourhood $j : U \rightarrow Y$ of each point $y \in Y$, namely $j^*((\text{Fr}_X)_* \mathcal{A}_\lambda) = \underline{\text{End}}_{\mathcal{O}_{X^{(1)}}}(\mathcal{E}_U)$ for some vector bundle $\mathcal{E}_U$ on $(U \times_Y X)^{(1)}$. Using the fact that $\mathcal{A}_\lambda$ was a quantization one can prove that $\mathcal{E}_U$ is a tilting object, namely $R^\bullet \text{Hom}(\mathcal{E}_U, \mathcal{E}_U) = \Gamma(j^* \mathcal{A}_\lambda)$ and the functor $R^\bullet \text{Hom}(\mathcal{E}_U, \bullet)$ gives an equivalence between the derived category of coherent sheaves on $(U \times_Y X)^{(1)}$ and finitely generated $\Gamma(j^* \mathcal{A}_\lambda)$ -modules. We have $\text{Pic}(X) = \text{Pic}(X')$ and the quantization $\mathcal{A}'_\lambda$ of X' is compatible with $\mathcal{A}_\lambda$ in the sense that it has a splitting bundle $\mathcal{E}'_U$, such that $\mathcal{E}_U$ is isomorphic to $\mathcal{E}'_U$ on the smooth part of Y . Finally one has $\Gamma(j^* \mathcal{A}_\lambda) = \Gamma(j^* \mathcal{A}'_\lambda)$ and we obtain derived equivalences

$$D^b(\text{Coh}((X \times_Y U)^{(1)})) \cong D^b(\Gamma(j^* \mathcal{A}_\lambda) - \text{mod}^{\text{fg}}) \cong D^b(\text{Coh}((X' \times_Y U)^{(1)})).$$

Bundles $\mathcal{E}_U$, $\mathcal{E}'_U$ are rigid and can be lifted to tilting objects in characteristic 0, providing a derived equivalence there. However the resulting equivalence we will obtain will depend on the original choice of $\lambda \in \text{Pic}(X)$ and one could try to keep track of this. One could also try to extend this sort of equivalence to a global one. In the same paper Kaledin proves that if $\pi : X \rightarrow Y$ is conical, then the corresponding tilting object on étale neighbourhood of the central point $y_0 \in Y$ comes as a pull-back of a $\mathbb{G}_m$ -equivariant sheaf $\mathcal{E}_0$ on X which then produces a global equivalence. However the Azumaya algebra $\mathcal{A}_\lambda$ is not globally split and it is not clear what is the global relation between $\mathcal{A}_\lambda$ and $\mathcal{E}_0$, as well as the relation between $\mathcal{E}_0$ and splitting bundles for $\mathcal{A}_X$ at the other points.

✂ This is the place where one can need the Azumaya algebras $\mathcal{D}_{X,\alpha}$. Namely one can give another description of the class $[\mathcal{A}_\lambda]$ in $\text{Br}(X^{(1)})$. In the case when X admits an open embedding to the cotangent bundle to a stack it is known that $[\mathcal{A}_\lambda] = [\mathcal{D}_{X,\alpha}]$ for a 1-form α . The form α can be written down, namely it is given by the contraction of the symplectic 2-form $\omega \in H^0(X, \Omega_X^2)$ with the Euler vector field ξ provided by the contracting $\mathbb{G}_m$ -action (see e.g. the proof of Proposition 10.3 in [BL]). In particular this is true for all Hamiltonian

reductions of the cotangent bundle of a vector space, which is the main set of examples we consider. This is also expected to be true for any conical symplectic resolution $\pi : X \rightarrow Y$.

Though the Azumaya algebra $(\mathrm{Fr}_X)_* \mathcal{A}_\lambda$ is usually not split even for the simplest examples, it is reasonable to expect that the class $[\mathcal{A}_\lambda]$ in the Brauer group $\mathrm{Br}(X^{(1)})$ (which does not depend on the choice of λ) descends to $Y^{(1)}$: namely that there exists a class $c \in \mathrm{Br}(Y^{(1)})$, such that $[\mathcal{A}_\lambda] = \pi^*(c)$. Let $\mathcal{C}$ be an Azumaya algebra on $Y^{(1)}$, such that $c = [\mathcal{C}]$. Then $\mathcal{A}_\lambda \otimes_{\mathcal{O}_{X^{(1)}}} \pi^* \mathcal{C}^{op}$ is split for any $\lambda \in \mathrm{Pic}(X)$ and so we obtain an equivalence

$$\mathrm{Coh}(X^{(1)}) \cong \mathrm{Coh}(\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op}),$$

where $\mathrm{Coh}(\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op})$ is the abelian category of $\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op}$ -modules which are $\mathcal{O}_{X^{(1)}}$ -coherent. We have $\Gamma(\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op}) = \Gamma(\mathcal{A}_\lambda) \otimes \mathcal{C}^{op}$ and for each $\lambda \in \mathrm{Pic}(X)$ we obtain a composite functor

$$R\Gamma_{\lambda, \mathcal{C}} : D^b(\mathrm{Coh}(X^{(1)})) \cong D^b(\mathrm{Coh}(\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op})) \xrightarrow{R\Gamma_\lambda} D^b(\Gamma(\mathcal{A}_\lambda) \otimes \mathcal{C}^{op} - \mathrm{mod}^{\mathrm{fg}}).$$

which is a derived equivalence whenever $R\Gamma_\lambda$ is. **To summarise:** if we want to define some derived equivalences between $D^b(\mathrm{Coh}(X^{(1)}))$ and $D^b(R_\lambda - \mathrm{mod}^{\mathrm{fg}})$ for some non-commutative resolutions R_λ indexed by $\lambda \in \mathrm{Pic}(X)$ there is a natural way to do that if the class $[\mathcal{A}_\lambda]$ descends to $Y^{(1)}$. The case of $[\mathcal{A}_\lambda]$ being of the form $[\mathcal{D}_{X_\alpha}]$ returns us to the question formulated in the beginning.

We prove that all classes $[\mathcal{D}_{X_\alpha}]$ descend étale locally on the base for all α if $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$ and we define a class of resolutions (which we call resolutions with conical slices) for which the descent is true globally (for big enough p). We also prove that there are a lot of examples of such, e.g. any resolution of singularities of a Hamiltonian reduction of a vector space by GIT-quotient with a non-trivial character is a resolution with conical slices (provided it is a resolution of singularities). In particular resolutions of singularities of hypertoric and Nakajima quiver varieties are resolutions with conical slices.

Remark 1.1.1. Following what we have written above, for regular λ (ones for which $R\Gamma_\lambda$ is an equivalence) we obtain equivalences

$$R\Gamma_{\lambda, \mathcal{C}} : D^b(\mathrm{Coh}(X^{(1)})) \xrightarrow{\sim} D^b(\Gamma(\mathcal{A}_\lambda) \otimes \mathcal{C}^{op} - \mathrm{mod}^{\mathrm{fg}})$$

for some Azumaya algebra $\mathcal{C}$. In this case $\mathcal{A}_\lambda$ also splits on some étale neighbourhoods of the fibers and so for any $y \in Y$, $R\Gamma_{\lambda, \mathcal{C}}$ provides an equivalence of sheaves with support at $\pi^{-1}(y)$ and modules with support at y :

$$R\Gamma_{\lambda, \mathcal{C}, y} : D^b(\mathrm{Coh}_{\pi^{-1}(y)}(X^{(1)})) \xrightarrow{\sim} D^b(\Gamma(\mathcal{A}_\lambda)_y - \mathrm{mod}^{\mathrm{fg}}).$$

It is known that for any symplectic resolution (over $\mathbb{C}$) $K^0(\mathrm{Coh}_{\pi^{-1}(y)}(X)) \otimes_{\mathbb{Z}} \mathbb{Q} \cong H^\bullet(X_y, \mathbb{Q})$ (Corollary 1.10, [Kal3]) and as K^0 remains the same for generic reduction we get a canonical basis in $H^\bullet(X_y, \mathbb{Q})$ corresponding to the basis of irreducible modules in $K^0(\Gamma(\mathcal{A}_\lambda)_y - \mathrm{mod}^{\mathrm{fg}})$ (depending on λ). This can be considered as a partial generalisation of the results of [BMR].

The program outlined by Bezrukavnikov and Okounkov and partially written down in Conjecture 1 of [ABM] proposes that there is some interesting structure behind equivalences $R\Gamma_\lambda$. As a part of the conjecture, categories $D^b(\mathrm{Coh}(\mathcal{A}_\lambda \otimes \pi^* \mathcal{C}^{op}))$ indexed by $\lambda \in \mathrm{Pic}(X)$, categories $D^b(\mathrm{Coh}(X_C^{(1)}))$ indexed by Namikawa's cones C , and derived equivalences $R\Gamma_{\lambda, \mathcal{C}}$ (for various X_C) should give rise to a representation of the Poincaré groupoid of a certain space $V_{\mathbb{C}}^\circ$, which is a complement of $V_{\mathbb{C}} = \mathrm{Pic}(X) \otimes_{\mathbb{Z}} \mathbb{C}$ to a certain affine hyperplane arrangement $\Sigma \subset V_{\mathbb{C}}$. However the precise statement of this conjecture is about the sheaves with support at the central fiber $\pi^{-1}(y_0)$ and our paper also provides a way to reformulate it for the global situation.

1.2. Plan of the paper. Logistically the paper can be divided into two parts. The first one is short and relatively simple, it studies the local side of the story. The second one takes almost all the space of the paper and is pretty technical. It attempts to give an answer for the global splitting in some cases.

In Sect. 2 we give a brief tour through the theory of “crystalline differential operators” in characteristic p and define Azumaya algebras $\mathcal{D}_{X,\alpha}$ on the Frobenius twist $X^{(1)}$. The content of this section intersects in a significant way with sections 4.1-4.3 in [OV]. Algebras $\mathcal{D}_{X,\alpha}$ are the main objects of study in this paper and we try to show how they are related to some more widely known objects in algebraic and differential geometry in characteristic p . In Sect. 2.1 we remind what the relative Frobenius twist $X^{(S)}$ of a scheme X over S is and also fix the notations which will be used later in the paper. In Sect. 2.2 we recall the statement of the Cartier isomorphism (Theorem 2.2.1) and define the Cartier operator $C : (F_X)_* \Omega_{X,cl}^1 \rightarrow \Omega_{X^{(1)}}^1$. In Sect. 2.3 we remind the definition of “crystalline differential operators” and discuss its Azumaya property (Proposition 2.3.1): namely the sheaf of differential operators $\mathcal{D}_X$ defines an Azumaya algebra on $T^*X^{(1)}$. In Sect. 2.4 to each differential form α on $X^{(1)}$ we associate an Azumaya algebra $\mathcal{D}_{X,\alpha}$ on $X^{(1)}$ as the restriction of $\mathcal{D}_X$ to the graph $\Gamma_\alpha \subset T^*X^{(1)}$ (see Definition 2.4.1). This way we obtain a map $c_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$. Sect. 2.5 establishes a connection between algebras $\mathcal{D}_{X,\alpha}$ and p -curvature. We first remind the definition of p -curvature of a sheaf with a flat connection and relate it to algebras $\mathcal{D}_{X,\alpha}$ (see e.g. Proposition 2.5.1 and Proposition 2.5.2). In particular we show how the Azumaya property of $\mathcal{D}_X$ implies the classical Cartier theorem (see Remark 2.5.2). For each differential 1-form α we define two groupoids: $\text{Spl}(X, \alpha)$ of splittings of $\mathcal{D}_{X,\alpha}$ and $LIC(X, \alpha)$ of line bundles with p -curvature α , and show that they are equivalent (Proposition 2.5.2). In Sect. 2.6 we provide a brief introduction to the theory of gerbes and show that the groupoids $\text{Spl}(U, \alpha)$ and $LIC(U, \alpha)$ for various opens $U \rightarrow X$ give rise to gerbes $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$ on $X^{(1)}$. The equivalence from Proposition 2.5.2 provides an equivalence of the corresponding gerbes (Proposition 2.6.1). In Sect. 2.7 we try to prove that the map c_X is additive. We construct some (a priori different) map $\bar{c}_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$ coming from a certain 4-term exact sequence (see Proposition 2.7.1). In particular we observe that its exactness essentially follows from the Azumaya property of $\mathcal{D}_X$. Finally, in Sect. 2.8 we introduce Picard groupoids and recall the equivalence between the category of Picard groupoids and complexes of sheaves with length 2. We deduce from this that $c_X = \bar{c}_X$ and so c_X is additive (Proposition 2.8.1). We end Sect. 2 with Corollary 2.8.1 which provides a set of α for which $c_X(\alpha) = 0$ and or, in other words, the Azumaya algebra $\mathcal{D}_{X,\alpha}$ is split.

In Sect. 3 we study if, given a proper morphism $\pi : X \rightarrow Y$ from a smooth variety X , the classes $[\mathcal{D}_{X,\alpha}] \in \text{Br}(X^{(1)})$ descend to $Y^{(1)}$ étale locally. We prove the following result (Theorem 3.2.1) which answers this question in almost full generality:

Theorem 1.2.1. *Let $\pi : X \rightarrow Y$ be a proper map from a smooth variety X such that $R^1\pi_*\mathcal{O}_X = R^2\pi_*\mathcal{O}_X = 0$. Then there exists an étale cover $U \rightarrow Y$, such that the pull-back of $\mathcal{D}_{X,\alpha}$ to $U \times_Y X$ is split.*

The core of our proof is Lemma 3.2.1 which shows that if we add a Frobenius-linear operator A to a surjective $\mathbb{k}$ -linear operator B between finite-dimensional vector spaces, we still get a surjective operator. The next important step is Lemma 3.2.2 where we prove that under assumptions $R^i\pi_*\mathcal{O}_X = 0$ for $i = 1, 2$, the Cartier operator $\pi_*C : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_X^1$ is a surjection. We then use Lemma 3.2.1 for $A = \bullet^{(1)}$ and $B = C$ to prove that the restriction of any differential 1-form α on $X^{(1)}$ on the formal neighbourhood of any fiber $\pi^{-1}(y)$ is of the form $\omega^{(1)} - C(\omega)$ for some ω . From this we deduce that $\mathcal{D}_{X,\alpha}$ is split on the formal neighbourhoods

of the fibers of π . Finally, Popescu's theorem allows us to extend the splitting to some étale neighbourhood of the fiber, which leads directly to the statement of Theorem 3.2.1.

Sect. 4 presents two approaches for studying the global descent. The first is *Picard obstruction*, it is defined in Sect. 4.1 and is a class in $H_{\text{ét}}^1(Y, \text{Pic}_{X/Y})[p]$. This obstruction is very natural, but seems somewhat useless for us, or at least we have not found a way to effectively work with it. Instead, in Sect. 4.2, for the case of an affine $Y = \text{Spec } A$, we define another type of obstructions, which we call $Q_{\pi, N}$ (see Definition 4.2.1). By definition for each N there is a natural projection $H^0(X, \Omega_X^1) \twoheadrightarrow Q_{\pi, N}$ and spaces $Q_{\pi, N}$ for different N are organised into a chain of surjections:

$$H^0(X, \Omega_X^1) \twoheadrightarrow Q_{\pi, 0} \twoheadrightarrow Q_{\pi, 1} \twoheadrightarrow \cdots \twoheadrightarrow Q_{\pi, N} \twoheadrightarrow \cdots$$

Moreover, if a differential 1-form α maps to 0 in some $Q_{\pi, N}$, the corresponding class $[D_{X, \alpha}] \in \text{Br}(X^{(1)})$ descends to $Y^{(1)}$ (see remark in the Definition 4.2.1). The idea of definition is based on an interesting observation: the Cartier operator $C : H^0(X, \Omega_{X, \text{cl}}^1) \rightarrow H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ does not map closed Kähler differentials $H^0(Y, \Omega_{Y, \text{cl}}^1)$ back to Kähler differentials $H^0(Y^{(1)}, \Omega_{Y^{(1)}}^1)$. Moreover, sometimes it can even happen that any globally defined 1-form on $X^{(N)}$ is obtained from a pull-back of a Kähler differential on Y by applying Cartier operator big enough amount of times. Spaces $Q_{\pi, N}$ are defined as the obstruction for this to be true. The main good thing about spaces $Q_{\pi, N}$ is that they have a natural structure of $(A)^{p^N}$ -module, such that the projection $H^0(X, \Omega_X^1) \twoheadrightarrow Q_{\pi, N}$ is $(A)^{p^N}$ -linear. This allows to study the corresponding coherent sheaves $\mathcal{Q}_{\pi, N}$ on $Y^{(N)}$ using the means of algebraic geometry. The relation between the descent of $[D_{X, \alpha}]$ and Kähler differentials is given by Lemma 4.2.1: namely, if $\alpha = \pi^* \theta$ for some Kähler differential θ , then $[D_{X, \alpha}]$ descends to Y . Also take notice of Remark 4.2.2 which comments on how the map c_X can be extended to the case of a singular X . Finally, very important is Lemma 4.2.2 where we show that given a smooth S , if $Q_{\pi, N} = 0$ then $Q_{\pi \times \text{id}_S, N} = 0$, where $\pi \times \text{id}_S : X \times S \rightarrow Y \times S$ is the product of $\pi : X \rightarrow Y$ with $\text{id}_S : S \rightarrow S$.

In Sect. 5 we develop the main technical tools for this paper, namely we define *the category of étale germs of resolutions* (Sect. 5.2), *étale equivalences* between *pointed resolutions* (Definition 5.2.2 and Definition 5.2.4) and *resolutions with conical slices* (Definition 5.4.3). The idea is to define a class of resolutions for which we could apply Lemma 4.2.2 and proceed by induction: we want to have a lot of points on the base of the resolution, that have a neighbourhood in a suitable topology which decomposes as a product of some smooth variety and a resolution of the same sort. The starting point is to take conical resolutions (Definition 5.4.1), then by Luna's étale slice theorem every non-central point has an étale decomposition into a product of its orbit of $\mathbb{G}_m$ -action and a slice to the orbit. But for our purposes this is not enough, as we will not have any information about the slice. However the right definition is not very far from this one, in addition we just need to ask for a contracting $\mathbb{G}_m$ -action in an étale neighbourhood of each non-central point (Definition 5.4.3). Some technical work needs to be done to make it easy to work with that definition. We define the notion of *étale equivalence* (Definition 5.2.4) between two pointed resolutions as an invertible morphism in a certain category $\text{Res}_*^{\text{ét}}$, which we call the category of étale germs of resolutions. We define a *resolution with conical slices* as a conical resolution $\pi : X \rightarrow Y$ such that $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$, $R^i \pi_* \mathcal{O}_X = 0$ for $i = 1, 2$, and at every non-central point y it is étale equivalent to some conical resolution $\pi' : X' \rightarrow Y'$ with the central point y' . In Sect. 5.3 we define the separability of a $\mathbb{G}_m$ -action (Definition 5.3.2). **Warning:** note that our definition of conical resolution by default includes separability assumption on the action of $\mathbb{G}_m$! In Sect. 5.5 we prove that at each non-central point a resolution with conical slices has a slice which is again a resolution with conical slices (Proposition 5.5.1). This is the statement that will allow us to make the induction step while studying spaces $Q_{\pi, N}$

for resolutions with conical slices. Finally in Sect. 5.6 we prove that given a resolution with conical slices in characteristic 0, general reduction to characteristic p is again a resolution with conical slices (Proposition 5.6.3).

The notion of a resolution with conical slices seems to be interesting even outside of the context of this paper. This is a class of resolutions for which any statement which is local in étale topology and which is invariant under multiplication on $\mathbb{A}^1$, is enough to check only for some neighbourhood of the central point. As we will see quite a few well-known resolutions are resolutions with conical slices (Sect. 8). The power of this tool is demonstrated on the example of spaces $Q_{\pi,N}$ (and sheaves $\mathcal{Q}_{\pi,N}$) and it looks plausible that they can be used in some other situations too.

The reason why we need étale slices and say not formal ones is that we need something like Remark 5.5.1 which says that there is an étale correspondence between the complement to a central point in $\pi : X \rightarrow Y$ and a finite union of products of resolutions with conical slices with $\mathbb{A}^1$. Finiteness is very important here if we want to prove something like Proposition 5.6.3 or have a universal bound for N such that $Q_{\pi,N} = 0$ on each slice.

In Sect. 6 we study weights of differential 1-forms on a conical resolution. The $\mathbb{G}_m$ -action on $\pi : X \rightarrow Y$ produces a positive grading on the ring of global functions on X (and Y). A sheaf of differential 1-forms Ω_X^1 is $\mathbb{G}_m$ -equivariant and so we also have grading on $H^0(X, \Omega_X^1)$ (by the weights of $\mathbb{G}_m$ -action), which turns it into a graded module over the ring of functions. Interesting question is: what can we say about this grading? For example: is it true that the grading is positive? Assuming that $R^i\pi_*\mathcal{O}_X = 0$ for $i = 1, 2$ we can say something about that (see Theorem 6.0.1). Here we write down explicitly one part of the result (Corollary 6.0.1):

Theorem 1.2.2. *Let $\pi : X \rightarrow Y$ be a conical resolution of singularities over an algebraically closed field of characteristic 0 with $\mathcal{O}_Y \xrightarrow{\sim} \pi_*\mathcal{O}_X$ and $R^1\pi_*\mathcal{O}_X = R^2\pi_*\mathcal{O}_X = 0$. Then all $\mathbb{G}_m$ -weights of $H^0(X, \Omega_X^1)$ are strictly positive.*

In other words, if there exists a differential 1-form α which is rescaled by the $\mathbb{G}_m$ -action, $t \circ \alpha = t^k \alpha$ for some integer k , then $k > 0$. The proof of Theorem 6.0.1 is quite interesting and uses both reductions to characteristic p and analytification of X over $\mathbb{C}$. We first prove, using the surjectivity of Cartier operator (Lemma 3.2.2), that for big enough p , for the reductions which are resolutions with conical slices, all weights of differential 1-forms are non-negative and that all $\mathbb{G}_m$ -invariant 1-forms are closed (Lemma 6.0.2). Then we use analytification and Dolbeault cohomology to show that there are no non-zero closed $\mathbb{G}_m$ -invariant 1-forms over $\mathbb{C}$ (Lemma 6.0.4). We state Conjecture 6.0.1 that the statement of Corollary 6.0.1 remains true even in characteristic p . In Sect. 6.1 we play around a little bit and try to cope with the potential existence of $\mathbb{G}_m$ -invariant forms in characteristic p . Namely we give a somewhat complicated definition of totally positive forms (Definition 6.1.2), which morally are differential 1-forms on X that have positive weight at a conical neighbourhood of each point. However there is some ambiguity in the choice of the conical neighbourhood and the actual definition is more complicated. The space $H^0(X, \Omega_X^1)^{\geq 0}$ of totally positive forms is invariant under multiplication by functions and we define the sheaf $\mathcal{M}_{\pi, \text{inv}}$ as the coherent sheaf on Y corresponding to the quotient $H^0(X, \Omega_X^1)/H^0(X, \Omega_X^1)^{\geq 0}$. Despite a messy definition, we are able to prove some statements about it: like Lemma 6.1.1 and Lemma 6.1.5. In Definition 6.1.3 we define a strictly positive resolution with conical slices: namely it is a resolution with conic slices for which $\mathcal{M}_{\pi, \text{inv}} = 0$. Finally we end the section with Proposition 6.2.2, which is an enhancement of Proposition 5.6.3 and states that a general reduction of a resolution with conical slices is strictly positive.

In Sect. 7 we prove our main theorem (Theorem 7.2.2):

Theorem 1.2.3. *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a map of schemes over $S = \operatorname{Spec} R$ of characteristic 0, such that the generic fiber $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a resolution with conical slices. Then there exists an étale open $S' \rightarrow S$ such that for any geometric point $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S'$*

- *the corresponding fiber $\pi_s : X_s \rightarrow Y_s$ is a strictly positive resolution with conical slices,*
- *the classes $[\mathcal{D}_{X_s, \alpha}]$ descend to Y_s for all $\alpha \in H^0(X_s, \Omega_{X_s}^1)$.*

The theorem follows directly from the Proposition 6.2.2 and Proposition 7.2.1, which states that for a strictly positive resolution with conical slices the sheaf $Q_{\pi, N}$ is 0. The proof of Proposition 7.2.1 proceeds by induction on $\dim X$, using Remark 5.5.1, while for the central point it uses the same trick with surjectivity of Cartier operator as in Theorem 6.0.1.

In Sect. 8 we prove that some classes of symplectic resolutions are resolutions with conical slices. For this we need to do some additional work. The existence of formal slices for symplectic resolutions was proved by Kaledin in [Kal2]. However the existence of a contracting $\mathbb{G}_m$ -action on the slice is still a conjecture (Conjecture 1.8 in [Kal1]) and we need étale slices anyways. For quiver varieties in Crawley-Boevey's paper [CrB1] it is proven that at any point the base $N_Q(\lambda, \alpha)$ looks étale locally like the central point of $N_{Q'}(0, \alpha')$ which has a natural contracting $\mathbb{G}_m$ -action. It is remarkable that Crawley-Boevey's result suits perfectly well to the framework of our definition of resolution with conical slices: he does not prove the existence of a slice, but provides an étale equivalence with some other variety which has a contracting $\mathbb{G}_m$ -action. Given a quiver Q with set of vertices I , dimension vector $\alpha \in \mathbb{N}^I$ and a vector $\lambda \in \mathbb{K}^I$, variety $N_Q(\lambda, \alpha)$ is defined as the Hamiltonian reduction of the space of representations of the double quiver $\overline{Q}$ with dimension vector α and with the level of the moment map equal to λ . Geometric invariant theory provides a projective map $\pi_\lambda^\theta : N_Q(\lambda, \alpha)^\theta \rightarrow N_Q(\lambda, \alpha)$ from the GIT quotient $N_Q(\lambda, \alpha)^\theta$. The sufficient conditions for the map π_λ^θ to be a resolution of singularities were given also by Crawley-Boevey and we summarise them in Proposition 8.2.1. Unfortunately étale equivalence in Crawley-Boevey's paper is proved only for bases of resolutions and not for resolutions themselves. We do some additional work to extend the equivalence on this level (see Proposition 8.2.3). Since for $\lambda = 0$ the resolution $\pi_0^\theta : N_Q(0, \alpha)^\theta \rightarrow N_Q(0, \alpha)$ is conical, we obtain Theorem 8.2.4, which says that the resolution π_0^θ is a resolution with conical slices. Returning to Theorem 7.2.2 we get that for a general reduction to characteristic p of the resolution $\pi_0^\theta : N_Q(0, \alpha)^\theta \rightarrow N_Q(0, \alpha)$ all the classes $[\mathcal{D}_{N_Q(0, \alpha)^\theta, \alpha}]$ descend to $N_Q(0, \alpha)$. In particular following Sect. 8.2.5 this is true for Nakajima quiver varieties and this answers the question raised in [BL], which motivated this paper.

In Sect. 8.3 we generalise the result of Crawley-Boevey to an arbitrary Hamiltonian reduction of a vector space. Namely given a symplectic vector representation of a reductive group G in a vector space (V, ω) we can take its GIT Hamiltonian reduction $\mathfrak{M}(G, V)_\lambda^\theta = \mu^{-1}(\lambda) //_\theta G$ and consider a natural map $\pi_\lambda^\theta : \mathfrak{M}(G, V)_\lambda^\theta \rightarrow \mathfrak{M}(G, V)_\lambda^0 = \mu^{-1}(0) // G$. We restrict to the case of $V \oplus V^*$ and a symplectic representation of G induced by some representation $G \rightarrow \operatorname{GL}(V)$. Closely following the argument of Crawley-Boevey, we prove that if π_0^θ is a resolution of singularities and the base is normal, it is a resolution with conical slices (see Theorem 8.3.1). However being more general our context differs in a lot of ways from the Crawley-Boevey's one and we need to change some of the arguments in a significant way. Hypertoric variety is Hamiltonian reduction of a cotangent bundle of a representation of an algebraic torus $\mathbb{G}_m^k$ and in Sect. 8.3.5 we apply Theorem 8.3.1 for this case. Hypertoric varieties can be combinatorially encoded by weighted, cooriented affine hyperplane arrangements $\mathcal{A}$ in some lattice (that depends on the representation). The case $\theta = 0$ corresponds to arrangements where all hyperplanes pass through 0. If such arrangement is unimodular (see Sect. 8.3.5 for definition) it possesses a simplification $\tilde{\mathcal{A}}$, which differs from $\mathcal{A}$ only by the choice of a non-trivial character θ . Hypertoric variety $\mathfrak{M}(\tilde{\mathcal{A}})$ is smooth, and the natural map $\pi_{\mathcal{A}} : \mathfrak{M}(\tilde{\mathcal{A}}) \rightarrow \mathfrak{M}(\mathcal{A})$ is known to be a resolution

of singularities. Applying Theorem 8.3.1 we get that resolutions of singularities of hypertoric varieties are resolutions with conical slices. Finally in Sect. 8.4 we prove the same thing for Slodowy slices in a nilpotent cone of a semisimple Lie algebra. Here the proof is different (and much simpler), it relies mostly on Lemma 5.5.1. In all these cases we obtain that for general reduction to characteristic p the classes $[\mathcal{D}_{X,\alpha}]$ descend to the base of the resolution.

1.3. Remaining questions. There are several natural questions which one can ask and which we did not cover in this paper. They remain to be the subject of a future research.

Question 1. Is it true that any conical symplectic resolution is a resolution with conical slices?

This question is very similar to the conjecture due to Kaledin (Conjecture 1.8 in [Kal1]) which claims that the transversal slice to a Poisson leaf has a contracting $\mathbb{G}_m$ -action. Though Kaledin's conjecture is about the formal slice, may be using some finiteness results recently proved by Namikawa (see [Na2]) one can obtain étale slices as well. Anyways it is not yet clear at all how to attack any of these two statements.

Question 2 For a conical symplectic resolution $\pi : X \rightarrow Y$ is it true that the class $[\mathcal{A}_\lambda]$ in $\text{Br}(X^{(1)})$ comes from a differential 1-form (meaning $[\mathcal{A}_\lambda] = [\mathcal{D}_{X,\alpha}]$ for some α)?

One even has a candidate for α — it should be the contraction of the symplectic form ω with the Euler vector field corresponding to the contracting $\mathbb{G}_m$ -action. This is a computation which still needs to be done in the case of general resolution $\pi : X \rightarrow Y$.

Question 3 Is Conjecture 6.0.1 true?

This is just an interesting question about possible pathologies in characteristic p . It also seems to be related to the recent results of [BMS] (see Remark 6.0.3).

1.4. Acknowledgements. We are grateful to Roman Bezrukavnikov for introducing this problem to us, as well as to Michael Finkelberg for his incessant interest in our work. We also would like to thank Bhargav Bhatt, Brian Conrad, Ofer Gabber, Dennis Gaitsgory, Victor Ginzburg, Borys Kadets, Dmitry Kaledin, Ivan Losev, Michael McBreen, Alexey Pakharev, Grigory Papayanov, Nicholas Proudfoot, Sam Raskin, Nick Rosenblum, Konstantin Tolmachov, Misha Verbitsky, Jacqueline Wang and Ben Webster, for all discussions and e-mails which in different ways helped us to finish this paper.

2. FROBENIUS TWIST AND DIFFERENTIAL OPERATORS

2.1. Frobenius twist. Let S be an arbitrary scheme over $\mathbb{F}_p$. The *absolute Frobenius morphism* $\text{Frob}_S : S \rightarrow S$ is given by

- id_S on underlying topological space,
- $\text{Frob}_S^\sharp(f) = f^p$ for $f \in \mathcal{O}_S$ on the level of structure sheaves.

For any S -scheme $\xi : X \rightarrow S$ the *relative Frobenius twist* $\xi^{(S)} : X^{(S)} \rightarrow S$ is defined as the pull-back of ξ with respect to Frob_S . Let $W_{X/S} : X^{(S)} \rightarrow X$ be the natural morphism in the corresponding Cartesian square. Absolute Frobenius is functorial on X , namely $\xi \circ \text{Frob}_X = \text{Frob}_S \circ \xi$ and by the universal property of the pull-back we obtain a unique decomposition $\text{Frob}_X = \text{Fr}_{X/S} \circ W_{X/S}$:

$$\begin{array}{ccccc}
 & & \text{Frob}_X & & \\
 & \curvearrowright & & \curvearrowright & \\
 X & \xrightarrow{\quad} & X^{(S)} & \xrightarrow{\quad} & X \\
 \searrow \xi & \text{Fr}_{X/S} \downarrow & \downarrow \xi^{(S)} & W_{X/S} \downarrow & \downarrow \xi \\
 & & S & \xrightarrow{\quad \text{Frob}_S \quad} & S
 \end{array}$$

Morphism $\mathrm{Fr}_{X/S} : X \rightarrow X^{(S)}$ is called the *relative Frobenius morphism*. We denote by $\bullet^{(S)}$ the pull-back of any object $\bullet$ on X to $X^{(S)}$ under $W_{X/S}$.

In this paper we consider only the case $S = \mathrm{Spec} \mathbb{k}$, where $\mathbb{k}$ is an algebraically closed field of characteristic p . In this case $\mathrm{Frob}_{\mathrm{Spec} \mathbb{k}}$ is an isomorphism (of abstract schemes) because $\mathbb{k}$ is algebraically closed and the map $x \mapsto x^p$ is a bijection. For $S = \mathrm{Spec} \mathbb{k}$ we denote $\mathrm{Fr}_{X/S}$, $W_{X/S}$ and $\bullet^{(S)}$ simply by Fr_X , W_X and $\bullet^{(1)}$. One can inductively define $\bullet^{(k)}$ by $(\bullet^{(k-1)})^{(1)}$. Note that W_X is an isomorphism of abstract schemes too.

For a smooth $\mathbb{k}$ -scheme X we denote $\Omega_{X/\mathrm{Spec} \mathbb{k}}^i$ simply by Ω_X^i . The morphism $\xi : X \rightarrow \mathrm{Spec} \mathbb{k}$ is automatically flat since $\mathbb{k}$ is a field, and so by base change the natural maps

$$\mathcal{T}_{X^{(1)}} \xrightarrow{\sim} W_X^* \mathcal{T}_X = \mathcal{T}_X^{(1)} \text{ and } (\Omega_X^i)^{(1)} = W_X^* \Omega_X^i \xrightarrow{\sim} \Omega_{X^{(1)}}^i$$

are isomorphisms. For a vector space V over $\mathbb{k}$ we denote by $V^{(1)}$ the vector space V with a twisted $\mathbb{k}$ -structure:

$$\lambda^{(1)} \cdot v = \lambda^{\frac{1}{p}} \cdot v \text{ for } a \in \mathbb{k} \text{ and } v \in V.$$

Note that $V^{(1)}$ is nothing but $\mathrm{Frob}_{\mathrm{Spec} \mathbb{k}}^* V$ if we consider V as a quasi-coherent sheaf on $\mathrm{Spec} \mathbb{k}$. For every vector space V we have the corresponding affine space $\mathbb{A}(V)$, considered as a group scheme with an additional action of $\mathbb{k}$. Then on the level of $\mathbb{k}$ -points two twists coincide:

$$\mathbb{A}(V)^{(1)}(\mathbb{k}) = V^{(1)}.$$

Returning to the commutative diagram, applying the base change once again and using one of the isomorphisms above, we obtain natural isomorphisms

$$H^0(X, \Omega_X^i)^{(1)} \simeq H^0\left(X^{(1)}, (\Omega_X^i)^{(1)}\right) \simeq H^0(X^{(1)}, \Omega_{X^{(1)}}^i).$$

This way to any differential i -form $\omega \in H^0(X, \Omega_X^i)$ on X one can associate a differential form $\omega^{(1)} \in H^0(X^{(1)}, \Omega_{X^{(1)}}^i)$ on $X^{(1)}$. The map $\omega \mapsto \omega^{(1)}$ is by definition $\mathrm{Frob}_{\mathbb{k}}$ -linear.

Note that following the definitions the subsheaf $\mathcal{O}_{X^{(1)}} \subset (\mathrm{Fr}_X)_* \mathcal{O}_X$ can be identified with the subsheaf $(\mathrm{Fr}_X)_*(\mathcal{O}_X^p) \subset (\mathrm{Fr}_X)_* \mathcal{O}_X$ of p -th powers of functions on X .

2.2. Cartier operator. Let $\Omega_X^\bullet$ be the algebraic de Rham complex of X . The differential of $(\mathrm{Fr}_X)_* \Omega_X^\bullet$ is $\mathcal{O}_{X^{(1)}}$ -linear: $d(f^p \omega) = f^p d\omega - p f^{p-1} \cdot df \wedge \omega = f^p d\omega$, so $(\mathrm{Fr}_X)_* \Omega_X^\bullet$ defines a coherent sheaf of DG-algebras on $X^{(1)}$. In particular, cocycle, coboundary and cohomology sheaves of $(\mathrm{Fr}_X)_* \Omega_X^\bullet$ have natural structures of coherent sheaves on $X^{(1)}$. By definition the i -th cocycle sheaf $Z^i((\mathrm{Fr}_X)_* \Omega_X^\bullet)$ is identified with $(\mathrm{Fr}_X)_* \Omega_{X,cl}^i$, where $\Omega_{X,cl}^i$ is the sheaf of closed differential forms of degree i on X . The cohomology sheaves $\bigoplus_i \mathcal{H}^i((\mathrm{Fr}_X)_* \Omega_X^\bullet)$ form a coherent sheaf of graded modules and DG-algebra structure on $(\mathrm{Fr}_X)_* \Omega_X^\bullet$ turns it into a coherent sheaf of graded algebras on $X^{(1)}$.

Theorem 2.2.1 (Cartier isomorphism, [Katz2], Theorem 7.2). *Let X be a smooth variety over an algebraically closed field $\mathbb{k}$ of characteristic p . Then there is a unique isomorphism of coherent sheaves of graded algebras on $X^{(1)}$:*

$$\bigoplus_i C_i^{-1} : \bigoplus_i \Omega_{X^{(1)}}^i \xrightarrow{\sim} \bigoplus_i \mathcal{H}^i((\mathrm{Fr}_X)_* \Omega_X^\bullet),$$

such that $C_0^{-1}(1) = 1$ and $C_1^{-1}(dg^{(1)}) = [g^{p-1} dg]$ for $g \in \mathcal{O}_X$.

Cartier isomorphism also produces for us an isomorphism $C_1 : \mathcal{H}^1((\mathrm{Fr}_X)_* \Omega_X^\bullet) \xrightarrow{\sim} \Omega_{X^{(1)}}^1$ in the other direction. We define the Cartier operator $C : (\mathrm{Fr}_X)_* \Omega_{X,cl}^1 \rightarrow \Omega_{X^{(1)}}^1$ as the composition of the natural projection $(\mathrm{Fr}_X)_* \Omega_{X,cl}^1 \rightarrow \mathcal{H}^1((\mathrm{Fr}_X)_* \Omega_X^\bullet)$ and C_1 . It is easy to see that $\mathrm{Ker} C$

is exactly the subsheaf of exact 1-forms and so is isomorphic to the image of $d : (\mathrm{Fr}_X)_* \mathcal{O}_X \rightarrow (\mathrm{Fr}_X)_* \Omega_{X,cl}^1$. Finally $\mathrm{Ker} d = \mathcal{O}_{X^{(1)}}$ and as a result we obtain the following short exact sequence:

$$0 \rightarrow (\mathrm{Fr}_X)_* \mathcal{O}_X / \mathcal{O}_{X^{(1)}} \xrightarrow{d} (\mathrm{Fr}_X)_* \Omega_{X,cl}^1 \xrightarrow{C} \Omega_{X^{(1)}}^1 \rightarrow 0.$$

Remark 2.2.1. Further we will sometimes omit $(\mathrm{Fr}_X)_*$ before Ω_X^1 , $\Omega_{X,cl}^1$, $\mathcal{O}_X$ and other quasi-coherent sheaves on X and will automatically consider them as sheaves on $X^{(1)}$. For example in this convention the short exact sequence above becomes even shorter and looks like this:

$$0 \rightarrow \mathcal{O}_X / \mathcal{O}_{X^{(1)}} \xrightarrow{d} \Omega_{X,cl}^1 \xrightarrow{C} \Omega_{X^{(1)}}^1 \rightarrow 0.$$

We tried our best to avoid a possible confusion specifying whether the sheaf is on X or $X^{(1)}$ where necessary.

2.3. Crystalline differential operators. Let X be a smooth scheme over an algebraically closed field $\mathbb{k}$ of characteristic p . The sheaf of algebras of “crystalline differential operators” on X is defined as the universal enveloping algebra $\mathcal{D}_X$ of the Lie algebroid $\mathcal{T}_X$ of vector fields on X . In other words, $\mathcal{D}_X$ is a sheaf of associative algebras containing $\mathcal{O}_X$, such that for a quasi-coherent $\mathcal{O}_X$ -module $\mathcal{F}$, an extension of the $\mathcal{O}_X$ -action on $\mathcal{F}$ to a $\mathcal{D}_X$ -action is equivalent to endowing $\mathcal{F}$ with a flat connection. More explicitly, $\mathcal{D}_X$ is a sheaf of algebras generated by the algebra of functions $\mathcal{O}_X$ and the $\mathcal{O}_X$ -module of vector fields $\mathcal{T}_X$, subject to the module and commutator relations $f \cdot \partial = f\partial$, $\partial \cdot f - f \cdot \partial = \partial(f)$ for $\partial \in \mathcal{T}_X, f \in \mathcal{O}_X$, and the Lie algebroid relations $\partial' \cdot \partial'' - \partial'' \cdot \partial' = [\partial', \partial'']$ for $\partial', \partial'' \in \mathcal{T}_X$. If we fix a local frame $\{\partial_1, \dots, \partial_n\}$ of vector fields on X , we obtain a decomposition $\mathcal{D}_X = \bigoplus_{I \in \mathbb{N}^n} \mathcal{O}_X \cdot \partial^I$ as a left $\mathcal{O}_X$ -module. For example for $X = \mathbb{A}^n$, the global sections of $\mathcal{D}_X$ are given by the Weyl algebra $W_n = \mathbb{k}[x_1, \dots, x_n, \partial_1, \dots, \partial_n] / \langle [\partial_i, x_i] = 1, [\partial_i, x_j]_{i \neq j} = [x_i, x_j] = [\partial_i, \partial_j] = 0 \rangle$.

Sheaf $\mathcal{O}_X$ is naturally a sheaf of left modules over $\mathcal{D}_X$. The corresponding action is defined by the multiplication for functions and by differentiation for vector fields: $g \circ f = gf$ and $\partial \circ f = \partial(f)$. It is easy to see that this defines a homomorphism $\theta : \mathcal{D}_X \rightarrow \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{O}_X)$ to the sheaf of algebras of $\mathbb{k}$ -linear endomorphisms of $\mathcal{O}_X$. While in the case of $\mathrm{char} \mathbb{k} = 0$ this map is an embedding, it turns out that for $\mathrm{char} \mathbb{k} = p$ this map has a huge kernel and consequently has a relatively small (more precisely $\mathcal{O}_X$ -coherent) image.

By the binomial formula, given a vector field $\partial \in \mathcal{T}_X$, the differential operator $\partial^p \in \mathcal{D}_X$ acts on functions as a derivation:

$$\partial^p(f \cdot g) = \sum_{i=0}^p \binom{p}{i} \partial^i(f) \cdot \partial^{p-i}(g) = \partial^p(f) \cdot g + f \cdot \partial^p(g).$$

This way it defines another vector field $\partial^{[p]} \in \mathcal{T}_X$, which is called the p -th restricted power of ∂ . For $\partial \in \mathcal{T}_X$ we set $\iota(\partial) := \partial^p - \partial^{[p]} \in \mathcal{D}_X$. By the definition of $\bullet^{[p]}$, the map ι lands in the kernel of the $\mathbb{k}$ -linear action $\theta : \mathcal{D}_X \rightarrow \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{O}_X)$.

Not very difficult computation shows that $\iota(f\partial) = f^p \iota(\partial)$ (e.g. see [Katz2], Proposition 5.3 or a simpler argument in [BMR], Lemma 1.3.1). So it defines a map $\iota : \mathcal{T}_{X^{(1)}} \rightarrow (\mathrm{Fr}_X)_* \mathcal{D}_X$ of quasi-coherent sheaves on $X^{(1)}$ ([BMR], Lemma 1.3.1): namely $\iota(\partial^{(1)}) = \partial^p - \partial^{[p]}$. Moreover one can show that $\mathrm{Ker} \theta$ is given exactly by the two-sided ideal $\mathcal{D}_X \cdot \iota(\mathcal{T}_X) \cdot \mathcal{D}_X$.

Let now $T^*X^{(1)}$ be the total space of the cotangent bundle of $X^{(1)}$ and let $q : T^*X^{(1)} \rightarrow X^{(1)}$ be the natural projection. Then $q_* \mathcal{O}_{T^*X^{(1)}}$ is naturally a quasi-coherent sheaf of commutative algebras on $X^{(1)}$. Note that since the map q is affine, the functor q_* induces an equivalence between the category of quasi-coherent sheaves of $q_* \mathcal{O}_{T^*X^{(1)}}$ -modules on $X^{(1)}$ and the category of quasi-coherent sheaves on $T^*X^{(1)}$. The natural pairing $\langle \cdot, \cdot \rangle : \mathcal{T}_{X^{(1)}} \otimes \Omega_{X^{(1)}}^1 \rightarrow \mathcal{O}_{X^{(1)}}$ defines

an embedding $\mathcal{T}_{X^{(1)}} \hookrightarrow q_*\mathcal{O}_{T^*X^{(1)}}$, which extends to a natural isomorphism $\mathrm{Sym}_{X^{(1)}}^\bullet \mathcal{T}_{X^{(1)}} \simeq q_*\mathcal{O}_{T^*X^{(1)}}$.

Proposition 2.3.1 ([BMR], Lemma 1.3.2, Proposition 2.3.3). *The map $\iota : \mathcal{T}_{X^{(1)}} \rightarrow \mathcal{D}_X$ lands in the center $Z(\mathcal{D}_X) \subset \mathcal{D}_X$. Moreover, its extension to $\mathrm{Sym}_{X^{(1)}}^\bullet \mathcal{T}_{X^{(1)}}$ provides an isomorphism of the center $Z(\mathcal{D}_X)$ and $q_*\mathcal{O}_{T^*X^{(1)}}$. So $\mathcal{D}_X$ is a $q_*\mathcal{O}_{T^*X^{(1)}}$ -central quasi-coherent sheaf of algebras on $X^{(1)}$ and in this way defines a quasi-coherent sheaf of algebras on $T^*X^{(1)}$. As a sheaf of algebras on $T^*X^{(1)}$, $\mathcal{D}_X$ is an Azumaya algebra. It has rank $p^{2 \dim X}$ and is non-trivial if $\dim X > 0$.*

2.4. Azumaya algebras $\mathcal{D}_{X,\alpha}$. According to the Proposition 2.3.1 there is a sheaf $\mathcal{D}_X$ of Azumaya algebras on $T^*X^{(1)}$. Let $\alpha \in H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ be a 1-form on $X^{(1)}$. Its graph $i_\alpha : \Gamma_\alpha \hookrightarrow T^*X^{(1)}$ is by definition the section given by α and considered as a subvariety of $T^*X^{(1)}$. Projection to $X^{(1)}$ provides an isomorphism $q : \Gamma_\alpha \simeq X^{(1)}$.

Definition 2.4.1. The Azumaya algebra $\mathcal{D}_{X,\alpha}$ on $X^{(1)}$ is defined as the restriction $i_\alpha^* \mathcal{D}_X$ of $\mathcal{D}_X$ to $\Gamma_\alpha \simeq X^{(1)}$.

This restriction can be described in more or less explicit way. The graph $\Gamma_\alpha \subset T^*X^{(1)}$ is defined by the system of equations $\partial^{(1)} - \langle \partial^{(1)}, \alpha \rangle = 0$ for all $\partial^{(1)} \in \mathcal{T}_{X^{(1)}}$, where the brackets denote the natural pairing $\langle \cdot, \cdot \rangle : \mathcal{T}_{X^{(1)}} \otimes \Omega_{X^{(1)}}^1 \rightarrow \mathcal{O}_{X^{(1)}}$. Remember that we have an isomorphism $\iota : q_*\mathcal{O}_{T^*X^{(1)}} \simeq Z(\mathcal{D}_X)$. The restriction of $\mathcal{D}_X$ to Γ_α as an Azumaya algebra is obtained by taking the quotient of $\mathcal{D}_X$ by the ideal generated by $\iota(\partial - \langle \partial^{(1)}, \alpha \rangle)$. In other words $\mathcal{D}_{X,\alpha} = \mathcal{D}_X / \langle \partial^p - \partial^{[p]} - \langle \partial, \alpha \rangle \rangle_{\partial \in \mathcal{T}_{X^{(1)}}}$, where $\langle \partial^{(1)}, \alpha \rangle$ lies in $\mathcal{O}_{X^{(1)}} = (\mathcal{O}_X)^p \subset \mathcal{O}_X \subset \mathcal{D}_X$.

Note that the association $\alpha \mapsto \mathcal{D}_{X,\alpha}$ defines a map $c_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \mathrm{Br}(X^{(1)})$ (for now just a map of sets).

2.5. Azumaya algebras $\mathcal{D}_{X,\alpha}$ and p -curvature. As we have seen, sheaves of algebras $\mathcal{D}_{X,\alpha}$ are some strange objects which have differential-geometric nature, but exist exclusively in characteristic p . There is another object of this sort, namely the p -curvature of a flat connection on X . In this subsection we are going to show how they are related.

Let $\mathcal{E}$ be a quasi-coherent sheaf on X and let $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_X^1$ be a connection on $\mathcal{E}$, which by definition means that locally $\nabla(fs) = s \otimes df + f \nabla(s)$ for any section $s \in \mathcal{E}(U)$ and function $f \in \mathcal{O}_X(U)$. Note that ∇ is not $\mathcal{O}_X$ -linear, but the difference of two connections $\nabla_1 - \nabla_2$ is, so connections on $\mathcal{E}$ form a torsor over $\mathcal{O}_X$ -linear maps from $\mathcal{E}$ to $\mathcal{E} \otimes \Omega_X^1$. Dually, any ∇ defines an $\mathcal{O}_X$ -linear map $\nabla : \mathcal{T}_X \rightarrow \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ and $\nabla_1 - \nabla_2$ lands in $\underline{\mathrm{End}}_{\mathcal{O}_X}(\mathcal{E})$ for any ∇_1, ∇_2 .

Sheaves $\mathcal{T}_X$ and $\underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ have natural structures of sheaves of Lie algebras of $\mathbb{k}$ -vector spaces. They are given by commutator of vector fields for $\mathcal{T}_X$ and the commutator $[a, b] = ab - ba$ for $\underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$. The classical curvature $\mathrm{curv}(\nabla) : \bigwedge_{\mathcal{O}_X}^2 \mathcal{T}_X \rightarrow \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ measures how far ∇ is from being a homomorphism of Lie algebras, namely

$$\mathrm{curv}(\nabla)(\partial_1, \partial_2) = [\nabla(\partial_1), \nabla(\partial_2)] - \nabla([\partial_1, \partial_2]).$$

A connection ∇ is called *integrable* or *flat* if the map $\mathrm{curv}(\nabla)$ is identically equal to 0. In this case $\nabla : \mathcal{T}_X \rightarrow \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ is a Lie algebra homomorphism.

In characteristic p both $\mathcal{T}_X$ and $\underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ possess a richer structure of p -th restricted Lie algebra, given by $\bullet^{[p]}$ and $\bullet^p$ correspondingly. Analogously to the classical case, the p -curvature map $\mathrm{curv}_p(\nabla) : \mathcal{T}_{X^{(1)}} \rightarrow (\mathrm{Fr}_X)_* \underline{\mathrm{End}}_{\mathbb{k}}(\mathcal{E})$ measures how much ∇ fails to preserve these new operations. It is defined by the following formula:

$$\mathrm{curv}_p(\nabla)(\partial^{(1)}) = \nabla(\partial)^p - \nabla(\partial^{[p]}).$$

By the same computation as before ([Katz2], Proposition 5.3) $\text{curv}_p(\nabla)$ is p -linear: $\text{curv}_p(\nabla)(f^{(1)}\partial^{(1)}) = f^p \text{curv}_p(\nabla)(\partial^{(1)})$. This explains the $\bullet^{(1)}$ and $(\text{Fr}_X)_*$ in the source and the target of the corresponding map.

Another feature of finite characteristic is that any connection $\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes \Omega_X^1$ is $\mathcal{O}_{X^{(1)}}$ -linear since $d(f^p) = 0$. So $\nabla : \mathcal{T}_X \rightarrow \underline{\text{End}}_{\mathbb{k}}(\mathcal{E})$ actually lands in $\mathcal{O}_{X^{(1)}}$ -linear endomorphisms $\underline{\text{End}}_{\mathcal{O}_{X^{(1)}}}((\text{Fr}_X)_*\mathcal{E})$. It follows immediately that p -curvature lands in $\underline{\text{End}}_{\mathcal{O}_{X^{(1)}}}((\text{Fr}_X)_*\mathcal{E})$ too. However it comes out that $\text{curv}_p(\nabla)(\partial^{(1)})$ is always $\mathcal{O}_X$ -linear:

Lemma 2.5.1. *p -curvature $\text{curv}_p(\nabla)$ lands in $\mathcal{O}_X$ -linear endomorphisms of $\mathcal{E}$. This way it defines a map $\text{curv}_p(\nabla) : \mathcal{T}_X^{(1)} \rightarrow (\text{Fr}_X)_*\underline{\text{End}}_{\mathcal{O}_X}(\mathcal{E})$.*

Proof. By the Leibniz rule,

$$\nabla(\partial)^p(f \cdot e) = \partial^p(f) \cdot e + f \cdot \nabla(\partial)^p(e).$$

On the other hand,

$$\nabla(\partial^{[p]})(f \cdot e) = \partial^{[p]}(f) \cdot e + f \cdot \nabla(\partial^{[p]})(e).$$

Now, since $\partial^p(f) = \partial^{[p]}(f)$, subtracting second equality from the first one, we get that

$$\nabla(\partial)^p(f \cdot e) - \nabla(\partial^{[p]})(f \cdot e) = f \cdot (\nabla(\partial)^p(e) - \nabla(\partial^{[p]})(e))$$

or, in other words,

$$\text{curv}_p(\nabla)(\partial^{(1)})(f \cdot e) = f \cdot \text{curv}_p(\nabla)(\partial^{(1)})(e).$$

□

Remark 2.5.1. In particular, if $\mathcal{E}$ is a line bundle, $\underline{\text{End}}_{\mathcal{O}_X}(\mathcal{E}) = \mathcal{O}_X$ and $\text{curv}_p(\nabla)$ defines a map $\text{curv}_p(\nabla) : \mathcal{T}_{X^{(1)}} \rightarrow (\text{Fr}_X)_*\mathcal{O}_X$. Moreover, one can show (see Proposition 5.2, [Katz2]) that p -curvature is not only $\mathcal{O}_X$ -linear, but also flat with respect to the induced connection ∇_{End} on $\underline{\text{End}}_{\mathcal{O}_X}(\mathcal{E})$. In the case when $\mathcal{E}$ is a line bundle this connection is trivial, namely $\nabla_{\text{End}}(\partial) = \theta(\partial)$, where $\theta(\partial) : \mathcal{O}_X \rightarrow \mathcal{O}_X$ is the derivation corresponding to ∂ . The flat sections are given by $\mathcal{O}_{X^{(1)}} \subset \mathcal{O}_X$ and so p -curvature defines a map $\text{curv}_p(\nabla) : \mathcal{T}_{X^{(1)}} \rightarrow \mathcal{O}_{X^{(1)}}$. Such a map is the same as a differential 1-form on $X^{(1)}$ which in this case we will call $\text{curv}_p(\nabla)$ as well.

Let $\text{MIC}(X)$ be the category of pairs $(\mathcal{E}, \nabla)$, where $\mathcal{E}$ is a quasi-coherent sheaf on X and ∇ is an integrable connection on $\mathcal{E}$. ∇ defines a natural $\mathcal{D}_X$ -action on $\mathcal{E}$, namely functions act by multiplication and vector fields act through ∇ : $\partial \mapsto \nabla(\partial) \in \text{End}_{\mathcal{O}_{X^{(1)}}}(\mathcal{E}) \subset \text{End}_{\mathbb{k}}(\mathcal{E})$. The integrability of the connection guarantees that this action is well-defined. As was mentioned before, the converse is also true: an action of $\mathcal{D}_X$ which extends the $\mathcal{O}_X$ -module structure on $\mathcal{E}$ endows $\mathcal{E}$ with a flat connection ($\nabla(\partial)$ acts as ∂). In other words we have an equivalence of categories between $\text{MIC}(X)$ and the category of $\mathcal{O}_X$ -quasi-coherent $\mathcal{D}_X$ -modules. It is easy to see from this point of view that $\text{curv}_p(\nabla)(\partial^{(1)})$ is just equal to $\nabla(\iota(\partial)) = \nabla(\partial^p - \partial^{[p]})$, where $\iota(\partial) = \partial^p - \partial^{[p]} \in \mathcal{D}_X$.

The p -curvature appears to be an important tool even while studying differential equations in characteristic 0, with the characteristic p proof of the quasi-unipotence of Gauss-Manin connection ([Katz2]) and Grothendieck-Katz conjecture ([Katz1], [FK]) being probably the best illustrating examples.

A much simpler result, now related to algebraic geometry in characteristic p , is the Cartier theorem, which relates p -curvature to the descent data for the relative Frobenius map:

Theorem 2.5.1 (Cartier, [Katz2], Theorem 5.1). *Let X be a smooth scheme of finite type over $\mathbb{k}$. Let $\text{MIC}(X)$ denote the category of quasi-coherent sheaves with integrable connection $(\mathcal{E}, \nabla)$.*

Remember also that we have the relative Frobenius morphism $\mathrm{Fr}_X : X \rightarrow X^{(1)}$ to the Frobenius twist of X . There is an equivalence of categories between the category of quasi-coherent sheaves on $X^{(1)}$ and the subcategory of $\mathrm{MIC}(X)$ consisting of pairs $(\mathcal{E}, \nabla)$, such that $\mathrm{curv}_p(\nabla)$ is equal to 0.

The natural question now is: what do non-zero p -curvatures correspond to? The algebras $\mathcal{D}_{X,\alpha}$ have to do with a partial answer to this question.

A differential 1-form $\alpha \in \Omega_{X^{(1)}}^1$ gives a map $\tilde{\alpha} : \mathcal{T}_X^{(1)} \rightarrow \mathcal{O}_{X^{(1)}}$. For any quasi-coherent sheaf $\mathcal{E}$ on X we have a natural map $\mathcal{O}_{X^{(1)}} \rightarrow (\mathrm{Fr}_X)_* \mathcal{O}_X \rightarrow (\mathrm{Fr}_X)_* \underline{\mathrm{End}}_{\mathcal{O}_X}(\mathcal{E})$. Taking the composition, we obtain a well-defined map $\tilde{\alpha} : \mathcal{T}_X^{(1)} \rightarrow (\mathrm{Fr}_X)_* \underline{\mathrm{End}}_{\mathcal{O}_X}(\mathcal{E})$ for any quasi-coherent sheaf $\mathcal{E}$. Let's denote by $\mathrm{MIC}_{\tilde{\alpha}}(X)$ the full subcategory of $\mathrm{MIC}(X)$ consisting of pairs $(\mathcal{E}, \nabla)$, such that $\mathrm{curv}_p(\nabla) = \tilde{\alpha}$. The following proposition is essentially a tautology and gives a partial answer to the question above.

Proposition 2.5.1. *There is an equivalence of categories between the category $\mathrm{MIC}_{\tilde{\alpha}}(X)$ and the category of $\mathcal{D}_{X,\alpha}$ -modules that are $\mathcal{O}_X$ -quasi-coherent.*

Proof. Recall the presentation of $\mathcal{D}_{X,\alpha}$ as a quotient of $\mathcal{D}_X$. Thus $\mathcal{D}_{X,\alpha}$ -modules are $\mathcal{D}_X$ -modules for which the action map factorises through $\mathcal{D}_{X,\alpha}$. So it is enough to check that $(\mathcal{E}, \nabla) \in \mathrm{MIC}_{\tilde{\alpha}}(X)$ if and only if $\partial^p - \partial^{[p]} - \langle \partial^{(1)}, \alpha \rangle$ acts as 0 (under the $\mathcal{D}_X$ -action on $\mathcal{E}$ defined by ∇) for any $\partial \in \mathcal{T}_X$. But

$$\nabla(\partial^p - \partial^{[p]} - \langle \partial^{(1)}, \alpha \rangle) = \mathrm{curv}_p(\nabla)(\partial^{(1)}) - \langle \partial^{(1)}, \alpha \rangle.$$

This expression is equal to 0 for any ∂ if and only if $\mathrm{curv}_p(\nabla) = \tilde{\alpha}$. Note that here again we have $\langle \partial^{(1)}, \alpha \rangle \in \mathcal{O}_{X^{(1)}} = (\mathcal{O}_X)^p \subset \mathcal{O}_X$. $\square$

This proposition can be considered as a direct generalisation of Cartier's theorem and we will see why very soon (in Remark 2.5.2).

Let $(\mathcal{E}, \nabla)$ be a line bundle with a flat connection. In local coordinates $\mathcal{E} \simeq \mathcal{O}_X$ and a connection ∇ is given by a closed 1-form ω , namely $\nabla = d + \omega$. One can also show that the explicit formula for p -curvature is then given by $\mathrm{curv}_p(d + \omega) = \omega^{(1)} - C(\omega)$. This relates p -curvature to Cartier morphism.

Let's moreover fix $\alpha \in \Omega_{X^{(1)}}^1$ and assume that $\mathrm{curv}_p(\nabla)$ is equal to α . By Proposition 2.5.1 we have a $\mathcal{D}_{X,\alpha}$ -action on $\mathcal{E}$, or, in other words, a homomorphism $\theta_{\nabla} : \mathcal{D}_{X,\alpha} \rightarrow \mathrm{End}_{\mathcal{O}_{X^{(1)}}}((\mathrm{Fr}_X)_* \mathcal{E})$. By Proposition 2.3.1, $\mathcal{D}_{X,\alpha}$ has rank $p^{2 \dim X}$ over $\mathcal{O}_{X^{(1)}}$. Moreover, an easy computation in local coordinates shows that the fiber of $\mathcal{D}_{X,\alpha}$ at each point is a matrix algebra of size $p^{\dim X}$ ([BMR], Lemma 2.2.1). Bundle $(\mathrm{Fr}_X)_* \mathcal{E}$ has rank $p^{\dim X}$, so $\mathrm{End}_{\mathcal{O}_{X^{(1)}}}((\mathrm{Fr}_X)_* \mathcal{E})$ has rank $p^{2 \dim X}$ and the fiber at each point is a matrix algebra of the same size. Map θ_{∇} is an isomorphism if and only if it is an isomorphism on each fiber, which in our case is equivalent to being non-zero, since matrix algebra is simple. But $\mathcal{O}_X \subset \mathcal{D}_{X,\alpha}$ always acts by multiplication on functions, so for any point this map can not be identically zero. We obtain an isomorphism $\theta_{\nabla} : \mathcal{D}_{X,\alpha} \rightarrow \mathrm{End}_{\mathcal{O}_{X^{(1)}}}((\mathrm{Fr}_X)_* \mathcal{E})$, or in other words we get that $(\mathrm{Fr}_X)_* \mathcal{E}$ is a splitting bundle for Azumaya algebra $\mathcal{D}_{X,\alpha}$. On the other hand, if we have a splitting $\theta : \mathcal{D}_{X,\alpha} \rightarrow \mathrm{End}_{\mathcal{O}_{X^{(1)}}}(\mathcal{E}')$, we get an embedding $(\mathrm{Fr}_X)_* \mathcal{O}_X \hookrightarrow \mathrm{End}_{\mathcal{O}_{X^{(1)}}}(\mathcal{E}')$, which shows that $\mathcal{E}'$ is in fact of the form $(\mathrm{Fr}_X)_* \mathcal{E}$ for some line bundle $\mathcal{E}$ on X (namely $\mathcal{E}$ is just $\mathcal{E}'$ considered as a module over $\mathcal{O}_X$). The restriction of θ to $(\mathrm{Fr}_X)_* \mathcal{T}_X$ provides $\mathcal{E}$ with an integrable connection with the p -curvature α .

To summarise the discussion let us introduce two categories. Let $\mathrm{Spl}(X, \alpha)$ be the category of splittings of $\mathcal{D}_{X,\alpha}$, namely by definition

- objects of $\text{Spl}(X, \alpha)$ are pairs $(\mathcal{E}', \theta)$, consisting of a vector bundle $\mathcal{E}'$ on $X^{(1)}$ and an isomorphism $\theta : \mathcal{D}_{X,\alpha} \rightarrow \text{End}_{\mathcal{O}_{X^{(1)}}}(\mathcal{E}')$,
- morphisms in $\text{Spl}(X, \alpha)$ are usual morphisms between vector bundles that respect the splittings. Note that such a morphism is necessarily an isomorphism.

The second category $LIC(X, \alpha)$ is the category of line bundles with an integrable connection with p -curvature α , namely

- objects of $LIC(X, \alpha)$ are pairs $(\mathcal{E}, \nabla)$, consisting of a line bundle $\mathcal{E}$ on X and an integrable connection ∇ , such that $\text{curv}_p(\nabla) = \alpha$,
- morphisms in $LIC(X, \alpha)$ are isomorphisms of line bundles which respect the connections.

Summarising the discussion above, we get the following

Proposition 2.5.2. *The category $\text{Spl}(X, \alpha)$ is equivalent to $LIC(X, \alpha)$ for any smooth variety X and any differential 1-form α on $X^{(1)}$.*

Remark 2.5.2. In particular taking $\alpha = 0$ we get that $(\mathcal{O}_X, d)$ produces a splitting bundle for $\mathcal{D}_{X,0}$, namely we have an isomorphism $\theta_d : \mathcal{D}_{X,0} \xrightarrow{\sim} \text{End}_{\mathcal{O}_{X^{(1)}}}((\text{Fr}_X)_* \mathcal{O}_X)$. So we see that $\mathcal{D}_{X,0}$ is canonically split for any X and is Morita-equivalent to $\mathcal{O}_{X^{(1)}}$. This canonical Morita-equivalence is given by tensor product of a quasi-coherent sheaf on $X^{(1)}$ with $(\text{Fr}_X)_* \mathcal{O}_X$, with the $\mathcal{D}_{X,0}$ -module structure given by d .

2.6. The gerbes $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$. Gerbes are 2-categorical analogues of principal G -bundles. They provide a convenient, though may be a little bit too abstract, language to study certain classes of objects which appear quite naturally in different areas of mathematics. $\mathbb{G}_m$ -gerbes play an important role in classical differential geometry, they serve as geometrifications (categorifications) of certain interesting characteristic classes (lying in $H^3(X, \mathbb{Z})$). Gerbes naturally show up in our situation too, as we will see soon.

Definition 2.6.1. A *gerbe* on a scheme X over an algebraically closed field $\mathbb{k}$ is a stack of groupoids $\mathcal{G}$ on étale site $X_{\text{ét}}$ which is

- *locally non-empty*, meaning that any point $x \in X(\mathbb{k})$ admits an étale neighbourhood U , such that $\mathcal{G}(U)$ is not empty;
- *transitive*, meaning that for any $U \in X_{\text{ét}}$ and any pair of objects A_1 and A_2 of $\mathcal{G}(U)$, there exists an étale cover $g : V \rightarrow U$, such that $g^*(A_1) \in \mathcal{G}(V)$ is isomorphic to $g^*(A_2)$.

Given a 1-form α on $X^{(1)}$ to each $U \in X_{\text{ét}}$ we can associate two groupoids: $\text{Spl}(U, \alpha|_U)$ and $LIC(U, \alpha|_U)$. We would like to show that this data glues up into two gerbes: gerbe of splittings of $\mathcal{D}_{X,\alpha}$ and gerbe of line bundles with an integrable connection with the p -curvature α . To do this we will first need to prove couple of lemmas.

Let $f : U \rightarrow X$ be an étale morphism. Since f is étale, it induces an isomorphism $f^* \Omega_X^1 \xrightarrow{\sim} \Omega_U^1$ and (using this isomorphism) produces a map $f : T^*U \rightarrow T^*X$ between total spaces of cotangent bundles. Let $f : T^*U^{(1)} \rightarrow T^*X^{(1)}$ also denote the corresponding map between the Frobenius twists. Recall that we have Azumaya algebras $\mathcal{D}_X$ and $\mathcal{D}_U$ on $T^*X^{(1)}$ and $T^*U^{(1)}$ correspondingly.

Lemma 2.6.1. *Let $f : U \rightarrow X$ be an étale morphism. Then $f^* \mathcal{D}_X$ is canonically isomorphic to $\mathcal{D}_U$.*

Proof. We use the presentation of $\mathcal{D}_U$ as universal enveloping algebras of the Lie algebroid $\mathcal{T}_U$ of vector fields. The statement of the lemma then immediately follows from the isomorphism $\mathcal{T}_U \xrightarrow{\sim} f^* \mathcal{T}_X$ and functoriality of Frobenius. $\square$

Corollary 2.6.1. *Let $f : U \rightarrow X$ be an étale morphism and let α be a differential 1-form on $X^{(1)}$. Then $f^*\mathcal{D}_{X,\alpha}$ is canonically isomorphic to $\mathcal{D}_{U,f^*\alpha}$.*

Proof. This follows from a simple observation that $f^{-1}(\Gamma_\alpha) = \Gamma_{f^*\alpha}$. $\square$

This corollary shows that the association $\alpha \rightarrow \mathcal{D}_{X,\alpha}$ is functorial in X with respect to étale morphisms. Let now $g : V \rightarrow X$ and $f : U \rightarrow X$ be two objects in X_{et} and $h : V \rightarrow U$ be a morphism between them. Such a morphism produces natural functors $\mathcal{S}(h) : \text{Spl}(U, f^*\alpha) \rightarrow \text{Spl}(V, g^*\alpha)$ and $\mathcal{L}(h) : \text{LIC}(U, f^*\alpha) \rightarrow \text{LIC}(V, g^*\alpha)$. Indeed if $(\mathcal{E}, \theta)$ is a splitting bundle for $\mathcal{D}_{U,f^*\alpha}$, then $(h^*\mathcal{E}, h^*\theta)$ is a splitting bundle for $h^*\mathcal{D}_{U,f^*\alpha}$, which by Corollary 2.6.1 is canonically isomorphic to $\mathcal{D}_{V,h^*(f^*\alpha)} = \mathcal{D}_{V,g^*\alpha}$. If now we have an object $(\mathcal{E}, \nabla)$ of $\text{LIC}(U, f^*\alpha)$, then

$$\begin{aligned} \text{curv}_p(h^*\nabla)(\partial) &= (h^*\nabla(\partial))^p - h^*\nabla(\partial^{[p]}) = \nabla(h_*\partial)^p - \nabla(h_*(\partial^{[p]})) = \\ &= \nabla(h_*\partial)^p - \nabla((h_*\partial)^{[p]}) = \text{curv}_p(\nabla)(h_*\partial) = f^*\alpha(h_*\partial) = h^*(f^*\alpha)(\partial) = g^*\alpha(\partial), \end{aligned}$$

so $\text{curv}_p(h^*\nabla) = h^*\alpha$ and $(h^*(\mathcal{E}), h^*\nabla)$ defines an object of $\text{LIC}(V, g^*\alpha)$. It is easy to see that these maps on objects extend to honest functors and that the data $\{U \mapsto \text{Spl}(U, f^*\alpha)\}$, $\{U \mapsto \text{LIC}(U, f^*\alpha)\}$ defines two presheaves of groupoids which we call $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$ correspondingly. The fact that $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$ are actually sheaves of groupoids essentially follows from faithfully flat descent.

As a result to a differential 1-form α on $X^{(1)}$ we associate two gerbes: $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$. From Proposition 2.5.2 it follows that they are actually isomorphic:

Proposition 2.6.1. *There is an equivalence of gerbes $\mathcal{S}(X, \alpha) \simeq \mathcal{L}(X, \alpha)$ for any smooth X and any differential 1-form α on $X^{(1)}$.*

A connected groupoid is defined up to equivalence by the automorphism group of any of its objects. A gerbe is a sheaf of groupoids and there is no uniquely defined automorphism group; instead there is a more complicated object which is called the *band* of a gerbe. The definition of the band for a general gerbe is somewhat abstract, but if the automorphisms of all objects of $\mathcal{G}(U)$ for all U are abelian, the band is given by a certain sheaf of groups on X_{et} . Let us briefly sketch its construction. To an $f : U \rightarrow X$ in X_{et} and to an object A of $\mathcal{G}(U)$ we can associate a sheaf $\underline{\text{Aut}}(A)$ on U_{et} : its sections on any open $g : V \rightarrow U$ are given by $\underline{\text{Aut}}(A)(V) = \text{Aut}(g^*A)$. The fact that $\mathcal{G}$ is a stack of groupoids guarantees that this is a sheaf. We now want to show that if $f : U \rightarrow X$ is an étale cover then there is a natural descent data for $\underline{\text{Aut}}(A)$. Let $p_1, p_2 : U \times_X U \rightarrow U$ be two projections. By the transitivity property there is a cover $h : V \rightarrow U \times_X U$, such that the pull-backs of p_1^*A and p_2^*A to V are isomorphic: there is an arrow $\theta : h^*(p_1^*(A)) \xrightarrow{\sim} h^*(p_2^*(A))$. This produces the isomorphism $\tilde{\theta} : \text{Aut}(h^*(p_1^*(A))) \xrightarrow{\sim} \text{Aut}(h^*(p_2^*(A)))$ which in fact does not depend on θ since by assumption the automorphism groups are abelian and a different choice of θ changes $\tilde{\theta}$ by conjugation. This way it descends to an isomorphism $\theta : p_1^*\underline{\text{Aut}}(A) \xrightarrow{\sim} p_2^*\underline{\text{Aut}}(A)$ and consequently defines the data for the descent of $\underline{\text{Aut}}(A)$ to X . In the similar manner, taking the suitable cover of the fiber product $U_1 \times_X U_2$ of two different covers U_1 and U_2 one can prove that the defined sheaf on X_{et} does not depend on either the cover U or the object A . So this sheaf is well-defined and following [Mo] we call it $\text{Band}(\mathcal{G})$.

The isomorphism classes of gerbes with a given abelian band $\mathcal{B}$ are in 1-to-1 correspondence with the elements of 2nd Čech cohomology $\check{H}^2(X_{et}, \mathcal{B})$. We will only sketch one direction of this bijection. Let $f : U \rightarrow X$ be a cover, such that $f^*\mathcal{G}$ is trivial, namely $f^*\mathcal{G} = */f^*\mathcal{B}$ where $*/G$ is the notation for a groupoid with one object with automorphism group G . Since $\mathcal{G}$ is a stack this isomorphism provides an equivalence $\theta_{12} : p_1^*(*/f^*\mathcal{B}) \xrightarrow{\sim} p_2^*(*/f^*\mathcal{B})$ of sheaves of

groupoids on $U \times_X U$. On the other hand $p_1^*(*/f^*\mathcal{B}) = */p_1^*f^*\mathcal{B} \xrightarrow{\sim} */p_2^*f^*\mathcal{B} = p_2^*(*/f^*\mathcal{B})$ since $f \circ p_1 = f \circ p_2$. The difference between these two isomorphisms is given by a section of $p_1^*f^*\mathcal{B} = p_2^*f^*\mathcal{B}$ or in other words by an element $g_{12} \in \mathcal{B}(U \times_X U)$. The Čech 2-cocycle $\gamma_{\mathcal{G}} \in \mathcal{B}(U \times_X U \times_X U)$ is then defined as $g_{12}g_{23}g_{13}^{-1} \in \mathcal{B}(U \times_X U \times_X U)$.

A gerbe with a given band $\mathcal{B}$ is often called a $\mathcal{B}$ -gerbe. The advantage of gerbes over the cohomology classes is that they have automorphisms and so can be glued together. Namely having a cover $U \rightarrow X$ and a gerbe on U with the gluing data we can uniquely glue out a gerbe on X . And while locally all the objects in the gerbe are isomorphic and the gerbe is trivial, globally it is usually not. Gerbes produce a geometrization for H^2 in the similar manner as torsors produce a geometrization for H^1 .

Let's now return to our example. We claim that $\text{Band}(\mathcal{S}(X, \alpha)) \simeq \mathbb{G}_m$. Indeed, locally Azumaya algebra $\mathcal{D}_{X,\alpha}$ is split, so we have $f^*(\mathcal{D}_{X,\alpha}) \simeq \text{End}(\mathcal{O}_{U(1)}^{p^{\dim X}})$ for some étale cover $f : U \rightarrow X$. The automorphisms of such splitting bundle are the maps $\mathcal{O}_{U(1)}^{\oplus p^{\dim X}} \rightarrow \mathcal{O}_{U(1)}^{\oplus p^{\dim X}}$ which are 1) invertible, 2) induce a trivial map on the $\text{End}(\mathcal{O}_{U(1)}^{p^{\dim X}})$. In other words they are given by the kernel of the map $\text{GL}_{p^{\dim X}}(\mathcal{O}_{U(1)}) \rightarrow \text{PGL}_{p^{\dim X}}(\mathcal{O}_{U(1)}) = \text{Aut}(\text{End}_{\mathcal{O}_{U(1)}}(\mathcal{O}_{U(1)}^{p^{\dim X}}))$ which is equal to $\mathbb{G}_m(\mathcal{O}_{U(1)})$. On the line bundles' side, any $(\mathcal{E}, \nabla)$ is locally isomorphic to $(\mathcal{O}_U, d + \omega)$ for some closed 1-form ω . Since $\text{curv}_p(\nabla) = \alpha$, ω also should satisfy a condition $\omega^{(1)} - \mathbb{C}(\omega) = \alpha$. The automorphism of $(\mathcal{E}, \nabla)$ is given by the automorphism of $\mathcal{E}$, preserving ∇ . If $f : \mathcal{O}_U \xrightarrow{\sim} \mathcal{O}_U$ is an automorphism of $\mathcal{O}_U$ given by an invertible function $f \in \mathcal{O}_X^\times$, then $f^*(d + \omega) = d + d \log f + \omega$. The kernel of $d \log : \mathcal{O}_U^\times \rightarrow \Omega_U^1$ is given by $(\mathcal{O}_U^\times)^p = \mathcal{O}_{U(1)}^\times \subset \mathcal{O}_U^\times$, or in other words by the values of the sheaf $\mathbb{G}_m$ on $U^{(1)}$.

As was mentioned before, there is a natural bijection between the set of isomorphism classes of $\mathcal{B}$ -gerbes and the group $\check{H}^2(X, \mathcal{B})$. Given an Azumaya algebra $\mathcal{A}$ on X we can associate to it a $\mathbb{G}_m$ -gerbe of its splittings $\mathcal{S}(\mathcal{A})$. Both $\mathcal{A}$ and $\mathcal{S}(\mathcal{A})$ give classes in $\check{H}^2(X, \mathbb{G}_m)$ and it is not hard to see that it is in fact the same class. Applying this to $\mathcal{A} = \mathcal{D}_{X,\alpha}$ we get that both $\mathcal{S}(X, \alpha)$ and $\mathcal{L}(X, \alpha)$ represent the same class $c_X(\alpha) = [\mathcal{D}_{X,\alpha}]$.

2.7. Additivity of the map c_X : cohomology. Here we present a more classical point of view on the map c_X . Namely we will obtain the homomorphism $c_X : H^0(X_{\text{ét}}^{(1)}, \Omega_{X(1)}^1) \rightarrow H^2(X_{\text{ét}}^{(1)}, \mathcal{O}_{X(1)}^\times)$ from a map $\Omega_{X(1)}^1 \rightarrow \mathcal{O}_{X(1)}^\times[2]$ in the derived category of étale sheaves on X . Such a map is the same as an element of $\text{Ext}^2(\Omega_{X(1)}^1, \mathcal{O}_{X(1)}^\times)$.

Proposition 2.7.1. *There is a 4-term exact sequence*

$$0 \rightarrow \mathcal{O}_{X(1)}^\times \xrightarrow{\text{Fr}_X^*} \mathcal{O}_X^\times \xrightarrow{d \log} \Omega_{X,\text{cl}}^1 \xrightarrow{\bullet^{(1)} - \mathbb{C}} \Omega_{X(1)}^1 \rightarrow 0.$$

Proof. For the usual proof not using gerbes or differential operators see ([Mi], Proposition 4.14). We will deduce the exactness from Proposition 2.3.1 and Proposition 2.5.2. The exactness in the second term is obvious: étale locally (as well as globally) $d \log f = 0 \Leftrightarrow df = 0 \Leftrightarrow f \in \mathcal{O}_{X(1)}^\times$. For the third term we need to prove that locally $\omega^{(1)} - \mathbb{C}(\omega) = 0 \Leftrightarrow \omega = d \log f$. As we have seen, the condition $\omega^{(1)} - \mathbb{C}(\omega) = 0$ for some cover U means that $(\mathcal{O}_U, d + \omega)$ has p -curvature 0 and is a splitting bundle for $\mathcal{D}_{U,0}$. So after refining the cover it should be isomorphic to $(\mathcal{O}_U, d)$ and this means exactly that $\omega = d \log f$ for some function f on U . It remains only to prove that the map $\bullet^{(1)} - \mathbb{C}$ is surjective. Let α be a differential 1-form on a cover $U^{(1)} \rightarrow X^{(1)}$. Then by the Azumaya property there is a refinement $V^{(1)} \rightarrow U^{(1)}$, such that the pull-back of $\mathcal{D}_{X,\alpha}$ is split. So there is a line bundle with a flat connection $(\mathcal{E}, \nabla)$ on V , such that $\text{curv}_p(\nabla) = \alpha$. Refining the cover we can assume that $(\mathcal{E}, \nabla) = (\mathcal{O}_U, d + \omega)$ and so $\omega^{(1)} - \mathbb{C}(\omega) = \alpha$. $\square$

By Yoneda's presentation for Ext-groups, such a sequence provides a class in $\text{Ext}^2(\Omega_{X^{(1)}}^1, \mathcal{O}_{X^{(1)}}^\times)$ which produces a map $\bar{c}_X : H^0(X_{et}^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow H^2(X_{et}^{(1)}, \mathcal{O}_{X^{(1)}}^\times)$. The image is contained in p -torsion, so c_X lands in $\text{Br}(X^{(1)})[p]$. We would like to know that $\bar{c}_X$ is the same as the map c_X that we defined before.

2.8. Additivity of the map c_X : Picard groupoids. Let $f : U \rightarrow X$ be an étale open set and let $(\mathcal{E}_1, \nabla_1), (\mathcal{E}_2, \nabla_2)$ be line bundles with integrable connections on U . Their tensor product $\mathcal{E}_1 \otimes \mathcal{E}_2$ is endowed with a natural flat connection $\nabla : \mathcal{E}_1 \otimes \mathcal{E}_2 \rightarrow \mathcal{E}_1 \otimes \mathcal{E}_2 \otimes \Omega_X^1$ given by $\nabla = \nabla_1 \otimes \text{id} + \text{id} \otimes \nabla_2$ or, in dual presentation, the map $\nabla : \mathcal{T}_{X^{(1)}} \rightarrow \text{End}_{\mathcal{O}_{U^{(1)}}}(\mathcal{E}_1 \otimes \mathcal{E}_2)$ is given by $\partial \mapsto \nabla_1(\partial) \otimes \text{id} + \text{id} \otimes \nabla_2(\partial)$. The p -curvature of the tensor product is easy to compute, namely:

$$\begin{aligned} \text{curv}_p(\nabla)(\partial) &= \nabla(\partial)^p - \nabla(\partial^{[p]}) = (\nabla_1(\partial) \otimes \text{id} + \text{id} \otimes \nabla_2(\partial))^p - \nabla_1(\partial^{[p]}) \otimes \text{id} - \text{id} \otimes \nabla_2(\partial^{[p]}) = \\ &= \nabla_1(\partial)^p \otimes \text{id} - \nabla_1(\partial^{[p]}) \otimes \text{id} + \text{id} \otimes \nabla_2(\partial)^p - \text{id} \otimes \nabla_2(\partial^{[p]}) = \text{curv}_p(\nabla_1)(\partial) + \text{curv}_p(\nabla_2)(\partial). \end{aligned}$$

In particular if $\mathcal{E}_1, \mathcal{E}_2$ were line bundles and $\text{curv}_p(\nabla_1) = \alpha_1, \text{curv}_p(\nabla_2) = \alpha_2$, then $\text{curv}_p(\nabla) = \alpha_1 + \alpha_2$. Note that this way $(\mathcal{E}_1, \nabla_1) \otimes (\mathcal{E}_2, \nabla_2) := (\mathcal{E}_1 \otimes \mathcal{E}_2, \nabla)$ is a splitting bundle for $\mathcal{D}_{X, \alpha_1 + \alpha_2}$. In local coordinates $(\mathcal{O}_U, d + \omega_1) \otimes (\mathcal{O}_U, d + \omega_2) = (\mathcal{O}_U, d + \omega_1 + \omega_2)$.

This operation induces a map of gerbes $\otimes : \mathcal{L}(X, \alpha_1) \times \mathcal{L}(X, \alpha_2) \rightarrow \mathcal{L}(X, \alpha_1 + \alpha_2)$. Note that the source here is a $\mathbb{G}_m \times \mathbb{G}_m$ -gerbe while the target is a $\mathbb{G}_m$ -gerbe. The kernel of the corresponding map $\mathbb{G}_m \times \mathbb{G}_m \rightarrow \mathbb{G}_m$ is given by $\{(f, f^{-1})\} \subset \mathbb{G}_m \times \mathbb{G}_m$ since $f \otimes f^{-1}$ acts trivially on $\mathcal{E}_1 \otimes \mathcal{E}_2$.

We see that the natural operation of the tensor product mixes all the p -curvatures, so we would like to consider $\mathcal{L}(X, \alpha)$ for all α at the same time. Namely we define the stack of groupoids $\mathcal{L}(X)$ by $\mathcal{L}(X)(U) = \sqcup_\alpha \mathcal{L}(X, \alpha)(U)$, or in other words $\mathcal{L}(X) = \sqcup_\alpha \mathcal{L}(X, \alpha)$. It is then endowed with a symmetric monoidal structure $\otimes : \mathcal{L}(X) \times \mathcal{L}(X) \rightarrow \mathcal{L}(X)$ which turns it into a *stack of Picard groupoids* or a *Picard stack*. For the formal definition and a very good introduction to Picard stacks see ([SGA], 1.4). Briefly, *Picard groupoid* is a groupoid $\mathcal{P}$ endowed with a functor $+$: $\mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$ together with functorial isomorphisms $\tau : X + Y \xrightarrow{\sim} Y + X$, $\sigma : (X + Y) + Z \xrightarrow{\sim} X + (Y + Z)$ which turn it into what is called *associative strictly commutative monoidal functor*, and such that the functor $X \mapsto X + Y$ is an isomorphism for all $Y \in \text{Ob}(\mathcal{P})$. The typical example of Picard groupoid is given by the groupoid $\text{Pic}(X)$ of line bundles on X with the natural structure of the tensor product. This example also explains the name for the definition.

Every Picard groupoid $\mathcal{P}$ has a unique unit object $e \in \text{Ob}(\mathcal{P})$ with an isomorphism $e + e \xrightarrow{\sim} e$ and isomorphisms $e + X \xrightarrow{\sim} X$ functorial in X , satisfying some natural axioms. For $\text{Pic}(X)$ it is given by trivial bundle $\mathcal{O}_X$, in our case of $\mathcal{L}(X)(U)$ — by $(\mathcal{O}_U, d)$. We have a group $\text{Aut}(e)$ which is abelian and which is isomorphic to $\text{Aut}(X)$ for any $X \in \text{Ob}(\mathcal{P})$ under the functor $Y \mapsto Y + X$. The set of isomorphism classes of objects of $\mathcal{P}$ under operation $+$ also forms a group which we call $\text{Iso}(\mathcal{P})$.

A *Picard stack* on X_{et} is by definition a stack of groupoids $\mathcal{P}$ on X_{et} endowed with an operation $+$: $\mathcal{P} \times \mathcal{P} \rightarrow \mathcal{P}$ together with functorial isomorphisms $\tau : X + Y \xrightarrow{\sim} Y + X$, $\sigma : (X + Y) + Z \xrightarrow{\sim} X + (Y + Z)$ which for any $U \in X_{et}$ turn $\mathcal{P}(U)$ into a Picard groupoid. Since $\mathcal{P}$ is a stack, the presheaf $U \mapsto \text{Aut}(e)(U)$ which associate to U the automorphism group of the unit object in $\mathcal{P}(U)$ is actually a sheaf which we call $\underline{\text{Aut}}(e)$. Presheaf of isomorphism classes of objects $U \mapsto \text{Iso}(\mathcal{P})(U)$ does not form a sheaf and we denote by $\underline{\text{Iso}}(\mathcal{P})$ its sheafification.

Following Deligne we denote by $C^{[-1, 0]}(X)$ the category of complexes $\mathcal{K}^\bullet$ of sheaves of abelian groups on X_{et} , such that $\mathcal{K}^i = 0$ for $i \notin [-1, 0]$. To every such complex $d : \mathcal{K}^{-1} \rightarrow \mathcal{K}^0$ we can associate a prestack of Picard groupoids $\text{pch}(\mathcal{K}^\bullet)$, such that

- for any U in X_{et} , we have $Ob(\text{pch}(\mathcal{K}^\bullet)(U)) = \mathcal{K}^0(U)$,
- for any two $x, y \in \mathcal{K}^0(U)$ a morphism from x to y is given an element $f \in \mathcal{K}^{-1}(U)$, such that $df = y - x$,
- composition of morphisms is given by the addition in $\mathcal{K}^{-1}(U)$,
- functor $+$ is given by the addition on $\mathcal{K}^{-1}$ and $\mathcal{K}^0$.

The Picard stack $\text{ch}(\mathcal{K}^\bullet)$ is defined as the sheafification of the prestack $\text{pch}(\mathcal{K}^\bullet)$. The sheaf $\mathcal{H}^0(\mathcal{K}^\bullet)$ then can be interpreted as $\underline{\text{Iso}}(\text{ch}(\mathcal{K}^\bullet))$, while $\mathcal{H}^{-1}(\mathcal{K}^\bullet)$ is nothing but the sheaf $\underline{\text{Aut}}(e)$. We have a 4-term exact sequence

$$0 \rightarrow \mathcal{H}^{-1}(\mathcal{K}^\bullet) \rightarrow \mathcal{K}^{-1} \xrightarrow{d} \mathcal{K}^0 \rightarrow \mathcal{H}^0(\mathcal{K}^\bullet) \rightarrow 0,$$

which can be presented by some class $\gamma \in \text{Ext}^2(\mathcal{H}^0(\mathcal{K}^\bullet), \mathcal{H}^{-1}(\mathcal{K}^\bullet))$. Note that $\overline{\gamma}$ also produces a natural map $\overline{\gamma}: H^0(X_{et}, \mathcal{H}^0(\mathcal{K}^\bullet)) \rightarrow H^2(X_{et}, \mathcal{H}^{-1}(\mathcal{K}^\bullet))$.

Let's now fix a global section $\alpha \in H^0(X_{et}, \mathcal{H}^0(\mathcal{K}^\bullet)) = \underline{\text{Iso}}(\text{ch}(\mathcal{K}^\bullet))(X)$ and consider the full substack $\text{ch}_\alpha(\mathcal{K}^\bullet) \subset \text{ch}(\mathcal{K}^\bullet)$, given by the property that objects $\text{ch}_\alpha(\mathcal{K}^\bullet)(U)$ are objects of $\text{ch}(\mathcal{K}^\bullet)(U)$ that are isomorphic to $\alpha|_U$. Substacks $\text{ch}_\alpha(\mathcal{K}^\bullet)$ for all α are the connected components of the Picard stack $\text{ch}(\mathcal{K}^\bullet)$. It is easy to see that by definition each substack $\text{ch}_\alpha(\mathcal{K}^\bullet)$ is an $\mathcal{H}^{-1}(\mathcal{K}^\bullet)$ -gerbe, so it gives a class $[\text{ch}(\alpha)] \in \hat{H}^2(X_{et}, \mathcal{H}^{-1}(\mathcal{K}^\bullet))$. Sections of the sheaf of objects $\underline{\text{Iso}}(\text{ch}_\alpha(\mathcal{K}^\bullet))$ does not anymore have a structure of abelian groups unless $\alpha = 0$. Note though that the sheaf $\underline{\text{Iso}}(\text{ch}_\alpha(\mathcal{K}^\bullet))$ has a natural structure of a $\underline{\text{Iso}}(\text{ch}_0(\mathcal{K}^\bullet))$ -torsor with respect to the $+$ operation on $\text{ch}(\mathcal{K}^\bullet)$. Also note that by definition $\underline{\text{Iso}}(\text{ch}_0(\mathcal{K}^\bullet))(X)$ is in fact equal to $\text{Im } d = \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet)$.

Lemma 2.8.1. *We have $[\text{ch}(\alpha)] = \overline{\gamma}(\alpha)$.*

Proof. The 4-term exact sequence above can be divided into two exact sequences

$$0 \rightarrow \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet) \xrightarrow{d} \mathcal{K}^0 \rightarrow \mathcal{H}^0(\mathcal{K}^\bullet) \rightarrow 0, \quad 0 \rightarrow \mathcal{H}^{-1}(\mathcal{K}^\bullet) \rightarrow \mathcal{K}^{-1} \rightarrow \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet) \rightarrow 0.$$

Map $\overline{\gamma}$ comes as the composition of the two boundary maps in the long exact sequences of cohomology, namely $\delta_1: H^0(X_{et}, \mathcal{H}^0(\mathcal{K}^\bullet)) \rightarrow H^1(X_{et}, \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet))$ and $\delta_2: H^1(X_{et}, \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet)) \rightarrow H^2(X_{et}, \mathcal{H}^{-1}(\mathcal{K}^\bullet))$. We claim that the class $\delta_1(\alpha) \in H^1(X_{et}, \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet))$ corresponds exactly to the $\mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet)$ -torsor, given by $\underline{\text{Iso}}(\text{ch}_\alpha(\mathcal{K}^\bullet))(X)$. Indeed, it can be computed using Čech cohomology: let $\varphi: U \rightarrow X$ be a cover and let α_0 be an element of $\mathcal{K}^0(U)$ which maps to $\varphi^*\alpha$, then $\alpha_{-1} = p_1^*\alpha_0 - p_2^*\alpha_0$ maps to 0 and so gives a section $H^0(U \times_X U, \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet))$ which for sufficiently small U is equal to $\delta_1(\alpha)$. The element $\alpha_{-1} = p_1^*\alpha_0 - p_2^*\alpha_0$ is, by definition, an isomorphism between objects $p_1^*\alpha_0$ and $p_2^*\alpha_0$ and it is easy to see that α_{-1} is in fact a Čech 1-cocycle representing $\mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet)$ -torsor $\underline{\text{Iso}}(\text{ch}_\alpha(\mathcal{K}^\bullet))(X)$. Now take $U \times_X U \times_X U$ with natural projections p_{12}, p_{13}, p_{23} to $U \times_X U$. α_{-1} is a cocycle, so $p_{12}^*\alpha_{-1} + p_{23}^*\alpha_{-1} - p_{13}^*\alpha_{-1} = 0 \in \mathcal{K}^{-1}/\mathcal{H}^{-1}(\mathcal{K}^\bullet)(U \times_X U \times_X U)$. Possibly refining the cover and taking any lift $\tilde{\alpha}_{-1}$ of α_{-1} in $\mathcal{K}^{-1}(U \times_X U)$, we get a class $p_{12}^*\tilde{\alpha}_{-1} + p_{23}^*\tilde{\alpha}_{-1} - p_{13}^*\tilde{\alpha}_{-1}$ in $\mathcal{H}^{-1}(\mathcal{K}^\bullet)(U \times_X U \times_X U)$ which for sufficiently small U coincides with $\overline{\gamma}(\alpha)$. It is also not hard to see that from the point of view of gerbes, this is exactly how the Čech 2-cocycle for the $\mathcal{H}^{-1}(\mathcal{K}^\bullet)$ -gerbe $\text{ch}_\alpha(\mathcal{K}^\bullet)$ was defined. So $[\text{ch}(\alpha)] = \overline{\gamma}(\alpha)$. $\square$

The complex that we are interested in is given by the middle arrow in the 4-term exact sequence from Proposition 2.7.1:

$$\mathcal{O}^\bullet := \mathcal{O}_X^\times \xrightarrow{d \log} \Omega_{X,cl}^1.$$

We claim that $\text{ch}(\mathcal{O}^\bullet) = \mathcal{L}(X)$. Indeed for any $U \in X_{et}$ we have a map from $\Omega_{X,cl}^1(U)$ to the set of objects of $\mathcal{L}(X)(U)$ given by $\omega \mapsto (\mathcal{O}_U, d + \omega)$ and, since any line bundle is locally trivial, this map is locally an isomorphism. Isomorphisms between bundles with connection are

given by invertible functions f , which change the connection by $d \log f$. Finally, $(\mathcal{O}_U, d + \omega_1) \otimes (\mathcal{O}_U, d + \omega_2) = (\mathcal{O}_U, d + \omega_1 + \omega_2)$, as well as $f_1 \otimes f_2$ acts on the tensor product by $f_1 f_2$, so the tensor product of line bundles corresponds to the addition on the complex $\mathcal{O}^\bullet$. Note that from the 4-term exact sequence we have $\underline{\text{Aut}}(e) = \mathcal{O}_{X^{(1)}}^\times$ and $\underline{\text{Iso}}(\mathcal{L}(X)) = \Omega_X^1$, where the last equality is given by the p -curvature. From this we also get that $\text{ch}_\alpha(\mathcal{O}^\bullet) = \mathcal{L}(X, \alpha)$ for any $\alpha \in H^0(X_{\text{et}}^{(1)}, \Omega_{X^{(1)}}^1)$ and in particular that $c_X(\alpha) = [\text{ch}(\alpha)]$. Moreover, applying Lemma 2.8.1 to $\mathcal{O}^\bullet$ we get that $[\text{ch}(\alpha)] = \overline{\gamma}(\alpha)$, which by definition is $\overline{c}_X(\alpha)$. We get the following proposition:

Proposition 2.8.1. *Let X be a smooth scheme over an algebraically closed field $\mathbb{k}$ of characteristic p and let $c_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$ be the map given by $\alpha \mapsto [\mathcal{D}_{X, \alpha}]$. Let also $\overline{c}_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$ be the map obtained from the 4-term exact sequence*

$$0 \rightarrow \mathcal{O}_{X^{(1)}}^\times \xrightarrow{\text{Fr}_X^*} \mathcal{O}_X^\times \xrightarrow{d \log} \Omega_{X, \text{cl}}^1 \xrightarrow{\bullet^{(1)} - \text{C}} \Omega_{X^{(1)}}^1 \rightarrow 0.$$

Then $c_X = \overline{c}_X$ and consequently is a homomorphism of abelian groups.

Corollary 2.8.1. *Let $\alpha = \omega^{(1)} - \text{C}(\omega)$. Then $c_X(\alpha) = 0$ and Azumaya algebra $\mathcal{D}_{X, \alpha}$ is split.*

Proof. By definition of the map $\overline{c}_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$ coming from the 4-term exact sequence we have $\text{Im}(\bullet^{(1)} - \text{C}) \subset \text{Ker}(\overline{c}_X)$. The statement of the corollary then follows from $c_X = \overline{c}_X$. It can also be seen directly, since by Proposition 2.5.2 $(\text{Fr}_X)_* \mathcal{O}_X, d + \omega$ is a splitting bundle for $\mathcal{D}_{X, \alpha}$. $\square$

3. DESCENT ÉTALE LOCALLY

3.1. Setup. Now we will return to a very particular question about algebras $\mathcal{D}_{X, \alpha}$, which shows up naturally in the theory of quantizations of symplectic resolutions in characteristic p .

Let $\pi : X \rightarrow Y$ be a resolution of singularities over an algebraically closed field $\mathbb{k}$ of characteristic $p > 0$, such that $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^i \pi_* \mathcal{O}_X = 0$, for $i = 1, 2$. Let α be a 1-form on $X^{(1)}$ and let $\mathcal{D}_{X, \alpha}$ be the associated Azumaya algebra (see Definition 2.4.1).

We would like to investigate the following question: can the class $c_X(\alpha) = [\mathcal{D}_{X, \alpha}]$ in the Brauer group $\text{Br}(X^{(1)})$ of $X^{(1)}$ be descended to a class in $\text{Br}(Y^{(1)})$? In other words, does the class $c_X(\alpha)$ lie in the image of the natural map $\text{Br}(Y^{(1)}) \rightarrow \text{Br}(X^{(1)})$?

3.2. Splitting étale locally on Y . In this subsection we prove that the classes $c_X(\alpha)$ descend to Y étale locally for any α , provided $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^i \pi_* \mathcal{O}_X = 0$, for $i = 1, 2$. By “descends étale locally” we mean that there exists an étale cover $f : U \rightarrow Y$ such that the pull-back $f^* c_X(\alpha)$ in $\text{Br}((X \times_Y U)^{(1)})$ descends to U , namely $f^* c_X(\alpha) = \pi^* [\mathcal{C}]$ for some Azumaya algebra $\mathcal{C}$ on U . Shrinking U we can assume that $\mathcal{C}$ is split, so equivalently we can ask if there exists $f : U \rightarrow Y$ such that the Azumaya algebra $f^* \mathcal{D}_{X, \alpha} = \mathcal{D}_{X \times_Y U, f^* \alpha}$ is split. The idea is to use Corollary 2.8.1 and to prove that the operator $\bullet^{(1)} - \text{C}$ is surjective locally on Y (in the same sense).

Remark 3.2.1. Note that under the assumption $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ the map $\pi^* : \text{Br}(Y^{(1)}) \rightarrow \text{Br}(X^{(1)})$ is an injection. Indeed, by the projection formula $\pi_* \pi^* \mathcal{A} = \pi_* \pi^* \mathcal{O}_{Y^{(1)}} \otimes_{\mathcal{O}_{Y^{(1)}}} \mathcal{A} = \mathcal{A}$ and if $\pi^* \mathcal{A}$ were split on $X^{(1)}$ we would also obtain a splitting of $\mathcal{A}$ on $Y^{(1)}$.

We start with the following linear algebraic lemma which is probably due to V. Drinfeld:

Lemma 3.2.1. *Let $A, B : V \rightarrow W$ be two maps between vector spaces over algebraically closed field $\mathbb{k}$ of characteristic $p > 0$, such that A is Frobenius-linear and B is linear and surjective. Then $A - B : V \rightarrow W$ is surjective as a map of abelian groups.*

Proof. We consider V and W as the sets of $\mathbb{k}$ -points of the corresponding affine spaces $\mathbb{A}(V)$ and $\mathbb{A}(W)$. Any linear operator T from V to W induces a natural map $\tilde{T}$ between corresponding affine spaces. There also exists a linear map $A' : V \rightarrow W$, such that the map $\text{Fr}_W \circ \tilde{A}' : \mathbb{A}(V) \rightarrow \mathbb{A}(W)$ coincides with A on the level of $\mathbb{k}$ -points. Since the differential of the Frobenius map at any point is equal to 0, we get that so does the differential of $\text{Fr}_W \circ \tilde{A}'$. From this we see that $d(\text{Fr}_W \circ \tilde{A}' - \tilde{B}) = -d\tilde{B}$. After identifying the tangent spaces at any point with V or W correspondingly, the differential $d\tilde{B}$ is given by B on the tangent space of any point. We get that $\text{Fr}_W \circ \tilde{A}' - \tilde{B}$ is a submersion and so is an open map. It follows that $\text{Im}(\text{Fr}_W \circ \tilde{A}' - \tilde{B})$ is open in $\mathbb{A}(W)$.

Let $U \subset \mathbb{A}(W)$ be an open set. Then for any closed point $w \in W(\mathbb{k})$ the subset $w - U \subset \mathbb{A}(W)$ is also open and so the intersection $U \cap (w - U)$ is non-empty. In other words w lies in $U(\mathbb{k}) + U(\mathbb{k}) \subset W(\mathbb{k})$. Taking $U = \text{Im}(\tilde{A} - \tilde{B})$, which is closed under addition, we get that $w \in U(\mathbb{k})$ and that $A - B$ is surjective. $\square$

Next lemma proves that the Cartier operator is surjective:

Lemma 3.2.2. *Let $\pi : X \rightarrow Y$ be a proper morphism of algebraic varieties over $\mathbb{k}$, such that X is smooth and $R^i\pi_*\mathcal{O}_X = 0$ for $i = 1, 2$. Then the Cartier operator $\pi_*\mathbf{C} : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_{X^{(1)}}^1$ is surjective.*

Proof. From $R^1\pi_*\mathcal{O}_X = R^2\pi_*\mathcal{O}_X = 0$ and short exact sequence of sheaves $0 \rightarrow \mathcal{O}_{X^{(1)}} \rightarrow \mathcal{O}_X \rightarrow \mathcal{O}_X/\mathcal{O}_{X^{(1)}} \rightarrow 0$, applying $R^\bullet\pi_*$ we see that $R^1\pi_*(\mathcal{O}_X/\mathcal{O}_{X^{(1)}}) = 0$.

By Remark 2.2 we have the following short exact sequence

$$0 \rightarrow \mathcal{O}_X/\mathcal{O}_{X^{(1)}} \xrightarrow{d} \Omega_{X,cl}^1 \xrightarrow{\mathbf{C}} \Omega_{X^{(1)}}^1 \rightarrow 0$$

and applying $R^\bullet\pi_*$ we get that the map $\pi_*\mathbf{C} : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_{X^{(1)}}^1$ is a surjection. $\square$

Note that if Y is affine the induced map on global sections $\mathbf{C} : H^0(Y^{(1)}, \pi_*\Omega_{X,cl}^1) \rightarrow H^0(Y^{(1)}, \pi_*\Omega_{X^{(1)}}^1)$ is also a surjection. Notice that the spaces of global sections $H^0(Y^{(1)}, \pi_*\Omega_{X,cl}^1)$, $H^0(Y^{(1)}, \pi_*\Omega_{X^{(1)}}^1)$ and the map $\bullet^{(1)} - \mathbf{C}$ fit pretty well in the conditions of Lemma 3.2.1. Indeed $\bullet^{(1)}$ is Frobenius-linear, while $\mathbf{C}$ is linear and surjective. The only thing is that when Y is affine, X is usually not proper and both spaces $H^0(Y^{(1)}, \pi_*\Omega_{X,cl}^1)$ and $H^0(Y^{(1)}, \pi_*\Omega_{X^{(1)}}^1)$ are infinite dimensional.

Nevertheless, if $Y = \text{Spec } \mathbb{k}$, X is proper and we obtain the following:

Proposition 3.2.1. *Let X be a proper smooth variety over $\mathbb{k}$, such that $H^1(X, \mathcal{O}_X) = H^2(X, \mathcal{O}_X) = 0$. Then for any differential 1-form $\alpha \in H^0(X, \Omega_X^1)$ the Azumaya algebra $\mathcal{D}_{X,\alpha}$ is split.*

Proof. We apply Lemma 3.2.1 to the map $\bullet^{(1)} - \mathbf{C} : H^0(X, \Omega_{X,cl}^1) \rightarrow H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$. The image of this map lies in the kernel of $c_X : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)})$, so it is enough to prove that $\bullet^{(1)} - \mathbf{C}$ is surjective. Map $\mathbf{C}$ is $\mathbb{k}$ -linear, $\bullet^{(1)}$ is Frobenius-linear and by Lemma 3.2.1 $\mathbf{C}$ is surjective. So we are in the context of Lemma 3.2.1, $\bullet^{(1)} - \mathbf{C}$ is surjective and we are done. $\square$

If Y is non-trivial, we slightly generalise the argument to prove that $\mathcal{D}_{X,\alpha}$ splits on the formal neighbourhoods of the fibers of π .

Proposition 3.2.2. *Let $\pi : X \rightarrow Y$ be a proper morphism of algebraic varieties over $\mathbb{k}$ such that X is smooth and $R^i\pi_*\mathcal{O}_X = 0$ for $i = 1, 2$. Then for any $\alpha \in H^0(X, \Omega_X^1)$ the algebra $\mathcal{D}_{X,\alpha}$ splits on the formal neighbourhood of every fiber of π .*

Proof. Let's take any closed point $y \in Y(\mathbb{k})$. The question is Zariski local on Y , so we can assume that Y is affine. Let $\widehat{\mathcal{O}}_y$ be the completion of the ring $\Gamma(Y, \mathcal{O}_Y)$ at the point y and let $\widehat{X}_y$ denote the formal neighbourhood of the fiber $\pi^{-1}(y)$. Let also Z denote the formal fiber: $Z = X \times_Y \text{Spec } \widehat{\mathcal{O}}_y$. By the Grothendieck's existence theorem ([EGA], III, Theorem 5.1.4) one has an equivalence of categories $\text{Coh}(\widehat{X}_y) \simeq \text{Coh}(Z)$. In particular, $\mathcal{D}_{X,\alpha}$ is split on $\widehat{X}_y^{(1)}$ if and only if it is split on $Z^{(1)}$. The Frobenius twist $Z^{(1)}$ is naturally identified with $X^{(1)} \times_{Y^{(1)}} \text{Spec } \widehat{\mathcal{O}}_{\text{Fr}_Y(y)}^{(1)}$. We have a natural map of schemes $\text{Fr}_Z : Z \rightarrow Z^{(1)}$ and the maps $\pi_*(\bullet^{(1)}) : \pi_*(\text{Fr}_Z)_* \Omega_Z^1 \rightarrow \pi_* \Omega_{Z^{(1)}}^1$ and $\pi_* \mathbb{C} : \pi_*(\text{Fr}_Z)_* \Omega_{Z,\text{cl}}^1 \rightarrow \pi_* \Omega_{Z^{(1)}}^1$. As before, to split $\mathcal{D}_{X,\alpha}$ on $Z^{(1)}$ it is enough to find a closed 1-form ω on Z such that $\omega^{(1)} - \mathbb{C}(\omega) = \alpha|_{Z^{(1)}}$.

Since π is proper, globally defined closed 1-forms on Z form a finitely generated module $M = \pi_*(\text{Fr}_Z)_* \Omega_{Z,\text{cl}}^1$ over $\widehat{\mathcal{O}}_{\text{Fr}_Y(y)}^{(1)}$. So does the module $N = \pi_* \Omega_{Z^{(1)}}^1$ of all globally defined 1-forms on $Z^{(1)}$. Let $\mathfrak{m}_{\text{Fr}_Y(y)} \subset \widehat{\mathcal{O}}_{\text{Fr}_Y(y)}^{(1)}$ be the maximal ideal of the point $\text{Fr}_Y(y) \in Y^{(1)}(\mathbb{k})$. We endow M and N with natural descending filtrations: $F_i M := \mathfrak{m}_{\text{Fr}_Y(y)}^i M$, $F_i N := \mathfrak{m}_{\text{Fr}_Y(y)}^i N$. Modules M and N are complete with respect to the topologies induced by $F_\bullet$.

The map $\omega \mapsto \omega^{(1)} - \mathbb{C}(\omega)$ is continuous with respect to these topologies. Namely $\mathbb{C}(f^{(1)} \cdot \omega) = f^{(1)} \mathbb{C}(\omega)$ and $(f^{(1)} \cdot \omega)^{(1)} = (f^{(1)})^p \omega^{(1)}$, so $(\bullet^{(1)} - \mathbb{C})(F_i M) \subset F_i N$. We get that it is enough to solve the equation $\omega^{(1)} - \mathbb{C}(\omega) = \alpha|_{Z^{(1)}}$ on the level of the associated graded modules $\text{gr}_{F_\bullet}(M)$ and $\text{gr}_{F_\bullet}(N)$.

Note that the twist $\bullet^{(1)}$ maps $F_i M$ not just to $F_i N$, but to $F_{pi} N$. It follows that for $i > 0$ the map $\text{gr}_i(\bullet^{(1)}) : \text{gr}_i(M) \rightarrow \text{gr}_i(N)$ is 0 and so $\text{gr}_i(\bullet^{(1)} - \mathbb{C}) = \text{gr}_i(-\mathbb{C})$ is surjective by Lemma 3.2.2. It remains only to prove that $\text{gr}_0(\bullet^{(1)} - \mathbb{C})$ is surjective. But the spaces $\text{gr}_0 M = M/F_1 M$ and $\text{gr}_0 N = N/F_1 N$ are finite-dimensional, $\text{gr}_0(\bullet^{(1)})$ is Frobenius-linear and $\text{gr}_0(\mathbb{C})$ is $\mathbb{k}$ -linear and surjective. Lemma 3.2.1 can be applied and so $\text{gr}_0(\bullet^{(1)} - \mathbb{C})$ is surjective too. We are done. $\square$

Remark 3.2.2. As we will see soon, in fact it is more important that $\mathcal{D}_{X,\alpha}$ splits on $Z^{(1)}$ than that it splits on the formal neighbourhood of the fiber.

Finally using Popescu's theorem (a fancy version of Artin approximation, which was introduced to the first author by Brian Conrad) we prove that $\mathcal{D}_{X,\alpha}$ also splits on some étale neighbourhood of each fiber.

Theorem 3.2.1. *Under the assumptions of Lemma 3.2.2, there exists an étale cover $U \rightarrow Y$ such that the pullback of $\mathcal{D}_{X,\alpha}$ to $U \times_Y X$ splits.*

Proof. Since $Y^{(1)}$ is quasi-compact, it is enough to prove that for each $y \in Y^{(1)}(\mathbb{k})$ Azumaya algebra $\mathcal{D}_{X,\alpha}$ splits on some étale neighbourhood of y . In particular, we can assume that Y is affine, $Y = \text{Spec } \Gamma(Y, \mathcal{O}_Y)$. Let $\mathcal{O}_y$ be the usual Zariski localisation of $\mathcal{O}_Y$ at the point y and let $\widehat{\mathcal{O}}_y$ be its formal completion. The field $\mathbb{k}$ is excellent like any other field, $\Gamma(Y, \mathcal{O}_Y)$ is a finitely generated algebra over $\mathbb{k}$ and is excellent, finally $\mathcal{O}_y$ is a localisation of $\mathcal{O}_Y$ and so is excellent too. Any excellent ring is a G-ring, namely if A is an excellent local ring, then the natural map $A \rightarrow \widehat{A}$ to its formal completion is regular. We get that the map $\mathcal{O}_y \rightarrow \widehat{\mathcal{O}}_y$ is regular. So does the natural map $\Gamma(Y, \mathcal{O}_Y) \rightarrow \mathcal{O}_y$ and consequently the composition $\Gamma(Y, \mathcal{O}_Y) \rightarrow \widehat{\mathcal{O}}_y$.

Popescu's theorem ([P1],[P2] and [P3]) states that a map between two Noetherian rings $A \rightarrow B$ is regular if and only if B is a filtered colimit of finitely generated smooth A -algebras. Since the map $\mathcal{O}_Y \rightarrow \widehat{\mathcal{O}}_y$ is regular, we get that $\widehat{\mathcal{O}}_y$ can be obtained as a filtered colimit $\varinjlim A_i$ of smooth finitely generated $\mathcal{O}_Y$ -algebras. Functor Spec sends colimits to limits, so we get that $\text{Spec } \widehat{\mathcal{O}}_y = \varprojlim \text{Spec } A_i$.

Let's now assume that the pull-back of $\mathcal{D}_{X,\alpha}$ to $\mathrm{Spec} \widehat{\mathcal{O}}_y^{(1)} \times_{Y^{(1)}} X^{(1)}$ splits. We have

$$\mathrm{Spec} \widehat{\mathcal{O}}_y^{(1)} \times_{Y^{(1)}} X^{(1)} \simeq (\varprojlim \mathrm{Spec} A_i^{(1)}) \times_{Y^{(1)}} X^{(1)} \simeq \varprojlim (\mathrm{Spec} A_i^{(1)} \times_{Y^{(1)}} X^{(1)})$$

and since $\mathcal{D}_{X,\alpha}$ is a coherent sheaf of algebras, it splits over some $\mathrm{Spec} A_i^{(1)} \times_{Y^{(1)}} X^{(1)}$ as well. Smooth map $\mathrm{Spec} A_i \rightarrow Y$ admits an étale quasi-section: there exists a surjection $A_i[f^{-1}] \twoheadrightarrow B_i$ for some $\mathcal{O}_Y$ -algebra B_i and a function $f \in \mathcal{O}_Y$, such that the corresponding map $\mathrm{Spec} B_i \rightarrow Y$ is étale. This is the étale neighbourhood we are looking for: $\mathrm{Spec} B_i \hookrightarrow \mathrm{Spec} A_i$ and $\mathcal{D}_{X,\alpha}$ splits on $\mathrm{Spec} B_i^{(1)} \times_{Y^{(1)}} X^{(1)}$.

So it is enough to prove that the pull-back of $\mathcal{D}_{X,\alpha}$ to $\mathrm{Spec} \widehat{\mathcal{O}}_y^{(1)} \times_{Y^{(1)}} X^{(1)}$ splits. But this was already done in Proposition 3.2.2. $\square$

4. OBSTRUCTIONS TO GLOBAL DESCENT

As we saw in Theorem 3.2.1, if $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$, the classes $[\mathcal{D}_{X,\alpha}]$ descend to $Y^{(1)}$ étale locally for all α . The way to prove it was to split them étale locally on $Y^{(1)}$, namely we proved that the pull-back of $\mathcal{D}_{X,\alpha}$ to $(U \times_Y X)^{(1)}$ splits for some étale cover $U \rightarrow Y$. Obviously this kind of argument will not help in the global situation, unless the algebra $\mathcal{D}_{X,\alpha}$ is globally split. Nevertheless one of the possible ways to proceed further could be to try to study the obstructions to global descent. In this section we define two different types of obstructions.

4.1. Picard obstruction. From the local-to-global principle it is natural to expect that obstructions to global descent should lie in some cohomology group of some natural sheaf on Y . It is true indeed and the corresponding sheaf is easy to describe, but on the other hand it seems to be almost impossible to work with. This approach does not take into account that our classes come from differential forms.

Let $\mathcal{A}$ be an Azumaya algebra on X and let $[\mathcal{A}]$ be its class in $\mathrm{Br}(X) = H^2(X_{\mathrm{\acute{e}t}}, \mathbb{G}_m)$. We have a natural map $\varphi : \mathrm{Br}(X) = H^2(X_{\mathrm{\acute{e}t}}, \mathbb{G}_m) \rightarrow H^0(Y_{\mathrm{\acute{e}t}}, R^2 \pi_* \mathbb{G}_m)$. It is easy to see that $\varphi([\mathcal{A}]) = 0$ if and only if $\mathcal{A}$ splits on $U \times_Y X$ for some étale cover $U \rightarrow Y$. In particular Theorem 3.2.1 says exactly that $\varphi([\mathcal{D}_{X,\alpha}]) = 0$.

Note that $R^0 \pi_* \mathbb{G}_m = \mathbb{G}_m$ and $R^1 \pi_* \mathbb{G}_m = \underline{\mathrm{Pic}}_{X/Y}$. From Leray spectral sequence $H_{\mathrm{\acute{e}t}}^p(Y, R^q \pi_* \mathbb{G}_m) \Rightarrow H_{\mathrm{\acute{e}t}}^{p+q}(X, \mathbb{G}_m)$, we get a short exact sequence

$$0 \rightarrow \mathrm{Br}(Y) \rightarrow \mathrm{Ker}(\varphi) \rightarrow H_{\mathrm{\acute{e}t}}^1(Y, \underline{\mathrm{Pic}}_{X/Y}) \rightarrow 0$$

and if $\varphi([\mathcal{A}]) = 0$ we get a well-defined class $[\widetilde{\mathcal{A}}]$ in $H_{\mathrm{\acute{e}t}}^1(Y, \underline{\mathrm{Pic}}_{X/Y})$, which is zero if and only if $[\mathcal{A}]$ lies in the image of $\mathrm{Br}(Y)$. We call this class the *Picard obstruction* corresponding to $\mathcal{A}$.

Unfortunately the only thing we can say about this class in general is that $[\widetilde{\mathcal{A}}]$ lies in the p -torsion subgroup $H_{\mathrm{\acute{e}t}}^1(Y, \underline{\mathrm{Pic}}_{X/Y}[p])$. In particular we do not know when this subgroup is zero, unless $Y = \mathrm{Spec} \mathbb{k}$, in that case we obtain the statement of Proposition 3.2.1.

4.2. Sheaves $\mathcal{Q}_{\pi,N}$. This class of obstructions works only for affine $Y = \mathrm{Spec} A$ and a proper map $\pi : X \rightarrow Y$ from a smooth variety X that satisfies $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$. We will construct a sequence of sheaves $\mathcal{Q}_{\pi,N}$ on $Y^{(N)}$ (or equivalently a sequence of A^{p^N} -modules) endowed with natural maps $\psi_N : (\mathrm{Fr}_Y^N)_* \pi_* \Omega_X^1 \rightarrow \mathcal{Q}_{\pi,N}$ such that if $\psi_N(\alpha^{(N-1)}) = 0$ for some N then the class $[\mathcal{D}_{X,\alpha}]$ descends to $Y^{(1)}$.

The first idea is to functorially extend the map $c_X : \Omega^1(X^{(1)}) \rightarrow \mathrm{Br}(X^{(1)})$ to the case of singular varieties, in particular Y . This can be done indeed, but only with the help of some

homotopy algebra, which we decided not to use (at least in this paper). Interested reader will find more details about that in Remark 4.2.2.

We use the following simple, but completely satisfactory for our purposes, lemma instead.

Lemma 4.2.1. *Let $\pi : X \rightarrow Y$ be a map from a smooth variety X to an affine variety Y . Let $\alpha \in H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ be a 1-form and suppose that there exists a Kähler differential θ on $Y^{(1)}$, such that $\alpha = \pi^*(\theta)$. Then the class $[\mathcal{D}_{X,\alpha}] \in \text{Br}(X^{(1)})$ descends to $Y^{(1)}$.*

Proof. Let ι be an embedding of $Y^{(1)}$ to a smooth variety $Z^{(1)}$, such that $\theta = \iota^*\theta_Z$ for some 1-form θ_Z on Z . If θ is represented in the form $\sum f_i dg_i$ then for ι one can take the map to the big enough affine space given by all f_i and g_i . We define an element $c_Y(\theta) \in \text{Br}(Y^{(1)})$ (this is just a notation, the map c_Y is not a priori defined since Y is not necessarily smooth, in particular in our definition it a priori depends on ι) by $c_Y = \iota^*c_Z(\theta_Z)$. Then

$$\pi^*c_Y(\theta) = \pi^*\iota^*c_Z(\theta_Z) = (\iota \circ \pi)^*c_Z(\theta_Z) = c_X((\iota \circ \pi)^*\theta_Z) = c_X(\pi^*\theta).$$

So the class $[\mathcal{D}_{X,\alpha}]$ descends to Y and we are done. $\square$

Remark 4.2.1. In particular, since $\mathcal{O}_Y \xrightarrow{\sim} \pi_*(\mathcal{O}_X)$ the class $[\mathcal{D}_{X,df}]$ for $f \in H^0(X^{(1)}, \mathcal{O}_{X^{(1)}})$ always descends to $Y^{(1)}$.

Remark 4.2.2. One can show that the class $c_Y(\theta) \in \text{Br}(Y)$ does not depend on the choice of the embedding ι and that this construction gives a well-defined map from the space of Kähler differentials $H^0(Y^{(1)}, \Omega_{Y^{(1)}}^1)$ to the Brauer group $\text{Br}(Y^{(1)})$. With some modifications this kind of definition can also be extended to the case of arbitrary non-smooth X which is not affine. Namely one can construct a well-defined map $c_X : \mathbb{H}^0(X^{(1)}, \mathbb{L}_{X^{(1)}}^\bullet) \rightarrow \text{Br}(X^{(1)})$ from the 0-th hyper-cohomology group of the cotangent complex of X . Moreover this map will be functorial in X . Note that $\mathbb{L}_X^\bullet$ has cohomology sheaves in negative degrees and a priori $\mathbb{H}^0(X, \mathbb{L}_X^\bullet)$ is not isomorphic to the space of global Kähler differentials $H^0(X, \Omega_X^1)$ (unless X is smooth or affine).

Now we return to our setting of affine $Y = \text{Spec } A$ and a proper map $\pi : X \rightarrow Y$ from a smooth variety X with $\mathcal{O}_Y \xrightarrow{\sim} \pi_*\mathcal{O}_X$ and $R^1\pi_*\mathcal{O}_X = R^2\pi_*\mathcal{O}_X = 0$. We have a natural A -module structure on $H^0(X, \Omega_X^1)$ given by multiplication of a differential form by a function, and, since π is proper, the module $H^0(X, \Omega_X^1) = H^0(Y, \pi_*\Omega_X^1)$ is finitely generated over A . Recall (see Remark 3.2.1) that we have an injective map $\pi^* : \text{Br}(Y^{(1)}) \rightarrow \text{Br}(X^{(1)})$. Let $\text{Br}(X^{(1)}/Y^{(1)})$ denote the quotient $\text{Br}(X^{(1)})/\text{Br}(Y^{(1)})$ and let $c_{X/Y} : H^0(X^{(1)}, \Omega_{X^{(1)}}^1) \rightarrow \text{Br}(X^{(1)}/Y^{(1)})$ be the map obtained from c_X by composition with the natural projection $\text{Br}(X^{(1)}) \rightarrow \text{Br}(X^{(1)}/Y^{(1)})$. It is clear that the class $[\mathcal{D}_{X,\alpha}]$ descends to Y if and only if $c_{X/Y}(\alpha) = 0$. Note that by Corollary 2.8.1 and Lemma 4.2.1

- $\mathbb{F}_p$ -vector subspace spanned by $\omega^{(1)} - C(\omega)$ for closed 1-forms ω on X ;
- A -submodule generated by $\pi^*(\theta)$ for Kähler differentials θ on $Y^{(1)}$.

lie in the kernel of $c_{X/Y}$. Let $S_\pi = \langle \pi^*(\theta), \omega^{(1)} - C(\omega) \rangle_{\theta, \omega}$ be the $\mathbb{F}_p$ -subspace of $H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ generated by these two types of 1-forms and let Q_π denote quotient $H^0(X^{(1)}, \Omega_{X^{(1)}}^1)/S_\pi$. We would like to show that $Q_\pi = 0$, in this case we would get that $[\mathcal{D}_{X,\alpha}]$ descends to $Y^{(1)}$ for any α . If Q_π had a structure of A -module it would define a coherent sheaf $\mathcal{Q}_\pi$ on $Y^{(1)}$ and to prove that $Q_\pi = 0$ it would be enough to show that all stalks of $\mathcal{Q}_\pi$ are zero. However, there is no structure on Q_π except that of an $\mathbb{F}_p$ -vector space.

So we try to be more clever and filter S_π by a system of subspaces $S_{\pi,0} \subset S_{\pi,1} \subset \dots \subset S_\pi$, such that $S_{\pi,N}$ has a module structure over $(A)^{p^N}$, or, in other words, $S_{\pi,N}$ defines a subsheaf $\mathcal{S}_{\pi,N} \subset (\text{Fr}_X^N)_*\pi_*\Omega_X^1$. The strategy then is to prove that $S_{\pi,N} = H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ for N big enough and that consequently $Q_\pi = 0$. Let's now define the sheaves $\mathcal{S}_{\pi,N}$.

The starting point is the following interesting observation: let $\Omega_{Y,cl}^1 \subset \pi_*\Omega_{X,cl}^1$ be the natural embedding of closed Kähler differentials to closed 1-forms on X . Let $\pi_*C : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_{X^{(1)}}^1$ be the Cartier operator. Does π_*C map $\Omega_{Y,cl}^1 \subset \pi_*\Omega_{X,cl}^1$ to $\Omega_Y^1 \subset \pi_*\Omega_X^1$? If the Cartier operator were defined for singular varieties, this would be true by functoriality. Surprisingly, usually this is not true at all, moreover in some cases (of the most interest for us) we can prove that all differential 1-forms on X can be obtained from Kähler differentials on Y by applying Cartier operator sufficiently many times.

Recall that by Lemma 3.2.2 Cartier operator $\pi_*C : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_{X^{(1)}}^1$ is surjective. We inductively define sheaves $\mathcal{C}_{\pi,n} \subset \pi_*\Omega_X^1$ by $\mathcal{C}_{\pi,0} = \Omega_X^1$ and $\mathcal{C}_{\pi,n} = C^{-1}(\mathcal{C}_{\pi,n-1}^{(1)})$, where $\mathcal{C}_{\pi,n-1}^{(1)}$ is the analogous subsheaf of $\pi_*\Omega_{X^{(1)}}^1$ for $\pi : X^{(1)} \rightarrow Y^{(1)}$. Sheaf $\mathcal{C}_{\pi,n}$ is the subsheaf of differential 1-forms in $\pi_*\Omega_X^1$ on which C^n is well-defined. Moreover from the p -linearity of Cartier operator it is easy to see that $(\text{Fr}_Y^n)_*\mathcal{C}_{\pi,n}$ defines a coherent subsheaf of $(\text{Fr}_Y^n)_*\pi_*\Omega_X^1$. So, in particular $\mathcal{C}_{\pi,0} = \pi_*\Omega_X^1$, $\mathcal{C}_{\pi,1} = \pi_*\Omega_{X,cl}^1$ and in general we have a sequence of surjections

$$(\text{Fr}_Y^n)_*\mathcal{C}_{\pi,n} \xrightarrow{C} (\text{Fr}_Y^{n-1})_*\mathcal{C}_{\pi,n-1}^{(1)} \xrightarrow{C} \dots \xrightarrow{C} (\text{Fr}_Y)_*\pi_*\Omega_{X^{(n-1)},cl}^1 \xrightarrow{C} \pi_*\Omega_{X^{(n)}}^1$$

of coherent sheaves on $Y^{(n)}$.

Let's now consider $\Omega_Y^1 \subset \pi_*\Omega_X^1$ and the intersection $\Omega_Y^1 \cap \mathcal{C}_{\pi,n}$. We have a natural map $C^n : \Omega_Y^1 \cap \mathcal{C}_{\pi,n} \rightarrow \pi_*\Omega_{X^{(n)}}^1$ and we define $\mathcal{S}_{\pi,n}$ as the image of this map. Note that $\mathcal{S}_{\pi,n-1}^{(1)} \subset \pi_*\Omega_{X^{(n)}}^1$ is embedded into $\mathcal{S}_{\pi,n}$, because $C(f^p g^{p-1} dg) = f^{(1)} dg^{(1)}$ and so for any $\theta \in \Omega_Y^{(1)}$ there exists $\theta' \in \Omega_Y^1$, such that $C(\theta') = \theta$. Using $\bullet^{(n-1)}$ we can identify $H^0(Y^{(n)}, \pi_*\Omega_{X^{(n)}}^1) = H^0(X^{(n)}, \Omega_{X^{(n)}}^1)$ with $H^0(X^{(1)}, \pi_*\Omega_{X^{(1)}}^1)$ (as $\mathbb{F}_p$ -vector spaces) and images of $H^0(Y^{(n)}, \mathcal{S}_{\pi,n}) \subset H^0(X^{(n)}, \Omega_{X^{(n)}}^1)$ for different n give the desired inductive system $\mathcal{S}_{\pi,0} \subset \mathcal{S}_{\pi,1} \subset \dots$

Let's now suppose that we have $\alpha \in \mathcal{S}_{\pi,N}$ for some N . Unraveling the definitions, this means that we can find the set of 1-forms $\omega_i \in H^0(X, \Omega_X^1)$ and a Kähler differential $\theta \in H^0(Y, \Omega_{Y^{(1)}}^1)$ that satisfy the following system of equations:

$$\left\{ \begin{array}{l} d\omega_i = 0, \forall i = 1, \dots, N \\ C\omega_1 = \alpha \\ C\omega_2 = \omega_1^{(1)} \\ \dots \\ C\omega_i = \omega_{i-1}^{(1)} \\ \dots \\ C\omega_N = \omega_{N-1}^{(1)} \\ \pi^*\theta = \omega_N^{(1)}, \theta \in H^0(Y, \Omega_{Y^{(1)}}^1) \end{array} \right.$$

Taking the sum of all equalities we get that $\pi^*\theta - \alpha = (\bullet^{(1)} - C)(\sum_{i=1}^N \omega_i)$. This means that $\alpha \in \mathcal{S}_{\pi}$ and the class $[\mathcal{D}_{X,\alpha}]$ descends to $Y^{(1)}$.

Definition 4.2.1. For $N \geq 0$ we define the obstruction sheaf $\mathcal{Q}_{\pi,N}$ as the quotient $\pi_*\Omega_{X^{(N)}}^1 / \mathcal{S}_{\pi,N}$. It is a coherent sheaf on $Y^{(N)}$ and since Y is affine we have a short exact sequence

$$0 \rightarrow \mathcal{S}_{\pi,N} \rightarrow H^0(X^{(1)}, \pi_*\Omega_{X^{(1)}}^1) \xrightarrow{\psi_N} \mathcal{Q}_{\pi,N} \rightarrow 0,$$

where $\mathcal{Q}_{\pi,N}$ is the twist $H^0(Y^{(N)}, \mathcal{Q}_{\pi,N})^{(1-N)}$ of the global sections of $\mathcal{Q}_{\pi,N}$. From the discussion above: if $\psi_N(\alpha) = 0$ for some N , the class $[\mathcal{D}_{X,\alpha}]$ descends to $Y^{(1)}$.

The following lemma is very important and gives a motivation for the upcoming definition of a resolution with conical slices (in Sect. 5):

Lemma 4.2.2. *Let $\pi : X \rightarrow Y$ be a resolution of singularities such that $\mathcal{Q}_{\pi,N} = 0$. Let S be an arbitrary smooth affine variety, then for $\pi \times \text{id}_S : X \times S \rightarrow Y \times S$ we have $\mathcal{Q}_{\pi \times \text{id}_S, N} = 0$.*

Proof. The statement of the lemma amounts to saying that for any differential 1-form $\alpha \in \Omega^1(X \times S)$, there is a solution to the system (4.2). We now use the formula

$$\Omega_{X \times S}^1 = \Omega_X^1 \boxtimes \mathcal{O}_S \oplus \mathcal{O}_X \boxtimes \Omega_S^1$$

to write α as a sum of 1-forms of the following kind: $\alpha_X \boxtimes f_S + f_X \boxtimes \alpha_S$. Moreover we can assume that there is only one summand, so that

$$\alpha = \alpha_X \boxtimes f_S + f_X \boxtimes \alpha_S.$$

By assumption, there is a solution $(\omega_{1,X}, \dots, \omega_{N,X}, \theta_Y)$ to the system (4.2) with α replaced by α_X . Since S is smooth and affine, $\mathbb{C} : \Omega_{S,cl}^1 \rightarrow \Omega_{S(1)}^1$ is surjective and induces a surjective map on global sections. So (4.2) must also have a solution for α_S (and id_S instead of π). We denote the corresponding sequence of 1-forms on S by $(\omega_{1,S}, \dots, \omega_{N,S}, \theta_S = \omega_{N,S})$. Also, using the assumption $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$, we can descend f_X to Y , namely find $f_Y \in \Gamma(Y, \mathcal{O}_Y)$ such that $f_X = \pi^* f_Y$. We now take

$$\omega_i = \omega_{i,X} \boxtimes f_S^{p_i} + f_X^{p_i} \boxtimes \omega_{i,S}, \quad \theta = \theta_Y \boxtimes f_S^{p_N} + f_Y^{p_N} \boxtimes \theta_S.$$

Using $\mathbb{C}(f^p \omega) = f^{(1)} \mathbb{C}(\omega)$ we see that the resulting sequence $(\omega_1, \dots, \omega_N, \theta)$ solves the system (4.2) for the form α . □

5. RESOLUTIONS WITH CONICAL SLICES

5.1. Setup. We would like to have a class of resolutions for which we could effectively apply Lemma 4.2.2. Namely, it would be great if a neighbourhood of any point $y \in Y$ could be decomposed as a product of something smooth and some other resolution $\pi' : X' \rightarrow Y'$ for which we already knew that $\mathcal{Q}_{\pi',N} = 0$ for some N . A good candidate for this is a conical resolution: if there is a contracting $\mathbb{G}_m$ -action on $\pi : X \rightarrow Y$, then, by Luna's slice theorem, étale locally at any non-central point, π can be decomposed as a product of the $\mathbb{G}_m$ -orbit and some slice $\pi' : X' \rightarrow Y'$. However we will not know much about π' , in particular there is no need for π' to be again conical. It turns out though that if we add by force the condition for this slice to be conical (and some other mild technical conditions), everything will work out relatively well: see Sect. 7 and Theorem 7.2.2 in particular. The condition on the slice to be conical can be elegantly reformulated as the existence of another contracting $\mathbb{G}_m$ -action in an étale neighbourhood of each point: see Definition 5.4.3 and Proposition 5.5.1 for the relation to slices. We call a resolution satisfying this a *resolution with conical slices*.

5.2. Category of étale germs of resolutions. In this subsection we construct what we call *the category of étale germs of resolutions*. The idea is to identify resolutions which look the same in the neighbourhood of some point of the base (where the point is a part of the data). We could probably consider something like the category of the resolution with a henselian base, but instead we just localise the category of all resolutions by the class of morphisms which are étale. We follow probably the most classical approach to localisation of categories, namely we prove that the class of étale maps satisfies the right Ore condition. In particular, as a result we get a well defined and quite natural notion of an étale equivalence between resolutions of singularities with a chosen point on the base.

Definition 5.2.1. By a *resolution (of singularities)* $\pi : X \rightarrow Y$ we mean a proper birational map from a smooth variety X to a possibly singular variety Y .

Remark 5.2.1. Please note that Y does not actually need to be singular in the definition, so in this sense our definition of resolution can differ from the more standard ones.

Definition 5.2.2. A *pointed resolution* (π, X, Y, y) is the data of a resolution $\pi : X \rightarrow Y$ and a distinguished point $y \in Y(\mathbb{k})$.

We define Res_* to be the category with objects given by pointed resolutions (π, X, Y, y) , and morphisms $\text{Mor}_{\text{Res}_*}((\pi', X', Y', y'), (\pi, X, Y, y))$ given by a pair of maps $p : Y' \rightarrow Y$ and $q : X' \rightarrow X$, such that $p(y') = y$ and the following square is commutative:

$$\begin{array}{ccc} X' & \xrightarrow{q} & X \\ \pi' \downarrow & & \downarrow \pi \\ Y' & \xrightarrow{p} & Y \end{array}$$

We would like to identify two pointed resolutions, which become isomorphic after taking some étale neighbourhoods of distinguished points. For this we can try to localise the category Res_* by the set of morphisms f for which the corresponding p and q are étale. This simple lemma about étale morphisms will appear to be very useful in this context.

Lemma 5.2.1. *Let $f : (\pi, X, Y, y) \rightarrow (\pi', X', Y', y')$ be a morphism such that the corresponding morphisms p and q are étale. Then $X' = Y' \times_Y X$ and the corresponding square is Cartesian.*

Proof. By definition we have a commutative diagram

$$\begin{array}{ccc} X' & \xrightarrow{q} & X \\ \pi' \downarrow & & \downarrow \pi \\ Y' & \xrightarrow{p} & Y \end{array}$$

which gives a natural map $\iota : X' \rightarrow Y' \times_Y X$.

$$\begin{array}{ccccc} X' & & & & \\ & \searrow \iota & & \nearrow q & \\ & & Y' \times_Y X & \xrightarrow{\tilde{p}} & X \\ & \searrow \pi & \downarrow \tilde{\pi} & & \downarrow \pi \\ & & Y' & \xrightarrow{p} & Y \end{array}$$

We have a decomposition $q = \tilde{p} \circ \iota$, where $\tilde{p} : Y' \times_Y X \rightarrow X$ is the pull-back of p to X . The map $\tilde{p}$ is étale as a pull-back of p , q is étale by assumption, therefore we get that ι is étale too.

Let $Y_0 \subset Y$ and $Y'_0 \subset Y'$ be the smooth loci where π and π' are 1-to-1, and let $j : Y_0 \rightarrow X$, $j' : Y'_0 \rightarrow X'$ be the corresponding embeddings. We have two open embeddings $\tilde{j} : Y'_0 \times_Y Y_0 \rightarrow Y' \times_Y X$ and $j : Y'_0 \times_Y Y_0 \rightarrow X'$. Both Y_0 and Y'_0 are open and dense in X and X' , so $Y'_0 \times_Y Y_0$ is open and dense in both X' and $Y' \times_Y X$. Note that the restriction of ι to $Y'_0 \times_Y Y_0$ is just the identity map. We also have that ι should necessarily be surjective, since $\iota(X') = \iota(Y'_0 \times_Y Y_0) = \overline{Y'_0 \times_Y Y_0} = Y' \times_Y X$. Summarising the above, we get that ι is étale, surjective, and is an isomorphism on some dense open set, therefore it is an isomorphism on the whole and the corresponding square is Cartesian. $\square$

Definition 5.2.3. $\text{Et}((\pi, X, Y, y), (\pi', X', Y', y')) \subset \text{Mor}_{\text{Res}_*}((\pi, X, Y, y), (\pi', X', Y', y'))$ is defined as the subset of morphisms for which p and q are étale.

Let Sch denote the category of schemes over $\mathbb{k}$ and let $\text{Ar}_*(\text{Sch})$ be the category of pointed arrows in Sch . Namely objects of $\text{Ar}_*(\text{Sch})$ are maps $f : X \rightarrow Y$ with a distinguished point $y \in Y(\mathbb{k})$ and morphisms between two such arrows (f, X, Y, y) and (g, X', Y', y') are given by commutative squares

$$\begin{array}{ccc} X & \xrightarrow{q} & X' \\ f \downarrow & & \downarrow g \\ Y & \xrightarrow{p} & Y' \end{array}$$

such that $p(y) = y'$. It is easy to see that Res_* is a full subcategory of $\text{Ar}_*(\text{Sch})$.

The category $\text{Ar}_*(\text{Sch})$ has fiber products: if $A = (f, X, Y, y)$, $B_1 = (g_1, Z_1, W_1, w_1)$, $B_2 = (g_2, Z_2, W_2, w_2)$ and $f_1 : B_1 \rightarrow A$, $f_2 : B_2 \rightarrow A$ are two morphisms, then the corresponding fiber product $B_1 \times_A B_2$ is given by the map $g_1 \times g_2 : Z_1 \times_X Z_2 \rightarrow W_1 \times_Y W_2$ with a distinguished point $w_1 \times w_2 \in (W_1 \times_Y W_2)(\mathbb{k})$. However Res_* does not have fiber products in general, the problem is that $\text{Res}_* \subset \text{Ar}_*(\text{Sch})$ is not closed under the fiber product in $\text{Ar}_*(\text{Sch})$. To illustrate this we can take some smooth Y , and let $A = (\text{id}, Y, Y, y)$ be the identity map on Y , $B_1 = (\pi, \text{Bl}_y(Y), Y, y)$ the blow-up at the point y and $B_2 = (\text{id}, \text{Spec } \mathbb{k}, \text{Spec } \mathbb{k}, \text{Spec } \mathbb{k})$ the identity map for a point, with the maps f_1 and f_2 given by the projection $\text{Bl}_y(Y) \rightarrow Y$ and the point $y : \text{Spec } \mathbb{k} \rightarrow Y$ correspondingly. Then the fiber product $B_1 \times_A B_2$ is given by $\pi : \pi^{-1}(\text{Bl}_y(Y)) \rightarrow \text{Spec } \mathbb{k}$, which is not birational if $\dim Y \geq 2$. But for some pairs f_1, f_2 fiber product still exists, in particular when one of the morphisms lies in Et :

Lemma 5.2.2. *Let $A = (\pi, X, Y, y)$, $B_1 = (\eta_1, Z_1, W_1, w_1)$, $B_2 = (\eta_2, Z_2, W_2, w_2)$ and f_1, f_2 be objects and a pair of morphisms in Res_* :*

$$\begin{array}{ccc} B_1 & & B_2 \\ & \searrow f_1 & \swarrow f_2 \\ & A & \end{array}$$

Assume that f_2 lies in $\text{Et}(B_2, A)$. Then the fiber product $B_1 \times_A B_2$ exists and is given by $(\eta_1 \times \eta_2, Z_1 \times_X Z_2, W_1 \times_Y W_2, w_1 \times w_2)$.

Proof. We have the following diagram:

$$\begin{array}{ccccc} & Z_1 \times_X Z_2 & \xrightarrow{\tilde{q}_1} & Z_2 & \\ & \downarrow \eta_1 \times \eta_2 & & \downarrow \eta_2 & \\ Z_1 & \xrightarrow{q_1} & X & \xrightarrow{\pi} & W_2 \\ \downarrow \eta_1 & & \downarrow \pi & & \downarrow \eta_2 \\ & W_1 \times_Y W_2 & \xrightarrow{\tilde{p}_1} & W_2 & \\ & \downarrow \tilde{p}_2 & & \downarrow p_2 & \\ W_1 & \xrightarrow{p_1} & Y & & \end{array}$$

The quadruple $(\eta_1 \times \eta_2, Z_1 \times_X Z_2, W_1 \times_Y W_2, w_1 \times w_2)$ is the fiber product of (η_1, Z_1, W_1, w_1) and (η_2, Z_2, W_2, w_2) over (π, X, Y, y) in the category of arrows in the category of schemes Sch.

So the only thing we need to check is that

$$\eta_1 \times \eta_2 : Z_1 \times_X Z_2 \rightarrow W_1 \times_Y W_2$$

is a resolution of singularities. By Lemma 5.2.1 Z_2 is equal to $X \times_Y W_2$, so

$$Z_1 \times_X Z_2 = Z_1 \times_X X \times_Y W_2 = Z_1 \times_Y W_2 = Z_1 \times_{W_1} (W_1 \times_Y W_2),$$

Moreover the bottom, right and top faces of the cube are cartesian squares and it is easy to see that the left face is a cartesian square too. Therefore, $\eta_1 \times \eta_2$ is just the pull-back of η_1 under $\tilde{p}_2$, which is étale (as a pull-back of étale map). We see that $\eta_1 \times \eta_2$ is also a resolution. $\square$

Remark 5.2.2. Note that $\tilde{f}_2 : B_1 \times_A B_2 \rightarrow B_1$ lies in $\text{Et}(B_1 \times_A B_2, B_1)$. So étale maps are closed under pull-backs in Res_* as well.

Remark 5.2.3. There is another important example when the fiber product exists, namely when $A = \underline{\text{Spec}} \mathbb{k} = (\text{id}, \underline{\text{Spec}} \mathbb{k}, \underline{\text{Spec}} \mathbb{k}, \underline{\text{Spec}} \mathbb{k})$. It is easy to see that $\underline{\text{Spec}} \mathbb{k}$ is a final object of Res_* and so the corresponding fiber product is just the product. It is given by the quadruple $B_1 \times B_2 = (\pi_1 \times \pi_2, Z_1 \times Z_2, W_1 \times W_2, w_1 \times w_2)$ which is a resolution of singularities if π_1, π_2 are.

Lemma 5.2.3. *Collection of subsets $\text{Et}((\pi, X, Y, y), (\pi', X', Y', y'))$ inside the sets of morphisms $\text{Mor}_{\text{Res}_*}((\pi, X, Y, y), (\pi', X', Y', y'))$ satisfies the right Ore condition.*

Proof. This follows essentially from Lemma 5.2.2: having the diagram

$$\begin{array}{ccc} B_1 & & B_2 \\ & \searrow f_1 & \swarrow f_2 \\ & A & \end{array}$$

such that f_2 lies in $\text{Et}(B_2, A)$ we can form another diagram


$$\begin{array}{ccc} & B_1 \times_A B_2 & \\ \tilde{f}_2 \swarrow & & \searrow \tilde{f}_1 \\ B_1 & & B_2 \end{array}$$

such that $\tilde{f}_2$ lies in $\text{Et}(B_1 \times_A B_2, B_1)$. $\square$


Last lemma allows us to localise Res_* at the class of morphisms Et . We call the result of localisation Res_*^{et} the category of étale germs of resolutions. We remind ([GM], III.2, Lemma 8) that the set of morphisms in Res_*^{et} between objects B_1 and B_2 is given by classes of equivalences of diagrams

$$\begin{array}{ccc} & A & \\ f_2 \swarrow & & \searrow f_1 \\ B_1 & & B_2 \end{array}$$

where f_2 is étale. An equivalence between pairs (f_1, f_2) and (g_1, g_2) is given by an object A'' and a pair of morphisms $h_1 \in \text{Et}(A'', A')$, $h_2 \in \text{Et}(A'', A)$, such that the following diagram commutes:



The composition $(f_1, f_2) \circ (g_1, g_2)$ is given by the fiber product:



Morphisms in $\mathbf{Res}_*^{\text{et}}$ that we are interested in the most are isomorphisms. We call them *étale equivalences*.

Definition 5.2.4. Let (π, X, Y, y) and (π', X', Y', y') be two pointed resolutions. We say that (π, X, Y, y) is *étale equivalent* to (π', X', Y', y') if they are isomorphic as objects of $\mathbf{Res}_*^{\text{et}}$. Unwinding the definitions it means that there exists a pointed resolution (π'', X'', Y'', y'') and a correspondence $(\pi, X, Y, y) \xleftarrow{f_1} (\pi'', X'', Y'', y'') \xrightarrow{f_2} (\pi', X', Y', y')$ (given by correspondences $Y \xleftarrow{p_1} Y'' \xrightarrow{p_2} Y'$, and $X \xleftarrow{q_1} X'' \xrightarrow{q_2} X'$, $p_1(y'') = y$, $p_2(y'') = y'$ such that the diagram

$$\begin{array}{ccccc}
 X & \xleftarrow{q_1} & X'' & \xrightarrow{q_2} & X' \\
 \pi \downarrow & & \pi'' \downarrow & & \pi' \downarrow \\
 Y & \xleftarrow{p_1} & Y'' & \xrightarrow{p_2} & Y'
 \end{array}$$

is commutative) with maps p_1, p_2, q_1, q_2 being étale. We denote the equivalence by $(\pi, X, Y, y) \sim_{\text{et}} (\pi', X', Y', y')$.

Remark 5.2.4. Note that by Lemma 5.2.2 we necessarily have $X \times_Y Y'' = X'' = Y'' \times_Y X'$.

5.3. Separable $\mathbb{G}_m$ -action. Let $\mathbb{k}$ be an algebraically closed field (not necessarily of finite characteristic), let X be a variety over $\mathbb{k}$ (possibly singular), endowed with a $\mathbb{G}_m$ -action. We have a natural non-zero tangent vector $\partial_t \in T_{\{1\}}\mathbb{G}_m$. For any point $x \in X(\mathbb{k})$ we have the differential of the action map $a_x : T_{\{1\}}\mathbb{G}_m \rightarrow T_x X$, which extends to the map of sheaves $a : \mathcal{O}_X \rightarrow \text{Der}(\mathcal{O}_X)$, where $\text{Der}(\mathcal{O}_X)$ is the sheaf of $\mathbb{k}$ -linear derivations of $\mathcal{O}_X$. Consider the dual map $a^* : \Omega_X^1 \rightarrow \mathcal{O}_X$, where Ω_X^1 is the sheaf of Kähler differentials on X . The image $\mathcal{I}_{a^*} = \text{Im}(a^*) \subset \mathcal{O}_X$ is some sheaf of ideals on X .

Definition 5.3.1. The subscheme $X^{d\mathbb{G}_m} \subset X$ of *pseudo-fixed* points is the subscheme, defined by the sheaf of ideals $\mathcal{I}_{a^*}$. This is a $\mathbb{G}_m$ -invariant closed subscheme of X .

Remark 5.3.1. The subscheme $X^{d\mathbb{G}_m} \subset X$ can also be viewed in a more geometric way. The vector ∂_t defines a section $a(\partial_t)$ of the tangent bundle $TX \rightarrow X$ and the corresponding subscheme $\Gamma_{a(\partial_t)}$ in the total space of the tangent bundle TX . The subscheme of pseudo-fixed

points $X^{d\mathbb{G}_m} \subset X$ is then just the (scheme-theoretic) intersection of $\Gamma_{a(\partial_t)}$ and the zero-section $\Gamma_0 \subset TX$.

$X^{d\mathbb{G}_m}$ is the subscheme of points where the differential of the $\mathbb{G}_m$ -action is zero. If $\text{char } \mathbb{k} = 0$ it is true that a point is fixed if and only if the differential of the action is zero (e.g. see [Dr]), so $X^{d\mathbb{G}_m}(\mathbb{k}) = X^{\mathbb{G}_m}(\mathbb{k})$. But if $\text{char } \mathbb{k} \neq 0$ these two subsets can easily happen to be different.

Definition 5.3.2. The $\mathbb{G}_m$ -action on X is called *separable* if all pseudo-fixed points are fixed:

$$X^{d\mathbb{G}_m}(\mathbb{k}) = X^{\mathbb{G}_m}(\mathbb{k}).$$

The simplest example when this property can fail is given by the composition of any non-trivial $\mathbb{G}_m$ -action and $\text{Fr}_{\mathbb{G}_m} : \mathbb{G}_m \rightarrow \mathbb{G}_m$. Then the differential of the action is 0 at any point and $X^{d\mathbb{G}_m} = X$.

Another simple example to consider is the affine line $\mathbb{A}^1 = \text{Spec } k[z]$ with the action of $\mathbb{G}_m$, given by some character $n \in \mathbb{Z}$: $t \circ z = t^n z$. We claim that this action is separable if and only if $(n, p) = 1$. Indeed $a_0(\partial_t) = \partial_t(t^n)|_1 \cdot \partial_z = n\partial_z$. More generally a linear diagonalised $\mathbb{G}_m$ -action on an affine space $\mathbb{A}^n$ is separable if and only if p does not divide the weights of the coordinates.

Remark 5.3.2. If we have a contracting $\mathbb{G}_m$ -action on an affine variety $Y = \text{Spec } A$, then there is a way to guarantee that the action is separable, looking at the weights of the action. There exists a $\mathbb{G}_m$ -equivariant embedding $Y \hookrightarrow \mathbb{A}^n$ (with the contracting action on $\mathbb{A}^n$) mapping y_0 to 0. Then, if the weights of the coordinates for the action on $\mathbb{A}^n$ are not divisible by p , $(\mathbb{A}^n)^{d\mathbb{G}_m} = (\mathbb{A}^n)^{\mathbb{G}_m}$. We have an embedding $Y^{d\mathbb{G}_m} \hookrightarrow (\mathbb{A}^n)^{d\mathbb{G}_m} = (\mathbb{A}^n)^{\mathbb{G}_m}$ and so $Y^{d\mathbb{G}_m} = Y^{\mathbb{G}_m}$. In other words, if there exists a set of homogeneous generators of A with degrees not divisible by p , the action is separable.

5.4. Definition of a resolution with conical slices. Let $\mathbb{k}$ be an algebraically closed field. We first define what we mean by the conical resolution. Note that in our definition there is a separability assumption on the $\mathbb{G}_m$ -action, which make the definition different from the usual one in the case when the characteristic of $\mathbb{k}$ is positive.

Definition 5.4.1. Resolution $\pi : X \rightarrow Y$ is called *conical*, if it can be endowed with a separable $\mathbb{G}_m$ -action, contracting Y to a single fixed point $y_0 \in Y(\mathbb{k})$ which is then called the *central point* of the resolution.

Here we say that π can be endowed with a $\mathbb{G}_m$ -action if there is a $\mathbb{G}_m$ -action on X and $\mathbb{G}_m$ -action on Y intertwined by π .

Definition 5.4.2. Let $\pi : X \rightarrow Y$ be a resolution. We say that $y \in Y(\mathbb{k})$ has a *conical étale neighbourhood* if there exists (π', X', Y', y') , such that $\pi' : X' \rightarrow Y'$ is a conical resolution, y' is its central point and $(\pi, X, Y, y) \sim_{et} (\pi', X', Y', y')$.

Definition 5.4.3. A *resolution with conical slices* is a conical resolution $\pi : X \rightarrow Y$ such that $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$, $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$ and every point $y \in Y(\mathbb{k})$ has an étale conical neighbourhood.

Remark 5.4.1. Note that from the condition $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ it follows that Y is normal.

Lemma 5.4.1. *Let $\pi : X \rightarrow Y$ be a resolution with conical slices. Let $(\pi, X, Y, y) \sim_{et} (\pi', X', Y', y')$ be an étale conical neighbourhood of a point $y \in Y(\mathbb{k})$. Then $\pi' \mathcal{O}_{X'} = \mathcal{O}_{Y'}$ and $R^1 \pi'_* \mathcal{O}_{X'} = R^2 \pi'_* \mathcal{O}_{X'} = 0$.*

Proof. The étale equivalence $(\pi, X, Y, y) \sim_{et} (\pi', X', Y', y')$ is given by a commutative diagram

$$\begin{array}{ccccc} X & \xleftarrow{q_1} & X'' & \xrightarrow{q_2} & X' \\ \pi \downarrow & & \pi'' \downarrow & & \pi' \downarrow \\ Y & \xleftarrow{p_1} & Y'' & \xrightarrow{p_2} & Y' \end{array}$$

where all horizontal maps are étale. From the flat base change we get that $\pi''\mathcal{O}_{X''} = \mathcal{O}_{Y''}$ and $R^1\pi''_*\mathcal{O}_{X''} = R^2\pi''_*\mathcal{O}_{X''} = 0$. Applying the flat base change one more time we get that $\pi'\mathcal{O}_{X'} = \mathcal{O}_{Y'}$ and $R^1\pi'_*\mathcal{O}_{X'} = R^2\pi'_*\mathcal{O}_{X'} = 0$ on some open neighbourhood $U' \subset Y'$ of the point y' , such that U' is contained in $p_2(Y'')$. Since U' is open and contains y' , the map $\mathbb{G}_m \times U' \rightarrow Y'$, given by the $\mathbb{G}_m$ -action, is surjective (since the orbit of any point contains y' in its closure and so intersects U'). From this, using the natural $\mathbb{G}_m$ -equivariant structure on $\mathcal{O}_{X'}$ we get that $\pi'\mathcal{O}_{X'} = \mathcal{O}_{Y'}$ and $R^1\pi'_*\mathcal{O}_{X'} = R^2\pi'_*\mathcal{O}_{X'} = 0$ on the whole Y' . $\square$

5.5. Construction of a slice. This subsection is devoted to the mentioned relation between Definition 5.4.3 and the existence of conical slices.

Let X be a variety over $\mathbb{k}$ and $x \in X(\mathbb{k})$. We denote by $\underline{X}_{\{x\}}$ the trivial pointed resolution (id_X, X, X, x) . We also define $\underline{\mathbb{A}}^1$ to be $\underline{\mathbb{A}}^1_{\{0\}}$ in the previous notation. Note that $\underline{\mathbb{A}}^1 \sim_{et} \underline{\mathbb{G}}_{m\{1\}}$, where equivalence is given by the composition of the embedding $\mathbb{G}_m \rightarrow \mathbb{A}^1$ and the translation by -1 . More generally, if X is of pure dimension n and x is a smooth point of X , $\underline{X}_{\{x\}} \sim_{et} (\underline{\mathbb{A}}^1)^n$. We also remind that if (ξ, Z, W, w) is a pointed resolution, then $(\xi, Z, W, w) \times \underline{\mathbb{A}}^1 = (\xi \times \text{id}, Z \times \mathbb{A}^1, W \times \mathbb{A}^1, w \times \{0\})$.

Lemma 5.5.1. *If there exists an étale equivalence $(\xi_1, Z_1, W_1, w) \times \underline{\mathbb{A}}^1 \sim_{et} (\xi_2, Z_2, W_2, w_2) \times \underline{\mathbb{A}}^1$, then there also exists an étale equivalence $(\xi_1, Z_1, W_1, w_1) \sim_{et} (\xi_2, Z_2, W_2, w_2)$.*

Proof. Let $\theta : T \rightarrow U$ with $u \in U(\mathbb{k})$ be a resolution which realizes the étale equivalence:

$$\begin{array}{ccccc} Z_1 \times \mathbb{A}^1 & \xleftarrow{q_1} & T & \xrightarrow{q_2} & Z_2 \times \mathbb{A}^1 \\ \downarrow \xi_1 \times \text{id} & & \theta \downarrow & & \downarrow \xi_2 \times \text{id} \\ W_1 \times \mathbb{A}^1 & \xleftarrow{p_1} & U & \xrightarrow{p_2} & W_2 \times \mathbb{A}^1 \end{array}$$

By definition, we have étale maps $W_1 \times \mathbb{A}^1 \xleftarrow{p_1} U \xrightarrow{p_2} W_2 \times \mathbb{A}^1$ and a point $u \in U(\mathbb{k})$ that maps to $w_1 \times \{0\}$ and $w_2 \times \{0\}$ respectively. For $i = 1, 2$ consider the natural isomorphisms $T_{w_i \times \{0\}}(W_i \times \mathbb{A}^1) \xrightarrow{\sim} T_{w_i}W_i \oplus T_{\{0\}}\mathbb{A}^1$. The maps p_1 and p_2 are étale, so for any point $z \in U(\mathbb{k})$ they give isomorphisms of Zariski tangent spaces $T_{p_1(z)}W_1 \oplus T_{\{0\}}\mathbb{A}^1 \cong T_zU \cong T_{p_2(z)}W_2 \oplus T_{\{0\}}\mathbb{A}^1$. We denote by $V_{z,1}$ and $V_{z,2}$ the two 1-dimensional subspaces of T_zU , which come as preimages of $T_{\{0\}}\mathbb{A}^1$ under these two isomorphisms. For any function $f : U \rightarrow \mathbb{A}^1$ and any $z \in U(\mathbb{k})$ we have two maps $g_{z,1}, g_{z,2}$, given by the restriction of $df_z : T_zU \rightarrow T_{f(z)}\mathbb{A}^1$ to subspaces $V_{z,1}$ and $V_{z,2}$. Consider the open subvariety $W_f \subset U$ of points z , such that $f(z) = 0$ and $g_{z,1}$ and $g_{z,2}$ are isomorphisms. By definition, both maps $\text{pr}_i \circ p_i|_{W_f} : W_f \rightarrow W_i$ (where pr_i is the projection $W \times \mathbb{A}^1 \rightarrow W_i$) are unramified at any point z of W_f , therefore are étale. Choosing f such that $f(u) = 0$ and $g_{u,1}, g_{u,2}$ are isomorphisms, we get that u is contained in $W_f(\mathbb{k})$. We get an étale correspondence $W_1 \xleftarrow{p_1} W_f \xrightarrow{p_2} W_2$, such that $p_i(u) = w_i$.

Now by Remark 5.2.4 the fiber products $Z_1 \times_{W_1} W_f$ and $Z_2 \times_{W_2} W_f$ are equal and give a resolution $\xi_f : Z_f \rightarrow W_f$. The natural maps $Z_1 \xleftarrow{q_1} Z_f \rightarrow q_2$ are étale since they are base

changes of p_1 and p_2 . We obtain an étale equivalence $(\xi_1, Z_1, W_1, w_1) \leftarrow (\xi_f, Z_f, W_f, u) \rightarrow (\xi_2, Z_2, W_2, w_2)$. $\square$

Lemma 5.5.2. *Let (ξ, Z, W, w) be a pointed resolution with an étale equivalence $(\xi, Z, W, w) \times \mathbb{A}^1 \sim_{et} (\pi', X', Y', y')$ where (π', X', Y', y') is conical. Then there exists another conical resolution (ξ', Z', W', w') with an étale equivalence $(\xi', Z', W', w') \sim_{et} (\xi, Z, W, w)$.*

Proof. Let $\theta : T \rightarrow U$ with $u \in U(\mathbb{k})$ be a resolution which realises the étale equivalence between $(\xi, Z, W, w) \times \mathbb{A}^1$ and (π', X', Y', y') :

$$\begin{array}{ccccc} Z \times \mathbb{A}^1 & \xleftarrow{q_1} & T & \xrightarrow{q_2} & X' \\ \downarrow \xi \times \text{id} & & \downarrow \theta & & \downarrow \pi' \\ W \times \mathbb{A}^1 & \xleftarrow{p_1} & U & \xrightarrow{p_2} & Y' \end{array}$$

We have a $\mathbb{G}_m$ -action on $W \times \mathbb{A}^1$, given by the trivial action on W and the dilation on $\mathbb{A}^1$. We want to find a slice to the orbit of w but in Y' , not in $W \times \mathbb{A}^1$. For this we consider the orbit $w \times \mathbb{G}_m \hookrightarrow W \times \mathbb{A}^1$. We look at its preimage $p_1^{-1}(w \times \mathbb{G}_m) \hookrightarrow U$ and take a connected component which contains y' in its image under p_2 . We denote by $S \subset Y'$ the image under p_2 . S is a subvariety of Y' of dimension 1, passing through y' . For any function $f : Y' \rightarrow \mathbb{A}^1$, such that $f(y') = 0$, we have a map $df|_{T_{y',S}} : T_{y',S} \rightarrow T_0\mathbb{A}^1$.

Recall that we have a $\mathbb{G}_m$ -action on Y' with a fixed point y' . So we also have a $\mathbb{G}_m$ -action on the ring $\mathcal{O}_{Y'}$ and, as a representation of $\mathbb{G}_m$, it decomposes as a direct sum over all possible characters: $\mathcal{O}_{Y'} = \bigoplus_{i \in \mathbb{Z}} V_i$, where $t(v) = t^i v$ for $v \in V_i$. Remember that Y' is conical, so we have $V_i = 0$ for $i < 0$ and $V_0 = \mathbb{k}$. Let $\mathfrak{m}_{y'} \subset \mathcal{O}_{Y'}$ be the maximal ideal corresponding to the point y' , then we also have an isomorphism of $\mathbb{G}_m$ -representations $\mathfrak{m}_{y'} \cong \bigoplus_{i > 0} V_i$. The surjective map to the Zariski cotangent space $\mathfrak{m}_{y'} \rightarrow \mathfrak{m}_{y'}/\mathfrak{m}_{y'}^2$ is $\mathbb{G}_m$ -equivariant. Now choose a homogeneous element in $\mathfrak{m}_{y'}/\mathfrak{m}_{y'}^2$, which is nonzero on $T_{y',S}$ (considered as a linear function on $T_{y',Y'}$). For that, take any possibly non-homogeneous element which is nonzero, then one of its homogeneous components is nonzero too. If we take any equivariant lift $f \in \mathfrak{m}_{y'}$ under the projection $\mathfrak{m}_{y'} \twoheadrightarrow \mathfrak{m}_{y'}/\mathfrak{m}_{y'}^2$, by construction we will have that $df|_{T_{y',S}} : T_{y',S} \rightarrow T_0\mathbb{A}^1$ is an isomorphism. Fiber $Z' = f^{-1}(0)$ contains y' and since f was homogeneous it is also endowed with a contracting $\mathbb{G}_m$ -action which lifts to $W' = \pi'^{-1}(Z')$. Putting $w' = y'$ and $\xi' = \pi'|_{W'}$ we get a conical pointed resolution (ξ', Z', W', w') .

To construct an étale equivalence between (ξ', Z', W', w') and (ξ, Z, W, w) we proceed as in Lemma 5.5.1 using the function $f \circ p_2 : U \rightarrow \mathbb{A}^1$ on U . Namely take $W'' = f \circ p_2^{-1}(0)$ and $w'' = u$. Shrinking W'' to $W_{f \circ p_2}$ as in Lemma 5.5.1 and repeating the argument we get the desired equivalence. $\square$

Lemma 5.5.3. *Let $\pi : X \rightarrow Y$ be a resolution with conical slices and let $y \neq y_0$ be a point different from the central one. Then there exists a pointed resolution (ξ, Z, W, w) and an étale equivalence $(\pi, X, Y, y) \sim_{et} (\xi, Z, W, w) \times \mathbb{A}^1$.*

Proof. We consider the contracting $\mathbb{G}_m$ -action on Y . The point y is non-central, so the stabilizer of y is a finite group scheme over $\mathbb{k}$. Since the action is separable, it is étale. Hence Luna's slice theorem [Lu] can be applied: there is a subvariety W in Y containing y and an étale map $\mathbb{G}_m \times W \rightarrow Y$. Resolution $\pi : X \rightarrow Y$ is $\mathbb{G}_m$ -equivariant, so this decomposition also lifts to an étale map $\mathbb{G}_m \times Z \rightarrow X$, where $Z = \pi^{-1}(W)$. This gives an étale equivalence $(\pi, X, Y, y) \sim_{et} (\xi, Z, W, w) \times \underline{\mathbb{G}_m}_{\{1\}} \sim_{et} (\xi, Z, W, w) \times \mathbb{A}^1$. $\square$

Proposition 5.5.1 (Conical slice). *Let $\pi : X \rightarrow Y$ be a resolution with conical slices and $y \neq y_0$ a point different from the central one. Then there exists a resolution with conical slices $\xi : Z \rightarrow W$ with the central point w_0 and an étale equivalence $(\pi, X, Y, y) \sim_{et} (\xi, Z, W, w_0) \times \mathbb{A}^1$.*

Proof. By Lemma 5.5.3 (π, X, Y, y) is étale equivalent to $(\xi, Z, W, w_0) \times \mathbb{A}^1$ for some (ξ, Z, W, w_0) and by Lemma 5.5.2 it can be chosen to be conical. We now need to prove that any point $w \neq w_0$ has a conical étale neighborhood. Let $t \in \mathbb{G}_m(\mathbb{k})$, then w has a conical étale neighbourhood if and only if tw is. Taking a suitable t we can assume that w is in the image of the correspondence between (π, X, Y, y) and (ξ, Z, W, w_0) , and that this correspondence restricts to an étale equivalence between $(\xi, Z, W, w) \times \mathbb{A}^1$ and $(\pi, X, Y, \tilde{y})$ for some point $\tilde{y} \in Y(\mathbb{k})$. Now, again by Lemma 5.5.3 and Lemma 5.5.2, $(\pi, X, Y, \tilde{y})$ is étale equivalent to $(\xi', Z', W', w'_0) \times \mathbb{A}^1$ for some conical resolution (ξ', Z', W', w'_0) with the central point w'_0 . We get that

$$(\xi, Z, W, w) \times \mathbb{A}^1 \sim_{et} (\xi', Z', W', w'_0) \times \mathbb{A}^1$$

and from Lemma 5.5.1 it follows that $(\xi, Z, W, w) \sim_{et} (\xi', Z', W', w'_0)$ and so any point $\xi : Z \rightarrow W$ admits a conical étale neighbourhood. The equalities $\xi_* \mathcal{O}_Z = \mathcal{O}_W$ and $R^1 \xi_* \mathcal{O}_Z = R^2 \xi_* \mathcal{O}_Z = 0$ follow from Lemma 5.4.1 and the flat base change (e.g. see Proposition 5.6.2 and apply it to $S = \mathbb{A}^1$) for the trivial family of resolutions $\pi \times \text{id} : X \times \mathbb{A}^1 \rightarrow Y \times \mathbb{A}^1$ over $\mathbb{A}^1$. $\square$

We end this subsection with the following important remark.

Remark 5.5.1. Let $\pi : X \rightarrow Y$ be a resolution with conical slices and let $y_0 \in Y(\mathbb{k})$ be the central point. Then by Proposition 5.5.1 we have an étale correspondence between Y and $\bigsqcup_y W_y \times \mathbb{A}^1$, where we take the union over all points $y \in Y(\mathbb{k}) \setminus y_0$ and W_y is the transversal slice to the $\mathbb{G}_m$ -orbit of y , constructed in Proposition 5.5.1. Note that it covers $Y \setminus y_0$. Moreover it extends to an étale correspondence between resolution $\pi : X \setminus \pi^{-1}(y_0) \rightarrow Y \setminus y_0$ and $\bigsqcup_y \xi_y \times \text{id} : Z_y \times \mathbb{A}^1 \rightarrow W_y \times \mathbb{A}^1$. Note that since Y is quasi-compact, the union $\bigsqcup_y \xi_y \times \text{id} : Z_y \times \mathbb{A}^1 \rightarrow W_y \times \mathbb{A}^1$ can be replaced by a finite subunion.

5.6. Reductions to characteristic p . Let R be a domain finitely generated over $\mathbb{Z}$ of characteristic 0 (meaning $p \cdot R \neq 0$ for all primes p). Let $p : \mathcal{X} \rightarrow S$ be a scheme of finite type over $S = \text{Spec } R$. Let $K = \text{Frac } R$ be the field of fractions and let $\mathbb{K} = \overline{K}$ be its algebraic closure. We have the geometric generic point map $\eta : \text{Spec } \mathbb{K} \rightarrow S$ given by the embedding $R \subset \mathbb{K}$. We have structure maps $p_S : S \rightarrow \text{Spec } \mathbb{Z}$, $p_{\mathcal{X}} : \mathcal{X} \rightarrow \text{Spec } \mathbb{Z}$. Let $\mathbb{k}_s$ be an algebraically closed field and $s : \text{Spec } \mathbb{k}_s \rightarrow S$ be a geometric point with the function field $\mathbb{k}_s$. Let X_s be the corresponding fiber: $X_s = \mathcal{X} \times_S \text{Spec } \mathbb{k}_s$. Taking $s = \eta$ we obtain the geometric generic fiber X_η , whose image is dense in X .

Consider now a proper map $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ such that $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a resolution over $\text{Spec } \mathbb{K}$ and $\mathcal{Y}$ is affine over S . Then for every $s : \text{Spec } \mathbb{k}_s \rightarrow S$, such that X_s is smooth and π_s is birational we have a resolution $\pi_s : X_s \rightarrow Y_s$ over $\text{Spec } \mathbb{k}_s$.

Proposition 5.6.1 ([EGA] IV₄, 17.7.8(ii)). *Let $\mathcal{X}$, X_η , X_s be as before and assume that X_η is smooth. Then there is a Zariski open $S' \subset S$ such that for any geometric point $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ the corresponding fiber X_s is smooth.*

Let $Y_\eta^\circ \subset Y_\eta$ be the dense open subset on which π_η is 1-to-1. The map $\pi_\eta^{-1} : Y_\eta^\circ \rightarrow X_\eta$ is defined over some finite extension $K \subset L$ and can be extended to some étale open $S' \rightarrow S$. Since the image of Y_η° is dense in $\mathcal{Y} \times_S S'$, shrinking S' we can also assume that the image of Y_s° in X_s for any $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ is dense and open as well. This proves that there exists an étale open $S' \rightarrow S$ such that for any geometric point $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ the map $\pi_s : X_s \rightarrow Y_s$ is a resolution with singularities.

Let's now assume that for $\pi_\eta : X_\eta \rightarrow Y_\eta$ we have $\mathcal{O}_{Y_\eta} \xrightarrow{\sim} (\pi_\eta)_* \mathcal{O}_{X_\eta}$ and $R^1(\pi_\eta)_* \mathcal{O}_{X_\eta} = R^2(\pi_\eta)_* \mathcal{O}_{X_\eta} = 0$. Replacing S with some Zariski open $S' \subset S$, we can assume that X and Y are both flat over S . In this case we have the base change theorem:

Proposition 5.6.2 ([EGA] III, 6.9.10). *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a morphism of schemes over S and $\mathcal{F}$ be a quasicoherent sheaf of $\mathcal{O}_{\mathcal{X}}$ -modules, flat over S . Assume that $\mathcal{O}_{\mathcal{Y}}$ -modules $R^i \pi_* \mathcal{F}$ are also flat over S . Let $s : \text{Spec } \mathbb{k}_s \rightarrow S$ be a geometric point and $s_Y : Y_s \rightarrow \mathcal{Y}$, $s_X : X_s \rightarrow \mathcal{X}$ be the corresponding morphisms from fibers. Let $\pi_s : X_s \rightarrow Y_s$ be the pull-back of π . Then*

$$R^i(\pi_s)_* (s_X^* \mathcal{F}) \simeq s_Y^* (R^i \pi_* \mathcal{F})$$

Remark 5.6.1. Applying the proposition to our situation and $\mathcal{F} = \mathcal{O}_{\mathcal{X}}$, we get that $(R^i \pi_s)_* \mathcal{O}_{X_s} \simeq s_Y^* (R^i \pi_* \mathcal{O}_{\mathcal{X}})$, in particular $(\pi_s)_* \mathcal{O}_{X_s} \simeq s_Y^* \mathcal{O}_{\mathcal{X}}$. We have a natural morphism $\mathcal{O}_{\mathcal{Y}} \rightarrow \pi_* \mathcal{O}_{\mathcal{X}}$ which is an isomorphism on a dense open subscheme of $\mathcal{Y}$ (containing the image of Y_η). The set of points where this morphism is not an isomorphism is closed, so is contained in some proper closed subscheme of $\mathcal{Y}$. Its image under projection $p_{Y/S} : Y \rightarrow S$ is constructible and since it does not contain the generic point, it is contained in some closed subscheme of S properly contained in S . Let $S' \subset S$ be the complement, then for any $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ we have $\mathcal{O}_{Y_s} \xrightarrow{\sim} (\pi_s)_* \mathcal{O}_{X_s}$. Similarly, shrinking S' more, we can also achieve $R^1(\pi_s)_* \mathcal{O}_{X_s} = R^2(\pi_s)_* \mathcal{O}_{X_s} = 0$ for all $s : \text{Spec } \mathbb{k}_s \rightarrow S$.

Let's now deal with the $\mathbb{G}_m$ -actions, namely with the separability (see Definition 5.3.2). Analogously to Definition 5.3.1 we define the subscheme $\mathcal{X}^{d\mathbb{G}_m} \subset \mathcal{X}$ of pseudo-fixed points. Namely let $\mathbb{G}_m = \text{Spec } R[t, t^{-1}]$ be the multiplicative group scheme over S , let $e : S \rightarrow \mathbb{G}_m$ be the embedding of the identity and $\mathcal{T}_e \mathbb{G}_m = e^* \mathcal{T}_{\mathbb{G}_m}$ be the tangent space at e . We have a nowhere-zero section $\partial_t \in \Gamma(S, \mathcal{T}_e \mathbb{G}_m)$ which gives the trivialisation $\mathcal{T}_e^* \mathbb{G}_m \xrightarrow{\sim} \mathcal{O}_S$. Consider $p^* \mathbb{G}_m$ as a group scheme over $\mathcal{X}$ (namely take $\mathcal{X} \times_S \mathbb{G}_m$) and let $\text{Lie}(p^* \mathbb{G}_m)$ be the corresponding Lie algebroid (considered just as a coherent sheaf). Since $p^* \mathbb{G}_m$ is just the fiber product, it is clear that $\text{Lie}(p^* \mathbb{G}_m) = p^*(\text{Lie}(\mathbb{G}_m)) = p^*(\mathcal{O}_S) = \mathcal{O}_{\mathcal{X}}$. The derivative of the $\mathbb{G}_m$ -action gives a map $a_S : \mathcal{O}_{\mathcal{X}} \xrightarrow{\sim} \text{Lie}(p^* \mathbb{G}_m) \rightarrow \text{Der}_R(\mathcal{O}_{\mathcal{X}})$ and the dual map $a_S^* : \Omega_{\mathcal{X}/S}^1 \rightarrow \mathcal{O}_{\mathcal{X}}$, where $\text{Der}(\mathcal{O}_{\mathcal{X}})$ and $\Omega_{\mathcal{X}/S}^1$ are the sheaves of R -linear derivations of $\mathcal{O}_{\mathcal{X}}$ and relative Kähler differentials on $\mathcal{X}$ respectively. The image of a_S^* is some sheaf of ideals on $\mathcal{X}$ which we denote by $\mathcal{I}_{\mathcal{X}, a_R}$. Similarly, for any geometric point s we have a map $a_s^* : \Omega_{\mathcal{X}/\mathbb{k}_s}^1 \rightarrow \mathcal{O}_{X_s}$ and a sheaf of ideals $\mathcal{I}_{X_s, a_s} \subset \mathcal{O}_{X_s}$. We define the subschemes of pseudo-fixed points $\mathcal{X}^{d\mathbb{G}_m} \subset \mathcal{X}$ and $X_s^{d\mathbb{G}_m} \subset X_s$ as subschemes determined by $\mathcal{I}_{\mathcal{X}, a_R}$ and $\mathcal{I}_{X_s, a_s}$ correspondingly.

Let $i_s : X_s \rightarrow \mathcal{X}$ be the natural morphism from the fiber, then the Cartesian square

$$\begin{array}{ccc} X_s & \xrightarrow{i_s} & \mathcal{X} \\ \downarrow & & \downarrow \\ \text{Spec } \mathbb{k}_s & \xrightarrow{s} & S \end{array}$$

induces an isomorphism $i_s^* \Omega_{\mathcal{X}/S}^1 \xrightarrow{\sim} \Omega_{\mathcal{X}/\mathbb{k}_s}^1$ which intertwines a_R^* and a_s^* . This gives the equality $i_s^* \mathcal{I}_{\mathcal{X}, a_R} = \mathcal{I}_{X_s, a_s}$ and the identification of the corresponding subschemes:

$$X_s^{d\mathbb{G}_m} = X_s \times_{\mathcal{X}} \mathcal{X}^{d\mathbb{G}_m}.$$

Recall that a $\mathbb{G}_m$ -action on a scheme X over an algebraically closed field $\mathbb{k}$ is called separable if $X^{d\mathbb{G}_m}(\mathbb{k}) = X^{\mathbb{G}_m}(\mathbb{k})$. Recall also that in the case $\text{char } \mathbb{k} = 0$ the action is always separable.

Lemma 5.6.1. *Let $\mathcal{X}$ be a scheme over $S = \operatorname{Spec} R$ endowed with a $\mathbb{G}_m$ -action. Let $S^{\text{non-sep}} \subset S$ be the subset, consisting of the images of geometric points $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S$ such that the induced action on X_s fails to be separable. Let $p_S : S \rightarrow \operatorname{Spec} \mathbb{Z}$ be the natural projection, then $p_S(S^{\text{non-sep}}) \subset \operatorname{Spec} \mathbb{Z}$ is a finite set.*

Proof. Let $\mathcal{Z} = \mathcal{X}^{d\mathbb{G}_m} \setminus \mathcal{X}^{\mathbb{G}_m}$ be the open subscheme of $\mathcal{X}^{\mathbb{G}_m}$. $S^{\text{non-sep}}$ is by definition the image of $\mathcal{Z}$ under $p : \mathcal{X} \rightarrow S$. Let $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S$ be a geometric point, where $\mathbb{k}_s$ is of characteristic 0. We know that in this case $\mathcal{X}^{d\mathbb{G}_m}(\mathbb{k}_s) = X_s^{d\mathbb{G}_m}(\mathbb{k}_s) = X_s^{\mathbb{G}_m}(\mathbb{k}_s) = \mathcal{X}^{\mathbb{G}_m}(\mathbb{k}_s)$. This means that $\mathcal{Z}(\mathbb{k}_s) = \emptyset$ for any such s and as a conclusion the image of $\mathcal{Z}$ under the natural projection $p_{\mathcal{Z}} : \mathcal{Z} \rightarrow \operatorname{Spec} \mathbb{Z}$ does not contain $\operatorname{Spec} \mathbb{Q}$. On the other hand the image of $\mathcal{Z}$ is constructible subset of $\operatorname{Spec} \mathbb{Z}$, so is a finite set. $\square$

In particular there exist an étale (as well as Zariski) open $S' \rightarrow S$ such that for any geometric point $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S'$ the action on X_s and Y_s is separable.

Proposition 5.6.3. *Let $S = \operatorname{Spec} R$, where R is a finitely generated $\mathbb{Z}$ -domain of characteristic 0. Let $\eta : \operatorname{Spec} \mathbb{K} \rightarrow S$ be the geometric generic point. Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a map of schemes over S , such that $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a resolution of singularities with conical slices. Then there exists an étale open $S' \rightarrow S$ such that for any geometric point $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S'$ the base change $\pi_s : X_s \rightarrow Y_s$ is a resolution with conical slices.*

Proof. The $\mathbb{G}_m$ -action on π_η is defined over some finite extension $K \subset L$, and it can be then extended to an action on π over some étale open $S' \rightarrow S$ finite type over S . Moreover, we can assume that the action on $\mathcal{Y} \times_S S'$ is contracting. Let $y_0 : S' \rightarrow \mathcal{X}$ be the central point.

By Proposition 5.6.1 and the comment about birationality below it, we can find étale open $S' \rightarrow S$ such that $\pi_s : X_s \rightarrow Y_s$ is a resolution of singularities for any $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S'$. By Remark 5.6.1 we can shrink it further so that $\mathcal{O}_{Y_s} \xrightarrow{\sim} (\pi_s)_* \mathcal{O}_{X_s}$ and $R^1(\pi_s)_* \mathcal{O}_{X_s} = R^2(\pi_s)_* \mathcal{O}_{X_s} = 0$. Finally Lemma 5.6.1 states that the $\mathbb{G}_m$ -action on π_s can also be assumed to be separable. It only remains to find étale conic neighbourhoods.

For that we proceed by induction on the dimension of $\mathcal{X}$. By Remark 5.5.2 we can find a finite collection of resolutions with conical slices $\theta_{i,\eta} : Z_{i,\eta} \rightarrow W_{i,\eta}$ and an étale correspondence between $\pi_\eta : X_\eta \setminus \pi^{-1}((y_0) \rightarrow Y_\eta \setminus (y_0))$ and $\bigsqcup \theta_{i,\eta} \times \operatorname{id} : Z_{i,\eta} \times \mathbb{A}^1 \rightarrow W_{i,\eta} \times \mathbb{A}^1$. Moreover we can assume that all this correspondences are defined over some étale open $S' \rightarrow S$, are étale, and moreover that each $\theta_i : Z_i \rightarrow W_i$ is a conical resolution (over S'). By induction we can assume that each $\theta_{i,s} : Z_{i,s} \rightarrow W_{i,s}$ is a resolution with conical slices and in particular that $w_s \in W_{i,s}$ has an étale conical neighbourhood. Since $\mathcal{Y} \times_S S'$ is affine over S' , shrinking S' we can also assume that any $y_s : \mathbb{k}_s \rightarrow \mathcal{Y} \times_S S'$ can be factorised through some global point $y : S' \rightarrow \mathcal{Y}$. Let's choose $\mathcal{W}_i \times \mathbb{A}^1$ which covers y under the correspondence and a point $(w, t) : S' \rightarrow \mathcal{W}_i \times \mathbb{A}^1$ that is in correspondence with y . Taking its product with $\mathbb{A}^1$ and the base change to $s : \operatorname{Spec} \mathbb{k} \rightarrow S'$ we obtain an étale conical neighbourhood of $y_s \in Y_s$. $\square$

6. $\mathbb{G}_m$ -WEIGHTS OF 1-FORMS ON A CONICAL RESOLUTION

Let R be a finitely generated $\mathbb{Z}$ -domain of characteristic 0 and $\eta : \operatorname{Spec} K \rightarrow S$ be its geometric generic point. Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a proper map of finite type schemes over $S = \operatorname{Spec} R$, such that $\mathcal{O}_{\mathcal{Y}} \xrightarrow{\sim} \pi_* \mathcal{O}_{\mathcal{X}}$ and $R^1 \pi_* \mathcal{O}_{\mathcal{X}} = R^2 \pi_* \mathcal{O}_{\mathcal{X}} = 0$. Assume that $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a conical resolution. Assume moreover that the corresponding $\mathbb{G}_m$ -action on π_η extends to π and that $\mathcal{Y}$ is contracted to a single point $y_0 : \operatorname{Spec} R \rightarrow \mathcal{Y}$ under this action. Consequently $\mathcal{Y}$ is affine and is equal to $\operatorname{Spec} A$ for some finitely generated R -algebra A . A comes with a natural positive grading $A = \bigoplus_{i \geq 0} A^i$, such that $A^0 \simeq R$ (given by y_0) and each A^i is of finite rank over R .

Replacing S with some étale open, following the proof of Proposition 5.6.3 we can assume that for any geometric point $s : \text{Spec } \mathbb{k}_s \rightarrow S$ we have a conical resolution $\pi_s : X_s \rightarrow Y_s$. The corresponding $\mathbb{G}_m$ -action induces an action on the space $V = H^0(X_s, \Omega_{X_s}^1)$, which decomposes into a direct sum over weights $k \in \mathbb{Z}$: $H^0(X_s, \Omega_{X_s}^1) = \bigoplus_k H^0(X_s, \Omega_{X_s}^1)^k$, where $t \circ \omega = t^k \omega$ for $\omega \in H^0(X_s, \Omega_{X_s}^1)^k$. Let also $H^0(X_s, \Omega_{X_s,cl}^1)^k \subset H^0(X_s, \Omega_{X_s}^1)^k$ denote the subspace of closed forms of weight k .

Theorem 6.0.1. *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be as above. Then $H^0(X_\eta, \Omega_{X_\eta}^1)^k = 0$ for $k \leq 0$, or in other words all weights of the space of 1-forms on X_η are strictly positive:*

$$H^0(X_\eta, \Omega_{X_\eta}^1) = \bigoplus_{k>0} H^0(X_\eta, \Omega_{X_\eta}^1)^k.$$

Moreover, there exists an étale open $S' \rightarrow S$ such that for any $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ the same is true for X_s :

$$H^0(X_s, \Omega_{X_s}^1) = \bigoplus_{k>0} H^0(X_s, \Omega_{X_s}^1)^k.$$

Proof. Let $p_{\mathcal{X}/S} : \mathcal{X} \rightarrow S$ and $p_{\mathcal{Y}/S} : \mathcal{Y} \rightarrow S$ denote the structure morphisms. For every i we have a quasi-coherent sheaf $(p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S}^i$ on $S = \text{Spec } R$ and a well-defined map $d : (p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S}^i \rightarrow (p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S}^{i+1}$. Its kernel is a subsheaf $(p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S,cl}^i \subset (p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}}^i$ of closed relative 1-forms. Recall that we had quasi-coherent sheaves $\pi_* \Omega_{\mathcal{X}/S}^i$ with subsheaves $\pi_* \Omega_{\mathcal{X}/S,cl}^i \subset \pi_* \Omega_{\mathcal{X}/S}^i$ of R -modules on $\mathcal{Y}$. The functor $(p_{\mathcal{Y}/S})_*$ induces isomorphisms:

$$(p_{\mathcal{Y}/S})_*(\pi_* \Omega_{\mathcal{X}/S}^i) \simeq (p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S}^i \quad \text{and} \quad (p_{\mathcal{Y}/S})_*(\pi_* \Omega_{\mathcal{X}/S,cl}^i) \simeq (p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S,cl}^i.$$

We will denote $(p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S}^1$ by $\mathcal{V}$ and $(p_{\mathcal{X}/S})_* \Omega_{\mathcal{X}/S,cl}^1$ by $\mathcal{V}_{cl}$. The corresponding R -modules of global sections will be denoted by V and V_{cl} . V has a natural A -module structure given by the multiplication of differential 1-forms on functions. Shrinking S , we can assume that V and V_{cl} are flat over R , and that the base-change holds for any $s : \text{Spec } \mathbb{k}_s \rightarrow S$. Namely for any $s : \text{Spec } \mathbb{k}_s \rightarrow S$ we have isomorphisms of $\mathbb{k}_s$ -vector spaces (sheaves on $\text{Spec } \mathbb{k}_s$):

$$(p_{X_s})_* \Omega_{X_s}^1 \simeq s^* \mathcal{V} \quad \text{and} \quad (p_{X_s})_* \Omega_{X_s,cl}^1 \simeq s^* \mathcal{V}_{cl},$$

where $p_{X_s} : X_s \rightarrow \text{Spec } \mathbb{k}_s$ is the structure map.

Now consider the $\mathbb{G}_m$ -action on $\mathcal{X}$. The sheaf $\Omega_{\mathcal{X}/S}^1$ has a natural $\mathbb{G}_m$ -equivariant structure, which gives $\mathbb{G}_m$ -equivariant structures on $\mathcal{V}$ and $\mathcal{V}_{cl}$ (with respect to trivial $\mathbb{G}_m$ -action on S). In other words we have a grading $V = \bigoplus_{k \in \mathbb{Z}} V^k$ which descends to the grading on $V_{cl} = \bigoplus_{k \in \mathbb{Z}} V_{cl}^k$. The grading on V agrees with the grading on A and turns V into a graded A -module. Since π is proper, V is a finitely generated over A , so each V^k is of finite rank over R . V_{cl}^k is an R -submodule of V^k , so is also finitely generated. Moreover, since A is positively graded, we have that $V^k = 0$ for all but finitely many $k < 0$. Consider the corresponding coherent sheaves $\mathcal{V}^k$ and $\mathcal{V}_{cl}^k$ on $\text{Spec } R$. The function $d_k(s) = \dim_{\mathbb{k}_s} s^* \mathcal{V}^k$ on the set of geometric points $s : \text{Spec } \mathbb{k}_s \rightarrow S$ is semi-continuous. In particular, if d_k is equal to n on generic point, then we can shrink S , such that the function $s \mapsto d_k(s)$ is constant and equal to n . Note that due to the flat base change $V_s^k := s^* \mathcal{V}^k$ is exactly the space $H^0(X_s, \Omega_{X_s}^1)^k$. Analogously we define $d_{k,cl}(s) := \dim_{\mathbb{k}_s} s^* \mathcal{V}_{cl}^k$. Let $(V_s^k)_{cl} \subset V_s^k$ denote the corresponding subspace. Note that from the flat base change $(V_s^k)_{cl} = H^0(X_s, \Omega_{X_s,cl}^1)^k$.

Let's assume that for some $k < 0$ we have $V_\eta^k = H^0(X_\eta, \Omega_{X_\eta}^1)^k \neq 0$. Then we can also find some $s : \text{Spec } \mathbb{k}_s \rightarrow S$ of characteristic $p > 0$ such that $d_k(s) > 0$. Since $V^k = 0$ for all but finitely many $k < 0$, there exists $N \gg 0$ such that $V_s^k = 0$ for all $k < -N$ and all s .

Lemma 6.0.2. *Let $\pi : X \rightarrow Y$ be a conical resolution over an algebraically closed field $\mathbb{k}$ of characteristic p , such that $R^1\pi_*\mathcal{O}_X = R^2\pi_*\mathcal{O}_X = 0$. Then all weights of $H^0(X, \Omega_X^1)$ are nonnegative and all $\mathbb{G}_m$ -invariant forms are closed.*

Proof. The action of $\mathbb{G}_m$ on $\pi : X \rightarrow Y$ gives an action of $\mathbb{G}_m^{(1)}$ on $\pi : X^{(1)} \rightarrow Y^{(1)}$. This produces an action of $\mathbb{G}_m^{(1)}$ on $H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$. Note that $H^0(X^{(1)}, (\mathrm{Fr}_X)_*\Omega_X^1)$ is also endowed with a natural action of $\mathbb{G}_m$ (without a twist), coming from the action on X . These actions differ by $\mathrm{Fr}_{\mathbb{G}_m}$, which agrees with the $\mathbb{k}$ -structures on both spaces. Cartier operator $\mathbb{C}$ is $\mathbb{k}$ -linear and commutes with the action, so it maps a component of weight k to a component of weight $\frac{k}{p}$. In particular all components with weights not divisible by p are killed by $\mathbb{C}$.

Let $V = H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$ and let $V^k = H^0(X^{(1)}, \Omega_{X^{(1)}}^1)^k$. Let $\alpha \in V^k$ be some homogeneous 1-form of weight $k < 0$. By Lemma 3.2.1 the map $\pi_*\mathbb{C} : \pi_*\Omega_{X,cl}^1 \rightarrow \pi_*\Omega_{X^{(1)}}^1$ is surjective and so does the induced map $\mathbb{C} : (V^{kp})_{cl} \rightarrow (V^k)^{(1)}$ for any k . We can identify $(V^k)^{(1)}$ with V^k using the Frobenius twist $\bullet^{(1)}$. But $V^{p^n k} = 0$ for n big enough, so we get that $V^k = 0$ for all $k < 0$.

It remains to treat the case $k = 0$. By Lemma 3.2.2, the map $(V^0)_{cl} \rightarrow V^0$ induced by Cartier morphism is a surjection. We get $\dim V^0 \leq \dim (V^0)_{cl}$, but since $(V^0)_{cl} \subset V^0$ the equality $(V^0)_{cl} = V^0$ follows. $\square$

After reshrinking S , Proposition 6.0.2 is true for all reductions $\pi_s : X_s \rightarrow Y_s$. From this we get that $V_\eta^k = 0$ for all $k < 0$ and that $V_\eta^0 = (V_\eta^0)_{cl}$ as well.

The field $\mathbb{K}$ of functions on the geometric generic point is an algebraic closure of finitely generated field over $\mathbb{Q}$ and admits an embedding $\mathbb{K} \rightarrow \mathbb{C}$ to complex numbers. Let $\xi : \mathrm{Spec} \mathbb{C} \rightarrow \mathrm{Spec} \mathbb{K}$ denote the corresponding morphism and let $\pi_\xi : X_\xi \rightarrow Y_\xi$ be the corresponding base change of $\pi_\eta : X_\eta \rightarrow Y_\eta$. From the flat base change we have $d_i(\eta) = d^i(\xi)$ and to prove that $V_\eta^0 = 0$ it is enough to prove that $H^0(X_\xi, \Omega_{X_\xi}^1)^{\mathbb{G}_m} = 0$ (where $V^{\mathbb{G}_m} \subset V$ denotes the invariants of $\mathbb{G}_m$ -action).

Let $\pi^{an} : X^{an} \rightarrow Y^{an}$ be the analytification of π_ξ , where X^{an} and Y^{an} are analytifications of X_ξ and Y_ξ . By assumption we have $R^1(\pi_\xi)_*\mathcal{O}_{X_\xi} = 0$ and Serre's GAGA provides isomorphisms

$$(R^i(\pi_\xi)_*\mathcal{O}_{X_\xi})^{an} \simeq R^i\pi_*^{an}\mathcal{O}_{X^{an}}.$$

of coherent sheaves on complex analytic space Y^{an} . In particular we have $R^1\pi_*^{an}\mathcal{O}_{X^{an}} = 0$.

Lemma 6.0.3. *Let $\pi : X \rightarrow Y$ be a proper morphism of complex analytic spaces with $R^1\pi_*\mathcal{O}_X = 0$. Then $R^1\pi_*\mathbb{C}_X = 0$.*

Proof. $R^1\pi_*\mathbb{C}_X = R^1\pi_*\mathbb{Z}_X \otimes \mathbb{C}_X$ and applying $R^\bullet\pi_*$ to the exponential sheaf sequence $0 \rightarrow \mathbb{Z}_X \rightarrow \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^\times \rightarrow 0$ we get that $R^1\pi_*\mathbb{Z}_X = 0$ and consequently $R^1\pi_*\mathbb{C}_X = 0$. $\square$

In our case Y^{an} is contractible ($\mathbb{G}_m$ -action contracts it to the central point), so using Serre-Leray spectral sequence we also get $H^1(X^{an}, \mathbb{C}) = 0$.

Let α be a $\mathbb{G}_m$ -invariant 1-form on X_ξ . We know that $d^0(\xi) = d_{cl}^0(\xi)$, so α is ∂ and $\bar{\partial}$ -closed. Therefore, its complex conjugate $\bar{\alpha} \in \Omega_{X^{an}}^{0,1}$ is $\bar{\partial}$ -closed and defines a cocycle in the Dolbeault resolution for $\mathcal{O}_{X^{an}}$, which gives rise to a $\mathbb{C}^\times$ -invariant class $[\bar{\alpha}]$ in $H^1(X^{an}, \mathcal{O}_{X^{an}})$.

Let p_X^{an} be the projection $p_X^{an} : X^{an} \rightarrow \mathrm{pt}$. By definition $H^i(X^{an}, \mathcal{O}_{X^{an}}) = R^i(p_X^{an})_*\mathcal{O}_{X^{an}}$. Obviously we have $p_X^{an} = p_Y^{an} \circ \pi_\xi^{an}$ and using Serre-Leray spectral sequence we get that $H^1(X^{an}, \mathcal{O}_{X^{an}}) = H^1(Y^{an}, \pi_*^{an}\mathcal{O}_{X^{an}}) = H^1(Y^{an}, \mathcal{O}_{Y^{an}}) = 0$, since Y_ξ is affine and Y^{an} is Stein. So $H^1(X^{an}, \mathcal{O}_{X^{an}}) = 0$ and $[\bar{\alpha}] = 0$. This means that there exists some $\mathbb{C}^\times$ -invariant smooth function f on X^{an} , such that $\bar{\partial}(f) = \bar{\alpha}$.

Lemma 6.0.4. *Let X be a complex manifold and let $f \in C^\infty(X, \mathbb{C})$ be a smooth function on X , such that $\partial\bar{\partial}(f) = 0$ in $\Omega_X^{\bullet,\bullet}$. Then, if $H^1(X, \mathbb{C}) = 0$, f is a sum of a holomorphic and an anti-holomorphic functions.*

Proof. By the local dd^c -lemma, statement is true locally on X . This means that the summation map $\mathcal{O}_X^{hol} \oplus \mathcal{O}_X^{ahol} \rightarrow \text{Ker } \partial\bar{\partial} \subset C^\infty(X, \mathbb{C})$ (of sheaves of $\mathbb{C}$ -vector spaces) is surjective. The kernel of this map is equal to the intersection $\mathcal{O}_X^{hol} \cap \mathcal{O}_X^{ahol} \subset C^\infty(X, \mathbb{C})$ which is the constant sheaf $\underline{\mathbb{C}}_X$. We get the short exact sequence

$$0 \rightarrow \underline{\mathbb{C}}_X \rightarrow \mathcal{O}_X^{hol} \oplus \mathcal{O}_X^{ahol} \rightarrow \text{Ker } \partial\bar{\partial} \rightarrow 0$$

and from $H^1(X, \mathbb{C}) = 0$ we see that the map on the global sections $H^0(X, \mathcal{O}_X^{hol}) \oplus H^0(X, \mathcal{O}_X^{ahol}) \rightarrow \Gamma(\text{Ker}(\partial\bar{\partial}))$ is also surjective. $\square$

We can apply the lemma to f : indeed, $\partial\bar{\partial}(f) = \partial\bar{\alpha} = d\bar{\alpha} = 0$. This means that $\bar{\alpha}$ is $\bar{\partial}$ of some anti-holomorphic $\mathbb{C}^\times$ -invariant anti-holomorphic function on X . Since π is proper any anti-holomorphic function on X is a pull-back of a(n) anti-holomorphic function on Y . In particular all $\mathbb{C}^\times$ -invariant anti-holomorphic functions on X are constant since Y is contracted to a single point. We get that $\bar{\alpha} = 0$ and consequently $\alpha = 0$. $\square$

In particular we get the following interesting corollary:

Corollary 6.0.1. *Let $\pi : X \rightarrow Y$ be a conical resolution of singularities over a field of characteristic 0 with $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$. Then all $\mathbb{G}_m$ -weights of $H^0(X, \Omega_X^1)$ are strictly positive.*

We do not know counterexamples to the statement of this corollary in characteristic p . We would not need to worry about strict positivity of weights of 1-forms (see Sect. 6.0.2) if this were true. This motivates for us the following conjecture:

Conjecture 6.0.1. *Let $\pi : X \rightarrow Y$ be a conical resolution over a field of any characteristic with $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$. Then all $\mathbb{G}_m$ -weights of $H^0(X, \Omega_X^1)$ are strictly positive.*

Remark 6.0.2. Note that from Hodge-de Rham spectral sequence, assuming $H^1(X, \mathcal{O}_X) = 0$, the de Rham cohomology $H_{\text{dR}}^1(X, \mathcal{O}_X)$ are just equal to $H^0(X, \Omega_X^1)/d(H^0(X, \mathcal{O}_X))$. Taking $\mathbb{G}_m$ -invariants, using the fact that there are no $\mathbb{G}_m$ -invariant exact forms, we get that $H_{\text{dR}}^1(X, \mathcal{O}_X)^{\mathbb{G}_m} = H^0(X, \Omega_X^1)^{\mathbb{G}_m}$. So Conjecture 6.0.1 is equivalent to $H_{\text{dR}}^1(X, \mathcal{O}_X)^{\mathbb{G}_m} = 0$ for any conical resolution $\pi : X \rightarrow Y$ with $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$.

Remark 6.0.3. Assume that our resolution $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ is defined over $R = \mathbb{Z}[1/N]$. Recent results by Bhatt, Morrow and Sholze ([BMS]) imply that $\dim H_{\text{dR}}^1(X_s, \mathbb{k}_s) \geq \dim H_{\text{sing}}^1(X_{\mathbb{C}}, \mathbb{F}_p)$ where $p = \text{char } \mathbb{k}_s$ and $\mathcal{X}$ is proper over $\text{Spec } R$. Though in our case $\mathcal{X}$ is not proper, one could expect the inequality to remain true if we replace the whole de Rham cohomology space with its $\mathbb{G}_m$ -invariant part: $\dim H_{\text{dR}}^1(X_s, \mathbb{k}_s)^{\mathbb{G}_m} \geq \dim H_{\text{sing}}^1(X_{\mathbb{C}}, \mathbb{F}_p)$. We do not know if this is true indeed, but provided it is, one could try to construct a counterexample to Conjecture 6.0.1 using that. By the universal coefficients formula, one just needs to find a conical resolution with non-trivial p -torsion in the 1-st homology group $H_1(X_{\mathbb{C}}, \mathbb{Z})$ (and then also find a model for π with good reduction at prime p). However we do not know if such examples exist.

6.1. Totally positive forms. Let $\pi : X \rightarrow Y$ be a resolution with conical slices over an algebraically closed field $\mathbb{k}$ of characteristic p , let $y \in Y(\mathbb{k})$ be a point and $(\pi, X, Y, y) \sim_{et} (\pi', X', Y', y')$ be an étale conic neighbourhood. We would like to have a good notion of a positive weight for 1-forms on X , even though there is no $\mathbb{G}_m$ -action and grading on the space of 1-forms on X itself.

By definition the étale equivalence is given by a commutative diagram

$$\begin{array}{ccccc} X & \xleftarrow{q_1} & X'' & \xrightarrow{q_2} & X' \\ \pi \downarrow & & \pi'' \downarrow & & \pi' \downarrow \\ Y & \xleftarrow{p_1} & Y'' & \xrightarrow{p_2} & Y' \end{array}$$

where all horizontal maps are étale. We have isomorphisms $q_1^* \Omega_X^1 \simeq \Omega_{X''}^1 \simeq q_2^* \Omega_{X'}^1$ as q_1, q_2 are étale and by the flat base change we also obtain isomorphisms $p_1^* \pi_* \Omega_X^1 \simeq \pi''_* \Omega_{X''}^1 \simeq p_2^* \pi'_* \Omega_{X'}^1$. By Lemma 5.4.1 we also know that $\mathcal{O}_{Y''} \xrightarrow{\sim} \pi''_* \mathcal{O}_{X''}$ and $R^1 \pi''_* \mathcal{O}_{X''} = R^2 \pi''_* \mathcal{O}_{X''} = 0$ as well as $\mathcal{O}_{Y'} \xrightarrow{\sim} \pi'_* \mathcal{O}_{X'}$ and $R^1 \pi'_* \mathcal{O}_{X'} = R^2 \pi'_* \mathcal{O}_{X'} = 0$. In particular $\pi' : X' \rightarrow Y'$ satisfies the assumptions of Lemma 6.0.2 and so all 1-forms on X' have positive weight.

To simplify the notations we denote $H^0(X, \Omega_X^1)$, $H^0(X', \Omega_{X'}^1)$ and $H^0(X'', \Omega_{X''}^1)$ by M , M' and M'' . Let also A , A' and A'' denote the rings of global functions $H^0(X, \mathcal{O}_X)$, $H^0(X', \mathcal{O}_{X'})$ and $H^0(X'', \mathcal{O}_{X''})$. The construction that we are going to give is Zariski local, so we can assume that $Y = \text{Spec } A$ and $Y'' = \text{Spec } A''$ (Y' is automatically affine since it is conical).

The torus action on Y' induces a grading on A' . Since Y' is conical, the weights of the action are non-negative and $\mathfrak{m}_{y'}$ is identified with $(A')^{>0} = \bigoplus_{k>0} (A')^k$, while $A' = \bigoplus_{k \geq 0} (A')^k$, where $(A')^0 = \mathbb{k}$. The module M' also possesses a grading and by Theorem 6.0.1 $(M')^{<0} = \bigoplus_{k<0} (M')^k = 0$. The strictly positive part $(M')^{>0}$ is naturally a submodule, moreover $\mathfrak{m}_{y'} \cdot M \subset M^{>0}$, so the quotient $M'/M'^{>0}$ is a finite-dimensional vector space over $\mathbb{k}$ and is supported at the point y' as a module over A' .

The étale equivalence above provides isomorphisms $M \otimes_A A'' \cong M'' \cong M' \otimes_{A'} A''$. Without loss of generality we can assume that the preimage of y' under p_2 consists of the single point y'' (if not we can throw out some divisors from Y'' that pass through the other points in the preimage, but do not pass through y''), so that $\mathfrak{m}_{y'} \cdot A'' \cong \mathfrak{m}_{y''}$. Under these assumptions we define an A'' -module $(M'')^{>0} = (M')^{>0} \otimes_{A'} A''$. Note that $\mathfrak{m}_{y'} \cdot M'' \subset (M'')^{>0}$ as $\mathfrak{m}_{y''} \cong \mathfrak{m}_{y'} \cdot A''$ and $\mathfrak{m}_{y'} \cdot M' \subset (M')^{>0}$. So, again, $M''/M''^{>0}$ is a finite-dimensional vector space supported at y'' .

As M is torsion free, the map $M \rightarrow M'' \cong M \otimes_A A''$ is an embedding. We define $M_{\sim \pi'}^{>0}$ as the intersection $M \cap (M'')^{>0}$. This is a submodule, and as $\mathfrak{m}_y \subset \mathfrak{m}_{y''}$, we have $\mathfrak{m}_y \cdot M \subset M_{\sim \pi'}^{>0}$. $M_{\sim \pi'}^{>0}$ is a submodule of differential 1-forms α on X , such that $q_1^* \alpha = \sum f_i \cdot q_2^* \alpha'_i$ where α'_i are forms of positive weight on X' and f_i are some functions on X'' .

Submodule $M_{\sim \pi'}^{>0}$ a priori depends on the choice of a conical neighbourhood (and a choice of equivalence) and we do not have a good argument for why in fact it does not. But this is not a big deal and instead we consider all conical neighbourhoods at once, namely we define a submodule $M_{\text{at } y}^{>0} \subset M$ as the sum of all $M_{\sim \pi'}^{>0}$ inside M . In other words differential 1-form α lies in $M_{\text{at } y}^{>0}$ if and only if $\alpha = \alpha_1 + \alpha_2 + \dots + \alpha_n$, where $\alpha_i \in M_{\sim \pi'_i}^{>0}$ for some conical étale neighbourhood $(\pi, X, Y, y) \sim_{et} (\pi'_i, X'_i, Y'_i, y'_i)$ (possibly different for each i).

Definition 6.1.1. Let $\pi : X \rightarrow Y$ be a resolution with conical slices. Differential 1-form $\alpha \in H^0(X, \Omega_X^1)$ is called *strictly positive at a point* $y \in Y(\mathbb{k})$ if α lies in $M_{\text{at } y}^{>0}$.

Remark 6.1.1. If the point y admits a conical neighbourhood (π', X', Y', y') for which all differential 1-forms have strictly positive weight, then $M_{\text{at } y}^{>0} = M$. Indeed, in this case $(M')^{>0} = M'$,

so $(M'')^{>0} = M''$ and as a result $M_{\sim\pi'}^{>0} = M$. In particular, if y belongs to the locus $Y^\circ \subset Y$ on which π is 1-to-1, then $(\pi, X, Y, y) \sim_{et} (\text{id}, \mathbb{A}^n, \mathbb{A}^n, 0)$. For $\mathbb{A}^n$ with contracting $\mathbb{G}_m$ -action all 1-forms have positive weight, so $M_{at\ y}^{>0} = M$.

Definition 6.1.2. Let $\pi : X \rightarrow Y$ be a resolution with conical slices. Differential 1-form $\alpha \in H^0(X, \Omega_X^1)$ is called *totally positive* if $\alpha \in M_{at\ y}^{>0}$ for all points $y \in Y(\mathbb{k})$. We denote the subspace of all totally positive forms on X by $H^0(X, \Omega_X^1)^{\gg 0}$.

Remark 6.1.2. If Conjecture 6.0.1 is true then $H^0(X, \Omega_X^1)^{\gg 0} = H^0(X, \Omega_X^1)$ and all differential 1-forms are totally positive.

In other words $H^0(X, \Omega_X^1)^{\gg 0} = \cap_y M_{at\ y}^{>0}$. This way the subspace of totally positive 1-forms is naturally an A -submodule. Since A is Noetherian and $M = H^0(X, \Omega_X^1)$ is finitely generated, $H^0(X, \Omega_X^1)^{\gg 0}$ is finitely generated too. We denote the corresponding coherent sheaf on $Y = \text{Spec } A$ by $M^{\gg 0}$ or $(\pi_* \Omega_X^1)^{\gg 0}$. We have an embedding of sheaves $(\pi_* \Omega_X^1)^{\gg 0} \subset \pi_* \Omega_X^1$ and we call the quotient sheaf $\pi_* \Omega_X^1 / (\pi_* \Omega_X^1)^{\gg 0}$ by $\mathcal{M}_{\pi, \text{inv}}$ and its global sections by $M_{\pi, \text{inv}} = H^0(X, \Omega_X^1) / H^0(X, \Omega_X^1)^{\gg 0}$.

Lemma 6.1.1. Let $i_y : \text{Spec } \mathbb{k} \hookrightarrow Y$ be the morphism given by a point $y \in Y(\mathbb{k})$. Then there is a natural surjection $i_y^*(\mathcal{M}_{\pi, \text{inv}}) \rightarrow M/M_{at\ y}^{>0}$.

Proof. As we have seen $\mathfrak{m}_{y'} \cdot M/M_{at\ y'}^{>0} = 0$, so for each point $y' \neq y$ we have $\mathfrak{m}_y \cdot M_{at\ y'}^{>0} = M_{at\ y'}^{>0}$. We have a short exact sequence

$$0 \rightarrow M^{\gg 0} \rightarrow M \rightarrow M_{\pi, \text{inv}} \rightarrow 0.$$

Tensoring it with $A/\mathfrak{m}_y$ we get exact sequence

$$M^{\gg 0}/\mathfrak{m}_y M^{\gg 0} \rightarrow M/\mathfrak{m}_y M \rightarrow M_{\pi, \text{inv}}/\mathfrak{m}_y M_{\pi, \text{inv}} \rightarrow 0$$

The last term in the sequence is by definition $i_y^*(\mathcal{M}_{\pi, \text{inv}})$. We also have a short exact sequence

$$0 \rightarrow M_{at\ y}^{>0} \rightarrow M \rightarrow M/M_{at\ y}^{>0} \rightarrow 0,$$

which being tensored with $A/\mathfrak{m}_y$ gives

$$M_{at\ y}^{>0}/\mathfrak{m}_y M_{at\ y}^{>0} \rightarrow M/\mathfrak{m}_y M \rightarrow M/M_{at\ y}^{>0} \rightarrow 0.$$

By definition of $M^{\gg 0}$ we have an embedding $M^{\gg 0} \subset M_{at\ y}^{>0}$. Moreover

$$\mathfrak{m}_y M^{\gg 0} = \mathfrak{m}_y \left(\bigcap_{y'} M_{at\ y'}^{>0} \right) = \mathfrak{m}_y M_{at\ y}^{>0} \cap \left(\bigcap_{y' \neq y} M_{at\ y'}^{>0} \right) \subset \mathfrak{m}_y M_{at\ y}^{>0},$$

so we also have an embedding $M^{\gg 0}/\mathfrak{m}_y M^{\gg 0} \hookrightarrow M_{at\ y}^{>0}/\mathfrak{m}_y M_{at\ y}^{>0}$. Finally commutative diagram

$$\begin{array}{ccccccc} M^{\gg 0}/\mathfrak{m}_y M^{\gg 0} & \longrightarrow & M/\mathfrak{m}_y M & \longrightarrow & i_y^*(\mathcal{M}_{\pi, \text{inv}}) & \longrightarrow & 0 \\ \downarrow & & \parallel & & \downarrow s & & \\ M_{at\ y}^{>0}/\mathfrak{m}_y M_{at\ y}^{>0} & \longrightarrow & M/\mathfrak{m}_y M & \longrightarrow & M/M_{at\ y}^{>0} & \longrightarrow & 0 \end{array}$$

given by the embedding $M^{\gg 0} \hookrightarrow M_{at\ y}^{>0}$, shows that the map s should be a surjection. $\square$

In particular if $\mathcal{M}_{\pi, \text{inv}} = 0$, then $M_{at\ y}^{>0} = M$ for all points $y \in Y(\mathbb{k})$.

Lemma 6.1.2. Let $j : U \hookrightarrow Y$ be an embedding of an open set and assume that for each $y \in U(\mathbb{k})$ we have $M/M_{at\ y}^{>0} = 0$. Then $j^* \mathcal{M}_{\pi, \text{inv}} = 0$.

Proof. All definitions are Zariski local on Y , so without loss of generality we can assume $U = Y$. But then by definition $M^{\gg 0} = \cap_y M_{\text{at } y}^{>0} = M$, so $M_{\pi, \text{inv}} = 0$. $\square$

Following the discussion in Remark 6.1.1, Y° is an example of such U . It follows from the lemma that $\mathcal{M}_{\pi, \text{inv}}$ is always supported on the exceptional locus $Y_{\text{exc}} = Y \setminus Y^\circ$.

Remark 6.1.3. If Conjecture 6.0.1 is true, then for any resolution with conical slices $\mathcal{M}_{\pi, \text{inv}} = 0$. If not, the sheaf $\mathcal{M}_{\pi, \text{inv}}$ is some bizarre invariant of the resolution $\pi : X \rightarrow Y$.

While considering several resolutions at the same moment, we will freely switch between the notation M and $H^0(X, \Omega_X^1)$, hopefully avoiding any confusions. Also if the source of étale equivalence needs to be specified we use a notation $M_{\pi \sim \pi'}^{>0}$, instead of $M_{\sim \pi'}^{>0}$.

Proposition 6.1.1. *Let $(\pi_1, X_1, Y_1, y_1) \sim_{\text{et}} (\pi_2, X_2, Y_2, y_2)$ be an étale equivalence between resolutions with conical slices. Then the stalk $(\mathcal{M}_{\pi_1, \text{inv}})_{y_1}$ is zero if and only if the stalk $(\mathcal{M}_{\pi_2, \text{inv}})_{y_2}$ is zero.*

Proof. Let $\pi : X \rightarrow Y$ be a resolution with conical slices and let $U \rightarrow Y$ be a surjective étale morphism. Let $V = U \times_Y X$ and let $\xi : V \rightarrow U$ be the corresponding resolution. Despite the fact that U is not necessarily conical and so does not need to be a resolution with conical slices, every point $u \in U(\mathbb{k})$ has an étale conical neighbourhood (e.g. coming from an étale conical neighbourhood for $p(u) \in Y(\mathbb{k})$) and so the definition of $\mathcal{M}_{\xi, \text{inv}}$ still makes sense.

We have a commutative square

$$\begin{array}{ccc} V & \xrightarrow{q} & X \\ \xi \downarrow & & \downarrow \pi \\ U & \xrightarrow{p} & Y \end{array}$$

where both p and q are étale and surjective.

Lemma 6.1.3. $\mathcal{M}_{\xi, \text{inv}} = 0$ if and only if $\mathcal{M}_{\pi, \text{inv}} = 0$

Proof. $\Leftarrow$. We will prove that $H^0(V, \Omega_V^1) = M_{\text{at } u}^{>0}$ for all points $u \in U(\mathbb{k})$. Let $y = p(u)$. Let $(\pi, X, Y, y) \sim_{\text{et}} (\pi', X', Y', y')$ be some étale conical neighbourhood of y . Then it also gives rise to an étale neighbourhood $(\xi, V, U, u) \sim (\pi', X', Y', y')$ of u :

$$\begin{array}{ccccc} & & V'' & & \\ & \swarrow & \downarrow \xi'' & \searrow & \\ V & & & & X'' \\ \downarrow \xi & \searrow & & \swarrow & \downarrow \pi'' \\ & U'' & & X & \\ & \downarrow & & \downarrow \pi & \\ U & & & & Y'' \\ & \swarrow & \searrow & & \\ & U & & Y & \end{array}$$

Here $\xi'' : V'' \rightarrow U''$ is the pull-back of $\pi'' : X'' \rightarrow Y''$ under $p : U \rightarrow Y$. Let $U = \text{Spec } B$. Now as $H^0(X, \Omega_X^1) = M_{\text{at } y}^{>0}$ we know that any 1-form α on X is a sum of forms α_i which lie in $M_{\pi \sim \pi'_i}^{>0}$ for some étale neighbourhoods $(\pi, X, Y, y) \sim (\pi'_i, X'_i, Y'_i, y'_i)$. It is easy to see from

the definitions that if α_i lies in $M_{\pi \sim \pi'_i}^{>0}$, then $q^*(\alpha_i)$ lies in $M_{\xi \sim \pi'_i}^{>0}$ (taking the pull-back of the corresponding expression on X'' to V''). But $H^0(V, \Omega_V^1) = H^0(X, \Omega_X^1) \otimes_A B$ and any differential 1-form β on V is of the form $\sum_j f_j q^*(\tilde{\alpha}_j)$ for some 1-forms $\tilde{\alpha}_j$ on X . So β lies in $M_{\text{at } u}^{>0}$ and we are done.

$\Rightarrow$ We will prove that $H^0(X, \Omega_X^1) = M_{\text{at } y}^{>0}$ for all points $y \in Y(\mathbb{k})$. Let u be a point which maps to y and let $(\xi, V, U, u) \sim_{\text{et}} (\xi', V', U', u')$ be an étale conical neighbourhood of u . It also gives rise to a neighbourhood $(\pi, X, Y, y) \sim_{\text{et}} (\xi', X', Y', y')$.

$$\begin{array}{ccccccc} V' & \longleftarrow & V'' & \longrightarrow & V & \xrightarrow{q} & X \\ \xi' \downarrow & & \xi'' \downarrow & & \xi \downarrow & & \pi \downarrow \\ U' & \longleftarrow & U'' & \longrightarrow & U & \xrightarrow{p} & Y \end{array}$$

We have A -submodule $M_{\pi \sim \xi'}^{>0} \subset H^0(X, \Omega_X^1)$ and B -submodule $M_{\xi \sim \xi'}^{>0} \subset H^0(X, \Omega_X^1)$. However both of them are defined as intersections with the same B'' -submodule of $H^0(V'', \Omega_{V''}^1)$: $M_{\pi \sim \xi'}^{>0} = H^0(X, \Omega_X^1) \cap H^0(V'', \Omega_{V''}^1)^{>0}$ and $M_{\xi \sim \xi'}^{>0} = H^0(V, \Omega_V^1) \cap H^0(V'', \Omega_{V''}^1)^{>0}$. From faithfully flat descent we get that $M_{\xi \sim \xi'}^{>0} = M_{\pi \sim \xi'}^{>0} \otimes_A B$. As $M_{\text{at } u}^{>0} = \cup_{\xi \sim \xi'} M_{\xi \sim \xi'}^{>0}$, from this we obtain that $M_{\text{at } u}^{>0} \subset M_{\text{at } y}^{>0} \otimes_A B$. But $M_{\text{at } u}^{>0} = H^0(V, \Omega_V^1)$, so again by faithfully flat descent $M_{\text{at } y}^{>0} = H^0(X, \Omega_X^1)$. $\square$

The statement of the proposition now easily follows from the lemma. If one of the stalks, say $(\mathcal{M}_{\pi_1, \text{inv}})_{y_1}$, is zero it means that $\mathcal{M}_{\pi_1, \text{inv}}$ is zero on some Zariski neighbourhood U of y_1 . As definition of $\mathcal{M}_{\pi_1, \text{inv}}$ was Zariski local we can replace Y_1 with U . An étale correspondence that gives the equivalence $(\pi_1, X_1 \times_{Y_1} U, U, y_1) \sim_{\text{et}} (\pi_2, X_2, Y_2, y_2)$ is surjective on some Zariski neighbourhoods U_1, U_2 of points y_1 and y_2 . Now, applying the lemma twice (for the first and then for the second arrow in the étale correspondence), we get that $\mathcal{M}_{\pi_2, \text{inv}}$ is zero on U_2 , so the stalk $(\mathcal{M}_{\pi_2, \text{inv}})_{y_2}$ is zero. $\square$

Lemma 6.1.4. *Let $\pi_1 : X_1 \rightarrow Y_1$ and $\pi_2 : X_2 \rightarrow Y_2$ be two resolutions with conical slices such that $\mathcal{M}_{\pi_1, \text{inv}}$ and $\mathcal{M}_{\pi_2, \text{inv}}$ are equal to 0. Let $\pi_1 \times \pi_2 : X_1 \times X_2 \rightarrow Y_1 \times Y_2$ be their product. Then $\mathcal{M}_{\pi \times \pi', \text{inv}} = 0$.*

Proof. We have $\Omega_{X_1 \times X_2}^1 = \Omega_{X_1}^1 \boxtimes \mathcal{O}_{X_2} \oplus \mathcal{O}_{X_1} \boxtimes \Omega_{X_2}^1$. We will show that $\pi_*(\mathcal{O}_{X_1} \boxtimes \Omega_{X_2}^1)$ lies in $(\pi_* \Omega_{X_1 \times X_2}^1)^{\geq 0}$. Let α be a section of $\pi_*(\mathcal{O}_{X_1} \boxtimes \Omega_{X_2}^1)$, and $(y_1, y_2) \in (Y_1 \times Y_2)(\mathbb{k})$ be a point. We can assume that α is decomposable: $\alpha = f \boxtimes \omega$. Since $\mathcal{M}_{\pi_2, \text{inv}} = 0$, ω is equal to $\omega_1 + \dots + \omega_n$ where $\omega_i \in M_{\pi'_i}^{>0}$ for some étale conical neighbourhoods $(\pi'_i, X'_i, Y'_i, y'_i)$ of $y_2 \in Y_2(\mathbb{k})$. Now consider any étale neighbourhood $(\tilde{\pi}_1, \tilde{X}_1, \tilde{Y}_1, \tilde{y}_1)$ of the point $y_1 \in Y_1(\mathbb{k})$. Each differential form $f \boxtimes \omega_i$ lies in $M_{\pi'_i \times \tilde{\pi}_1}^{>0}$, so $f \boxtimes \omega$ is strictly positive at (y_1, y_2) . Since the point (y_1, y_2) was arbitrary, we get that $f \boxtimes \omega$ is totally positive, and $\pi_*(\mathcal{O}_{X_1} \boxtimes \Omega_{X_2}^1) \subset (\pi_* \Omega_{X_1 \times X_2}^1)^{\geq 0}$. Analogously $\pi_*(\Omega_{X_1}^1 \boxtimes \mathcal{O}_{X_2}) \subset (\pi_* \Omega_{X_1 \times X_2}^1)^{\geq 0}$. Thus $\mathcal{M}_{\pi \times \pi', \text{inv}} = 0$. $\square$

Setting π_2 equal to $\text{id}_{\mathbb{A}^1} : \mathbb{A}^1 \rightarrow \mathbb{A}^1$ we get that $\mathcal{M}_{\pi} = 0$ implies $\mathcal{M}_{\pi \times \text{id}_{\mathbb{A}^1}} = 0$. The converse is also true:

Lemma 6.1.5. *Let $\pi : X \rightarrow Y$ be a resolution with conical slices such that $\mathcal{M}_{\pi \times \text{id}_{\mathbb{A}^1}, \text{inv}} = 0$. Then $\mathcal{M}_{\pi, \text{inv}} = 0$.*

Proof. Denote the natural projection $X \times \mathbb{A}^1 \rightarrow X$ by p and the natural embedding $X = X \times \{0\} \hookrightarrow X \times \mathbb{A}^1$ by ι . Let $\alpha \in H^0(X, \Omega_X^1)$ be a 1-form on X . We have $p^*\alpha \in H^0(X \times \mathbb{A}^1, \Omega_{X \times \mathbb{A}^1}^1)$. Since $\mathcal{M}_{\pi \times \text{id}_{\mathbb{A}^1}, \text{inv}} = 0$, for any $y \in Y(\mathbb{k})$ there exist 1-forms $\alpha_1, \dots, \alpha_n$ on $Y \times \mathbb{A}^1$ such that $p^*\alpha = \alpha_1 + \dots + \alpha_n$, and $\alpha_i \in M_{\sim \pi'_i}^{>0}$ for some étale conical neighbourhood $(\pi'_i, X'_i, Y'_i, y'_i)$ of $y \times \{0\} \in (Y \times \mathbb{A}^1)(\mathbb{k})$. Following the proof of Lemma 5.5.2, for each i we can find a $\mathbb{G}_m$ -invariant subvariety $W'_i \hookrightarrow Y'_i$ such that the conical neighbourhood above restricts to a conical étale neighbourhood $(\pi, X, Y, y) \sim_{\text{et}} (\xi'_i, Z'_i, W'_i, y'_i)$ where $Z'_i = X'_i \times_{Y'_i} W'_i$. Weights are preserved under restriction, so $\iota^*\alpha_i$ lies in $M_{\sim \xi'_i}^{>0}$ for each i . On the other hand, $\alpha = \iota^*p^*\alpha = \iota^*\alpha_1 + \dots + \iota^*\alpha_n$, so α is strictly positive at y . It follows that $\mathcal{M}_{\pi, \text{inv}} = 0$. $\square$

Definition 6.1.3. A resolution of singularities with conical slices $\pi : X \rightarrow Y$ is called *strictly positive* if the sheaf $\mathcal{M}_{\pi, \text{inv}}$ is zero. In other words, the defining property of strict positivity is that all 1-forms on X should be totally positive.

Lemma 6.1.4 and Lemma 6.1.5 show that $\pi : X \rightarrow Y$ is strictly positive if and only if $\pi \times \text{id}_{\mathbb{A}^1} : X \times \mathbb{A}^1 \rightarrow Y \times \mathbb{A}^1$ is.

6.2. Application to resolutions with conical slices. Recall the statement of Proposition 5.6.3: for a map $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ of schemes over $S = \text{Spec } R$, such that the geometric generic fiber is a resolution of singularities with conical slices, we can find a localisation $S' = \text{Spec } R[f^{-1}]$ such that for any geometric point $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ the base change $\pi_s : X_s \rightarrow Y_s$ is a resolution with conical slices as well.

Proposition 6.2.1. *In the context of Proposition 5.6.3, we can find an étale open $S' \rightarrow S$ such that for any $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ we have $\mathcal{M}_{\pi_s, \text{inv}} = 0$. In other words, for any $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ all differential 1-forms on X_s are totally positive.*

Proof. We proceed by induction on $\dim X_\eta$ using Remark 5.5.1. Following the proof of Proposition 5.6.3, we can find a finite collection of resolutions with conical slices $\theta_i : \mathcal{Z}_i \rightarrow \mathcal{W}_i$, defined over S' , such that there exists an étale correspondence between $\pi : \mathcal{X} \setminus \pi^{-1}((y_0) \rightarrow \mathcal{Y} \setminus (y_0))$ and $\bigsqcup \theta_i \times \text{id} : \mathcal{Z}_i \times \mathbb{A}^1 \rightarrow \mathcal{W}_i \times \mathbb{A}^1$, defined over S' as well. So by the induction hypothesis and Lemma 6.1.4 we can find S' such that $\mathcal{M}_{\pi_s, \text{inv}}$ is supported at the central point y_0 . Finally, for the central point the statement follows from Theorem 6.0.1. $\square$

Proposition 6.2.2. *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a map of schemes over $S = \text{Spec } R$ of characteristic 0 such that the generic fiber $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a resolution with conical slices. Then there exists an étale open $S' \rightarrow S$ such that for any geometric point $s : \text{Spec } \mathbb{k}_s \rightarrow S'$ the corresponding fiber $\pi_s : X_s \rightarrow Y_s$ is a strictly positive resolution with conical slices.*

Proof. Follows directly from Proposition 5.6.3 and Proposition 6.2.1. $\square$

7. DESCENT FOR RESOLUTIONS WITH CONICAL SLICES

7.1. Sheaf $\mathcal{Q}_{\pi, N}$ and étale equivalences. Recall that from Definition 4.2.1 we have the sheaf $\mathcal{Q}_{\pi, N} = \pi_* \Omega_{X(N)}^1 / \mathcal{S}_{\pi, N}$, where $\mathcal{S}_{\pi, N}$ is a subsheaf of 1-forms that are obtained as the N -th power of the Cartier operator applied to a Kähler differential on Y .

Recall the notion of étale equivalence from Definition 5.2.4. It would be useful to know that to prove $\mathcal{Q}_{\pi, N} = 0$ it is enough to prove $\mathcal{Q}_{\pi', N} = 0$ for some (π', X', Y', y') étale equivalent to (π, X, Y, y) . Some form of this is indeed true:

Lemma 7.1.1. *Given an étale equivalence $(\pi, X, Y, y) \sim_{\text{et}} (\pi', X', Y', y')$, $(\mathcal{Q}_{\pi, N})_y = 0$ if and only if $(\mathcal{Q}_{\pi', N})_{y'} = 0$.*

Proof. Let the equivalence be given by a commutative diagram

$$\begin{array}{ccccc} X & \xleftarrow{q_1} & X'' & \xrightarrow{q_2} & X' \\ \pi \downarrow & & \pi'' \downarrow & & \pi' \downarrow \\ Y & \xleftarrow{p_1} & Y'' & \xrightarrow{p_2} & Y' \end{array}$$

By definition q_i 's and p_i 's are étale, in particular flat. From flat base change we have isomorphisms $p_1^* \pi_* \Omega_{X(N)}^1 \simeq \pi''_* \Omega_{X''(N)}^1 \simeq p_2^* \pi'^* \Omega_{X'(N)}^1$. Moreover by functoriality of Cartier operator we also get isomorphisms $p_1^* \mathcal{S}_{\pi,N} \simeq \mathcal{S}_{\pi'',N} \simeq p_2^* \mathcal{S}_{\pi',N}$ and consequently $p_1^* \mathcal{Q}_{\pi,N} \simeq \mathcal{Q}_{\pi'',N} \simeq p_2^* \mathcal{Q}_{\pi',N}$. Let $y'' \in Y''(\mathbb{k})$ be a point which maps to y and y' . By faithfully flat descent we get that $(p_1^* \mathcal{Q}_{\pi,N})_{y''} = 0$ if and only if $(\mathcal{Q}_{\pi,N})_y = 0$. The same is true if we replace y with y' . So $(\mathcal{Q}_{\pi',N})_{y'} = 0 \Leftrightarrow (\mathcal{Q}_{\pi'',N})_{y''} = 0 \Leftrightarrow (\mathcal{Q}_{\pi,N})_y = 0$. $\square$

Recall also that we have the subsheaf $(\pi_* \Omega_X^1)^{\gg 0} \subset \pi_* \Omega_X^1$ (see Definition 6.1.2) of totally positive forms. We have a natural map $(\pi_* \Omega_{X(N)}^1)^{\gg 0} \rightarrow \mathcal{Q}_{\pi,N}$ and we denote the image by $\mathcal{Q}_{\pi,N}^{\gg 0}$. After all, for our argument to work we need to make some positivity assumptions, namely we want $\mathcal{Q}_{\pi,N}$ to be equal to $\mathcal{Q}_{\pi,N}^{\gg 0}$. In other words, we want each class in $\mathcal{Q}_{\pi,N}$ to be represented by some totally positive form. In particular, if $\pi : X \rightarrow Y$ is strictly positive (see Definition 6.1.3), $(\pi_* \Omega_{X(N)}^1)^{\gg 0} = \pi_* \Omega_{X(N)}^1$ and so $\mathcal{Q}_{\pi,N}^{\gg 0} = \mathcal{Q}_{\pi,N}$ for all N .

7.2. Descent for a resolution with conical slices. Let $\pi : X \rightarrow Y$ be a conical resolution with a central point y_0 and $Y = \text{Spec } A$. We recall the discussion about weights and Cartier operator from Lemma 6.0.2. The action of $\mathbb{G}_m$ on $\pi : X \rightarrow Y$ gives an action of $\mathbb{G}_m^{(1)}$ on $\pi : X^{(1)} \rightarrow Y^{(1)}$. This produces an action of $\mathbb{G}_m^{(1)}$ on $H^0(X^{(1)}, \Omega_{X^{(1)}}^1)$, of which we think as a standard one. Note that $H^0(X^{(1)}, (\text{Fr}_X)_* \Omega_X^1)$ is also endowed with a natural action of $\mathbb{G}_m$ (without twist), coming from the action on X . These actions differ by $\text{Fr}_{\mathbb{G}_m}$, and this agrees with natural $\mathbb{k}$ -structures on both spaces. Cartier operator C is $\mathbb{k}$ -linear and so maps a component of weight k to a component of weight $\frac{k}{p}$. In particular all components with weights not divisible by p are killed by C .

The $A^{(N)}$ -submodule $\mathcal{S}_{\pi,N} \subset H^0(X^{(N)}, \Omega_{X^{(N)}}^1)$ defines a $\mathbb{G}_m^{(N)}$ -invariant subspace (as the subspace $H^0(Y, \Omega_Y^1) \subset H^0(X, \Omega_X^1)$ is $\mathbb{G}_m$ -invariant and C is $\mathbb{k}$ -linear), so $\mathcal{S}_{\pi,N}$ has a structure of $\mathbb{G}_m$ -equivariant sheaf on $Y^{(N)}$. This way $\mathcal{Q}_{\pi,N}$ also obtains a $\mathbb{G}_m^{(N)}$ -equivariant structure, in particular its support should be a $\mathbb{G}_m^{(N)}$ -invariant subvariety of $Y^{(N)}$. Note that the central point of the resolution $\pi : X^{(N)} \rightarrow Y^{(N)}$ is the point $\text{Fr}_Y^N(y_0)$.

We are ready to prove the key lemma:

Lemma 7.2.1. *labelkey lemma Let $\pi : X \rightarrow Y$ be a conical resolution. Then $\mathcal{Q}_{\pi,N}$ is finite-dimensional over $\mathbb{k}$ if and only if $\mathcal{Q}_{\pi,N}$ is supported at the central point. In this case there exists $s > 0$, such that the image in $\mathcal{Q}_{\pi,N+s}$ of all 1-forms with positive weight is 0.*

Proof. Since $\mathcal{Q}_{\pi,N}$ is finite-dimensional over $\mathbb{k}$, the corresponding sheaf $\mathcal{Q}_{\pi,N}$ is supported at a finite set of points, but as the support should be $\mathbb{G}_m^{(N)}$ -invariant, this can be only the point $\text{Fr}_Y^N(y_0)$. For the other implication: $\mathcal{Q}_{\pi,N}$ is coherent, so if it is supported at the single point $\text{Fr}_Y^N(y_0)$, its global sections should be finite-dimensional.

In this case we see that $\mathcal{S}_{\pi,N}$ has finite codimension in $H^0(X^{(N)}, \Omega_{X^{(N)}}^1)$. In particular there is an integer r such that all forms of degree bigger than r lie in $\mathcal{S}_{\pi,N}$.

Let's take $\mathbb{G}_m^{(N)}$ -equivariant lift $Q_{\pi,N} \dashrightarrow H^0(X^{(N)}, \Omega_{X^{(N)}}^1)$ (as $\mathbb{k}$ -vector spaces). Let $\alpha_1, \dots, \alpha_n$ be a homogeneous basis of the image and let $d_i \geq 0$ be the weight of α_i . Without loss of generality we can assume that α is equal to some α_i with $d_i > 0$. Then by surjectivity of Cartier operator there exists a homogeneous closed form β_1 on $X^{(N-1)}$ of weight pd_i , such that $C(\beta_1) = \alpha_i$. Repeating this procedure s_i times, where $s_i > 0$ is any number with $p^{s_i}d_i > r$, we obtain a 1-form $\beta_{s_i} \in S_{\pi,N}$, such that $C^{s_i}(\beta_{s_i}) = \alpha_i$. It follows that $\alpha_i \in S_{\pi,N+s_i}$. Taking maximum of s_i over all i such that $d_i > 0$ and calling it s , we get that all 1-forms with positive weight lie in $S_{\pi,N+s}$ and so their image in $Q_{\pi,N+s}$ is 0. $\square$

Proposition 7.2.1. *Let $\pi : X \rightarrow Y$ be a strictly positive resolution with conical slices. Then $Q_{\pi,N} = 0$ for $N \gg 0$.*

Proof. We proceed by induction on $\dim X$. The base of induction is the case $\pi : \text{Spec } \mathbb{k} \rightarrow \text{Spec } \mathbb{k}$, where $Q_{\pi,N}^{\gg 0} = Q_{\pi,N} = 0$ is zero for all N . To prove that the sheaf $Q_{\pi,N}^{\gg 0}$ is zero in general we need to prove that its stalk at each point is. Let $y \neq y_0$ be a non-central point of Y . Then, by Lemma 5.5.1, (π, X, Y, y) is étale equivalent to the product $(\pi', X', Y', y'_0) \times \mathbb{A}^1$ for some resolution with conical slices $\pi' : X' \rightarrow Y'$ with the central point y'_0 . By Proposition 7.1.1 it is enough to prove that $(Q_{\pi' \times \text{id}, N})_{\text{Fr}_Y^N(y'_0) \times \{0\}} = 0$. Since π is strictly positive, by Lemma 6.1.4 and Proposition 6.1.1 we get that π' is strictly positive too. So, by induction, we can assume that there exists N , such that $Q_{\pi',N} = 0$. Now, by Lemma 4.2.2 we get that the stalk $(Q_{\pi' \times \text{id}, N})_{\text{Fr}_Y^N(y'_0) \times \{0\}}$ is zero. Moreover by Remark 5.5.1 we can choose such N uniformly. So there exists N such that $(Q_{\pi,N})_{\text{Fr}_Y^N(y)} = 0$ for all $y \neq y_0$. This means that the sheaf $Q_{\pi,N}$ is supported at $\text{Fr}_Y^N(y_0)$ and since it is also coherent we obtain that the space of global sections of $Q_{\pi,N}$ is finite-dimensional.

Let's take a $\mathbb{G}_m$ -equivariant lift $Q_{\pi,N} \dashrightarrow H^0(X^{(N)}, \Omega_{X^{(N)}}^1)$. Let $\alpha_1, \dots, \alpha_n$ be a homogeneous basis of the image of this lift. Since $\mathcal{M}_{\pi, \text{inv}} = 0$ we know that each α_i is totally positive. In particular it is strictly positive at y_0 , so $\alpha_i = \beta_{i1} + \dots + \beta_{in}$, where for each k there exists an étale conical neighbourhood $(\pi, X, Y, y_0) \sim_{\text{et}} (\pi'_{ik}, X'_{ik}, Y'_{ik}, y'_{ik})$, such that $\beta_{ik} \in M_{\pi'_{ik}}^{\gg 0}$ (see Definition 6.1.1). By definition it means that $q_1^*(\beta_{ik}) = \sum_j f_j \cdot q_2^*(\beta'_j)$ for some differential 1-forms $\beta'_j \in H^0(X'_{ik}, \Omega_{X'_{ik}}^1)$ with positive weight and some functions f_j on X''_{ik} . Using the isomorphisms $q_1^* Q_{\pi,N} \cong Q_{\pi'_{ik},N} \cong q_2^* Q_{\pi'_{ik},N}$ from the proof of Lemma 7.1.1 we see that it is enough to prove that the classes $[\beta'_j] \in Q_{\pi'_{ik},N}$ are zero. Moreover from the same isomorphisms we see that the stalk $(Q_{\pi'_{ik},N})_{\text{Fr}_Y^N(y'_{ik})}$ is finite-dimensional. Since the support should be $\mathbb{G}_m^{(N)}$ -invariant we see that it is actually supported at $\text{Fr}_Y^N(y'_{ik})$. So we are in the framework of Lemma ???. Fixing a decomposition as above for each α_i and a resolution for each β_{ik} , using Lemma ??? we see that there exists $s > 0$, such that the images in $Q_{\pi'_{ik},N+s}$ of all β'_j 's for all i and k are zero. So each α_i in $Q_{\pi,N+s}$ is 0 and as this was the basis of representatives for $Q_{\pi,N}$ and $Q_{\pi,N} \twoheadrightarrow Q_{\pi,N+s}$ we get that $Q_{\pi,N+s} = 0$. $\square$

Recall that if $Q_{\pi,N} = 0$, then $[\mathcal{D}_{X,\alpha}]$ descends to $Y^{(1)}$ for all differential 1-forms α (Definition 4.2.1). This way from Proposition 7.2.1 we obtain

Theorem 7.2.1. *Let $\pi : X \rightarrow Y$ be a strictly positive resolution with conical slices. Then the classes $[\mathcal{D}_{X,\alpha}]$ descend to Y for all $\alpha \in H^0(X, \Omega_X^1)$.*

Finally, by Proposition 6.2.2 we also get the following result, which is the main theorem in this paper:

Theorem 7.2.2. *Let $\pi : \mathcal{X} \rightarrow \mathcal{Y}$ be a map of schemes over $S = \operatorname{Spec} R$ of characteristic 0, such that the generic fiber $\pi_\eta : X_\eta \rightarrow Y_\eta$ is a resolution with conical slices. Then there exists an étale open $S' \rightarrow S$, such that for any geometric point $s : \operatorname{Spec} \mathbb{k}_s \rightarrow S'$*

- *the corresponding fiber $\pi_s : X_s \rightarrow Y_s$ is a strictly positive resolution with conical slices,*
- *the classes $[\mathcal{D}_{X_s,\alpha}]$ descend to Y_s for all $\alpha \in H^0(X_s, \Omega_{X_s}^1)$.*

8. EXAMPLES OF RESOLUTIONS WITH CONICAL SLICES

To construct examples we use Theorem 7.2.2: we prove that certain well-known resolutions in characteristic 0 are actually resolutions with conical slices. All of them are the so-called *symplectic resolutions* (see Sect. 8.1) and particular examples include Nakajima quiver varieties (Sect. 8.2.5), hypertoric varieties (Sect. 8.3.5) and Slodowy slices (Sect. 8.4). If we prove that all of them are resolutions with conical slices, Theorem 7.2.2 will say that for a general reduction s to characteristic p all classes $[\mathcal{D}_{X_s,\alpha}]$ in $\operatorname{Br}(X_s)$ descend to Y_s .

8.1. Background on symplectic resolutions. Let $\mathbb{K}$ be a field of characteristic 0.

Definition 8.1.1. A *symplectic resolution* is a smooth algebraic variety X over $\mathbb{K}$ equipped with a non-degenerate differential 2-form ω such that the affinisation map $\pi : X \rightarrow Y = \operatorname{Spec} H^0(X, \mathcal{O}_X)$ is a birational projective map.

Note that since $Y = \operatorname{Spec} H^0(X, \mathcal{O}_X)$, we have that $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and so Y is normal. Conversely, if Y is normal and affine, then $\mathcal{O}_Y \xrightarrow{\sim} \pi_* \mathcal{O}_X$ and so $Y = \operatorname{Spec} H^0(X, \mathcal{O}_X)$.

Since Ω is non-degenerate it is easy to see that the dimension of X is even and that $\omega^{\dim X/2}$ gives a trivialisation of the canonical bundle $K_X := \Omega_X^{\dim X}$. Grauert-Riemenschneider theorem (Theorem 4.3.9 in [La]) states that $R^\bullet \pi_*(K_X) = \pi_*(K_X)$ for any proper birational map $\pi : X \rightarrow Y$. Since in this case $K_X = \mathcal{O}_X$, we get that every symplectic resolution is necessarily rational, namely $R^\bullet \pi_*(\mathcal{O}_X) = \mathcal{O}_Y$.

In particular $R^1 \pi_* \mathcal{O}_X = R^2 \pi_* \mathcal{O}_X = 0$, so to prove that $\pi : X \rightarrow Y$ is a resolution with conical slices it is enough to construct: 1) a contracting $\mathbb{G}_m$ -action on π , 2) an étale conical neighbourhood for any point $y \in Y(\mathbb{K})$.

8.2. Marsden-Weinstein reductions for representations of quivers. In this subsection we mostly follow Crawley-Boevey's paper [CrB1] on normality of Marsden-Weinstein reductions for representations of quivers. The main results we use are Theorem 4.9 and Corollary 4.10 from [CrB1] but with a slight elaboration, see Proposition 8.2.3. We give a sketch of the content needed from [CrB1], so that it becomes clear that Proposition 8.2.3 follows easily from the mentioned results.

Let $\mathbb{K}$ be an algebraically closed field of characteristic 0 and let A be a finite-dimensional semisimple associative K -algebra. Since $\mathbb{K}$ is algebraically closed we have that A is just a direct sum of matrix algebras. Any left or right A -module and any A - A -bimodule is semisimple.

Let M be a finite-dimensional A - A -bimodule. A symplectic bilinear form ω on M is called *balanced* if $\omega(xa, y) = -\omega(y, ax) = -\omega(ya, x) = \omega(x, ya)$ for all $a \in A$ and $x, y \in M$. Subspace $U \subset M$ is called *Lagrangian* if $U = U^\perp$ and (co)isotropic if $U \subset U^\perp$ ($U^\perp \subset U$).

Not very difficult to prove is the following analogue of Darboux Theorem for bimodules with a balanced symplectic form:

Theorem 8.2.1 ([CrB1], Theorem 2.2). *If M is an A - A -bimodule with a balanced symplectic form ω , then any maximal isotropic sub-bimodule S of M has an isotropic bimodule complement. Both S and the complement are Lagrangian. Moreover there is an isomorphism of bimodules $M \cong S \oplus S^*$, under which $\omega((s, f), (s', f')) = f(s') - f'(s)$.*

Corollary 8.2.1 ([CrB1], Corollary 2.3). *If M is an A - A -bimodule with a balanced symplectic form ω , then any isotropic sub-bimodule S of M has a coisotropic bimodule complement.*

Definition 8.2.1. By a quadruple (A, M, ω, ζ) we will mean a data of a finite-dimensional semisimple associative $\mathbb{K}$ -algebra A , a finite-dimensional A - A -bimodule M , a balanced symplectic form ω on M , and a linear map $\zeta : A \rightarrow \mathbb{K}$ satisfying $\zeta(ab) = \zeta(ba)$ for all $a, b \in A$ (in other words $\zeta \in (A/[A, A])^* \subset A^*$).

Let $G = A^\times$ be the group of units of A . Since A is a direct sum of matrix algebras G is a product of copies of general linear groups. G acts by conjugation on any A - A -bimodule M via $g \circ m = gmg^{-1}$ (where the multiplication on the right hand side of the equation is given by the action of A from the left and the right on M). Note that $\text{Lie}(G) = A$, where the bracket on A is given by commutator.

Let $\mu : M \rightarrow A^*$ be the map defined by $\langle \mu(m), a \rangle = \omega(m, am)$. Note that $\omega(m, am) = \omega(ma, m) = -\omega(m, ma)$, so $\langle \mu(m), a \rangle = \frac{1}{2}\omega(m, [a, m])$, where $[a, m] = am - ma$.

Lemma 8.2.1 ([CrB1], Lemma 3.1). *μ is a moment map for the action of G on M .*

Proof. By the definition of a moment map we need to prove that $\langle d\mu_m(v), a \rangle = \omega_m(v, a_m)$ for all $m \in M$, $v \in T_m M$ and $a \in \text{Lie}(G)$. Here ω_m is the symplectic form on $T_m M$ induced by ω and $m \mapsto a_m$ is the vector field on M induced by a .

After the natural identification of $T_m M$ and $\text{Lie}(G)$ with A , the vector field a_m induced by a is given by the map $M \rightarrow M$, sending m to $[a, m] = am - ma$.

On the other hand from the formula for μ we see that $\langle d\mu_m(v), a \rangle = \omega(v, am) + \omega(m, av) = \omega(v, am) + \omega(ma, v) = \omega(v, [a, m]) = \omega(v, a_m)$. So μ is a moment map. $\square$

Definition 8.2.2. A categorical quotient

$$N(A, M, \omega, \zeta) = \mu^{-1}(\zeta) // G$$

is by definition the *Marsden-Weinstein reduction* associated to the quadruple (A, M, ω, ζ) .

Let Q be a finite quiver with a vertex set I , and let $h(a)$ and $t(a)$ denote the head and tail vertices of an arrow a in Q . For $\alpha \in \mathbb{N}^I$, the space of representations of Q with dimension vector α is by definition a $\mathbb{K}$ -vector space

$$\text{Rep}(\overline{Q}, \alpha) = \bigoplus_{a \in \overline{Q}} \text{Mat}(\alpha_{h(a)} \times \alpha_{t(a)}, \mathbb{K}).$$

Let $\overline{Q}$ be the double of Q , obtained from Q by adjoining a reverse arrow a^* for each arrow $a \in Q$. Finite-dimensional $\mathbb{K}$ -algebra $\text{End}(\alpha) = \prod_{i \in I} \text{Mat}(\alpha_i, \mathbb{K})$ is semisimple and acts naturally on $\text{Rep}(\overline{Q}, \alpha)$ both from the left and the right, turning it into $\text{End}(\alpha)$ - $\text{End}(\alpha)$ -bimodule. The space $\text{Rep}(\overline{Q}, \alpha)$ can be identified with the total space of cotangent bundle to $\text{Rep}(Q, \alpha)$. From this identification $\text{Rep}(\overline{Q}, \alpha)$ obtains the canonical symplectic form ω_α , which in explicit form is given by

$$\omega_\alpha(x, y) = \sum_{a \in Q} (\text{tr}(x_{a^*} y_a) - \text{tr}(x_a y_{a^*})).$$

It is easy to see that ω_α is balanced with respect to the action of $\text{End}(\alpha)$. The group $\text{GL}(\alpha) = \text{End}(\alpha)^\times = \prod_{i \in I} \text{GL}(\alpha_i)$ acts on $\text{Rep}(\overline{Q}, \alpha)$ by symplectomorphisms there is a moment map μ_α

$$\mu_\alpha(x) = \sum_{a \in \overline{Q}} [x_a, x_{a^*}] \in \text{End}(\alpha),$$

where $\text{End}(\alpha)$ is identified with its dual using the trace. Finally to a vector $\lambda \in \mathbb{K}^I$ we can associate a function $\zeta_\lambda : \text{End}(\alpha) \rightarrow \mathbb{K}$, defined by $\zeta_\lambda(\theta) = \sum_{i \in I} \lambda_i \text{tr}(\theta_i)$.

This way a quiver Q with vertex set I , elements $\alpha \in \mathbb{N}^I$ and $\lambda \in \mathbb{K}^I$, gives rise to a quadruple $(\text{End}(\alpha), \text{Rep}(\overline{Q}, \alpha), \omega_\alpha, \zeta_\lambda)$. We have an affine variety $N(\text{End}(\alpha), \text{Rep}(\overline{Q}, \alpha), \omega_\alpha, \zeta_\lambda)$ (see Definition 8.2.2) which in this case will be denoted simply by $N_Q(\lambda, \alpha)$.

Lemma 8.2.2 ([CrB1], Lemma 3.3). *Every quadruple (A, M, ω, ζ) is of the form $(\text{End}(\alpha), \text{Rep}(\overline{Q}, \alpha), \omega_\alpha, \zeta_\lambda)$ for some Q , α and λ .*

Proof. Using the analogue of Darboux theorem Theorem 8.2.1 we can find a maximal isotropic submodule S inside M and identify M with $S \oplus S^*$. Now we will give a construction of the quiver Q corresponding to S .

The vertex set I of Q is chosen to index the simple left A -modules up to isomorphism. The dimension vector α is defined by letting α_i to be equal to the dimension of the corresponding simple module. As A is a product of matrix algebras, we have $\text{End}(\alpha) = A$.

The number of arrows between X and Y in Q is defined as the multiplicity of $X \otimes Y^*$ in S (multiplicity can be 0). Note that if X and Y are simple left A -modules, then $X \otimes Y^*$ is a simple A - A -bimodule and every simple A - A -bimodule can be obtained this way. It follows that $S = \text{Rep}(Q, \alpha)$ and $M = \text{Rep}(\overline{Q}, \alpha)$. It also not hard to check that the described moment maps coincide.

Finally, ζ is an Ad-invariant element of $A^* \cong \text{End}(\alpha)^*$ and these are generated by the trace functions on each of the simple components of A . The coordinates of ζ in this basis define a vector $\lambda \in \mathbb{K}^I$, such that $\zeta = \zeta_\lambda$. $\square$

8.2.1. Properties of the map μ_α . For any $\lambda \in \mathbb{K}^I$ we denote by $\lambda \cdot \alpha$ the sum $\sum_{i \in I} \alpha_i \lambda_i$. Note that the action of $\text{GL}(\alpha)$ on $\text{Rep}(\overline{Q}, \alpha)$ factorises through $\mathbb{P}G(\alpha) = (\prod_{i \in I} \text{GL}(\alpha_i)) / \mathbb{G}_m$, where the scalars $\mathbb{G}_m$ are embedded diagonally. One-dimensional subspace $\mathbb{K} = \text{Lie}(\mathbb{G}_m) \subset \mathbb{K}^I$, where $\mathbb{K}^I$ is identified with $(\text{End}(\alpha) / [\text{End}(\alpha), \text{End}(\alpha)])^*$ using trace, is spanned by the vector $(\alpha_i)_{i \in I} \in \mathbb{K}^I$ since the trace of a scalar matrix t of size α_i is equal to $\alpha_i \cdot t$. In particular the image of μ_α lies in $\text{Lie}(\mathbb{G}_m)^\perp \subset \text{End}(\alpha)$ or in other words if λ is in the image of μ_α , then $\lambda \cdot \alpha = 0$.

Associated to a quiver Q there is also a quadratic form

$$q_Q(\alpha) = \sum_{i \in I} \alpha_i^2 - \sum_{a \in Q} \alpha_{h(a)} \alpha_{t(a)}$$

on $\mathbb{Z}^I$. We denote by $(\alpha, \beta)_Q$ the symmetric bilinear form with $(\alpha, \alpha)_Q = 2q(\alpha)$ and we define $p_Q(\alpha)$ as $1 - q_Q(\alpha)$. Associated to Q there is a subset $\Delta^+ \subset \mathbb{Z}^I$ of positive roots (positive roots of the corresponding Kac-Moody algebra) which corresponds exactly to the dimension vectors of indecomposable representations of Q (this is Kac theorem, see [Kac]).

Theorem 8.2.2 ([CrB2], Theorem 1.1). *For $\alpha \in \mathbb{N}^I$ the following are equivalent*

- (1) μ_α is a flat morphism.
- (2) $\mu_\alpha^{-1}(0)$ has dimension $\alpha \cdot \alpha - 1 + 2p(\alpha)$.
- (3) $p(\alpha) \geq \sum_{t=1}^r p(\beta^{(t)})$ for any decomposition $\alpha = \beta^{(1)} + \dots + \beta^{(r)}$ into positive roots.
- (4) $p(\alpha) \geq \sum_{t=1}^r p(\beta^{(t)})$ for any decomposition $\alpha = \beta^{(1)} + \dots + \beta^{(r)}$ into nonzero $\beta^{(t)} \in \mathbb{N}^I$

The way to study the fibers $\mu_\alpha^{-1}(\lambda)$ and varieties $N_Q(\lambda, \alpha)$, introduced by Crawley-Boevey, was to relate them to the representations of the certain algebra Π^λ , called the *deformed pre-projective algebra*. It is defined as

$$\Pi^\lambda = \mathbb{K}\overline{Q} / \left(\sum_{a \in \overline{Q}} [a, a^*] - \sum_{i \in I} \lambda_i e_i \right),$$

where $\mathbb{K}\overline{Q}$ is the path algebra of $\overline{Q}$. For $\lambda \in \mathbb{K}^I$, such that $\lambda \cdot \alpha = 0$ the variety $\mu_\alpha^{-1}(\lambda)$ is identified with the space of representations of Π^λ with the dimension vector α . Closed orbits of G -action on $\text{Rep}(\overline{Q}, \alpha)$ correspond to semisimple representations of $\overline{Q}$. So points of $N_Q(\lambda, \alpha) = \mu_\alpha^{-1}(\lambda) // G$ correspond to isomorphism classes of semisimple representations of Π^λ of dimension α .

For $\lambda \in \mathbb{K}^I$ let Δ_λ^+ denote the set of positive roots α with $\lambda \cdot \alpha = 0$. Following Crawley-Boevey we define $\Sigma_\lambda \subset \Delta_\lambda^+$ to be the set of $\alpha \in \Delta_\lambda^+$ such that

$$p(\alpha) > \sum_{t=1}^r p(\beta^{(t)})$$

for any decomposition of $\alpha = \beta^{(1)} + \dots + \beta^{(r)}$ where $r \geq 2$ and $\beta^{(t)}$ lie in Δ_λ^+ .

Theorem 8.2.3 ([CrB2], Theorem 1.2). *For $\lambda \in \mathbb{K}^I$ and $\alpha \in \mathbb{N}^I$ the following are equivalent*

- (1) *There is a simple representation of Π^λ of dimension vector α*
- (2) *α is an element of Σ_λ*

In this case μ_α^{-1} is a reduced and irreducible complete intersection of dimension $\alpha \cdot \alpha - 1 + 2p(\alpha)$ and the general element of $\mu_\alpha^{-1}(\lambda)$ is a simple representation of Π^λ .

We also will need the following corollary:

Corollary 8.2.2 ([CrB2], Corollary 1.4). *If $\alpha \in \Sigma_\lambda$, then $N_Q(\lambda, \alpha) = \mu_\alpha^{-1}(\lambda) // G$ is a reduced and irreducible scheme of dimension $2p(\alpha)$.*

Remark 8.2.1. Note that if $\alpha \in \Sigma_0$, then α lies in Σ_λ for all $\lambda \in \mathbb{K}^I$ such that $\lambda \cdot \alpha = 0$ and also satisfies the conditions of Theorem 8.2.2.

8.2.2. GIT and resolutions of singularities of $N_Q(\lambda, \alpha)$. Let θ be a character of $\mathbb{P}G(\alpha) = \text{GL}(\alpha)/\mathbb{G}_m = (\prod_{i=1}^I \text{GL}(\alpha_i))/\mathbb{G}_m$. θ can be considered as an element of $\mathbb{Z}^I$, such that $\theta \cdot \alpha = 0$. Natural action of $\text{GL}(\alpha)$ on $\text{Rep}(\overline{Q}, \alpha)$ factorises through $\mathbb{P}G(\alpha)$, so we can consider open subsets of θ -semistable and θ -stable points of $\text{Rep}(\overline{Q}, \alpha)$. The corresponding GIT quotient $N_Q(\lambda, \alpha)^\theta = \mu_\alpha^{-1}(\lambda) //_\theta G$ is defined as the categorical quotient $\mu^{-1}(\lambda)^{\theta-ss} // G$, where $\mu^{-1}(\lambda)^{\theta-ss}$ is by definition the intersection of $\mu^{-1}(\lambda)$ with the set of θ -semistable points $\text{Rep}(\overline{Q}, \alpha)^{\theta-ss}$. This is a variety endowed with a natural projective map $\pi_\lambda^\theta : N_Q(\lambda, \alpha)^\theta \rightarrow N_Q(\lambda, \alpha)$ called semisimplification, which is better seen from the equivalent definition of $N_Q(\lambda, \alpha)^\theta$ as

$$N_Q(\lambda, \alpha)^\theta = \text{Proj} \bigoplus_{m=0}^{\infty} \{f \in \mathcal{O}(\mu^{-1}(0)) \mid g \circ f = \theta(g)^m \cdot f, \text{ for any } g \in G\}.$$

The map $\pi_\lambda^\theta : N_Q(\lambda, \alpha)^\theta \rightarrow N_Q(\lambda, \alpha)$ is then just the projection from Proj to the Spec of its 0th component.

The stable locus $\mu^{-1}(\lambda)^{\theta-st} \subset \mu^{-1}(\lambda)^{\theta-ss}$ is defined as the subset of points with a finite stabiliser. Geometric invariant theory says that this is an open subset and if $\mu^{-1}(\lambda)^{\theta-st}$ is smooth, the quotient $\mu^{-1}(\lambda)^{\theta-st} // G$ is smooth too.

In the classical paper by King [Ki] it is proven that for the space of representations of a quiver θ -(semi)stability in the sense of GIT is the same as θ -(semi)stability in terms of slopes. In particular a simple representation is θ -stable for any character θ . It also follows that if α is indivisible, meaning that α_i do not have common divisors, one can find a character θ such that the set of θ -semistable points coincides with the set of θ -stable ones ([Ki], Remark 5.4).

If $\alpha \in \Sigma_\lambda$ then by Theorem 8.2.3 $\mu_\alpha^{-1}(\lambda)$ is irreducible and the general element is a simple representation of Π^λ , so is θ -stable. So the set of θ -semistable points is dense and it follows

that the semisimplification map

$$\pi_\lambda^\theta : N_Q^\theta(\lambda, \alpha) \rightarrow N_Q(\lambda, \alpha)$$

is a birational map of irreducible varieties. θ -stable Π^λ -modules have trivial endomorphisms, so by ([CrB3], Lemma 10.3) stable locus is smooth. If in addition α is indivisible, taking θ such that all θ -semistable points are θ -stable we get that $N_Q^\theta(\lambda, \alpha)$ is smooth and that π_λ^θ is a resolution of singularities. Moreover $\mu_\alpha^{-1}(\lambda) //_\theta \mathbb{P}G(\alpha)$ is a symplectic reduction, so is endowed with a natural symplectic form. Finally, in ([CrB1], Theorem 8.2) it is proven that $N(\lambda, \alpha)$ is normal for all λ and α . This turns π_λ^θ into a symplectic resolution.

We can summarise this discussion in the following:

Proposition 8.2.1 ([CrB1],[CrB2],[CrB3]). *Let $\alpha \in \mathbb{N}^I$ and $\lambda \in \mathbb{K}^I$ be such that*

- $\lambda \cdot \alpha = 0$,
- $\alpha \in \Sigma_\lambda$,
- α is indivisible.

Then the map $\pi_\lambda^\theta : N_Q^\theta(\lambda, \alpha) \rightarrow N_Q(\lambda, \alpha)$ is a symplectic resolution of singularities.

For a dimension vector $\alpha \in \mathbb{N}^I$ let $\mathfrak{h}_\alpha$ be the subspace of $\mathbb{K}^I$ consisting of λ with $\lambda \cdot \alpha = 0$. If $\lambda \in \mathfrak{h}_\alpha$ is such that $\lambda \notin \mathfrak{h}_\beta$ for all $0 < \beta < \alpha$ (meaning $\alpha - \beta$ and β are sums of positive roots), then $\alpha \in \Sigma_\lambda$ is equivalent to α being a positive root. In this case all elements of $\mu_\alpha^{-1}(\lambda)$ are simple Π^λ modules ([CH], Lemma 4.1). It follows that all points in $\mu_\alpha^{-1}(\lambda)$ are θ -stable, so $N_Q(\lambda, \alpha)$ is equal to $N_Q^\theta(\lambda, \alpha)$ and is smooth. If α is indivisible, the set of such λ is non-empty and by definition is the complement of $\mathfrak{h}_\alpha$ to the union of hyperplanes $\mathfrak{h}_\alpha \cap \mathfrak{h}_\beta$ over $0 < \beta < \alpha$.

Let $\alpha \in \Sigma_0$. Then (see Remark 8.2.1) $\alpha \in \Sigma_\lambda$ for all $\lambda \in \mathfrak{h}_\alpha$, map μ_α is flat and produces a flat family $\mu_\alpha^{-1}(\mathfrak{h}_\alpha) \rightarrow \mathfrak{h}_\alpha$. From Corollary 8.2.2 the induced family $\mu_\alpha^{-1}(\mathfrak{h}_\alpha) // G \rightarrow \mathfrak{h}_\alpha$ is equidimensional and is flat as well. If, in addition, α is indivisible, generic fiber is smooth and the whole family admits a simultaneous resolution

$$\pi_{\mathfrak{h}_\alpha}^\theta : \mu_\alpha^{-1}(\mathfrak{h}_\alpha) //_\theta \mathbb{P}G(\alpha) \rightarrow \mu_\alpha^{-1}(\mathfrak{h}_\alpha) // \mathbb{P}G(\alpha),$$

which restricts to resolutions

$$\pi_\lambda^\theta : N_Q^\theta(\lambda, \alpha) \rightarrow N_Q(\lambda, \alpha)$$

for each $\lambda \in \mathfrak{h}_\alpha$.

8.2.3. Étale local structure. Let (A, M, ω, ζ) be a quadruple (see Definition 8.2.1). Recall that we have $G = A^\times$ acting on M and with a moment map μ given by $\langle \mu(m), a \rangle = \omega(m, am)$. Points of $\mu^{-1}(\zeta) // G$ correspond to closed G -orbits on $\mu^{-1}(\zeta)$. Let $x \in \mu^{-1}(\zeta)$ be a point whose G -orbit $G \cdot x$ is closed. Note that the tangent space to the orbit is given by $[A, x] \subset M$ if we identify $T_x M$ with M .

Lemma 8.2.3 ([CrB1], Lemma 1). *$[A, x]$ is an isotropic subspace of M . In other words $[A, x] \subseteq [A, x]^\perp$.*

Proof.

$$\begin{aligned} \omega([a, x], [b, x]) &= \omega(ax, bx) - \omega(ax, xb) - \omega(xa, bx) + \omega(xa, xb) = \\ &= \omega(axb, x) + \omega(xb, ax) - \omega(x, abx) + \omega(x, axb) = \\ &= \omega(x, bax) - \omega(x, abx) = \langle \mu(x), ba - ab \rangle = \zeta(ba - ab) = 0 \end{aligned}$$

□

Moreover by Matsushima's theorem [Ma] the stabiliser G_x is reductive and from this it follows that $A_x = \{a \in A | ax - xa\}$ is a semisimple algebra ([CrB1], Lemma 4.2).

Let now L be an A_x - A_x -bimodule complement to A_x in A : $A = A_x \oplus L$. Note that $[A, x]$ is an A_x - A_x -subbimodule of M , so by Corollary 8.2.1 we can choose a coisotropic A_x - A_x -bimodule complement C , so that $M = [A, x] \oplus C$. Let $W = C \cap [A, x]^\perp$. Let $\mu_x : M \rightarrow A_x^*$ be the map obtained by composing μ with the restriction map $A^* \rightarrow A_x^*$. Let $\hat{\mu}$ denote the restriction of μ_x to W .

Lemma 8.2.4 ([CrB1], Lemma 4.3). *The restriction of symplectic form ω to W is non-degenerate and hence a symplectic form. It is balanced for the A_x - A_x -bimodule structure and the corresponding moment map is $\hat{\mu}$.*

This way each point $x \in M$ with a closed G -orbit provides a new quadruple (A_x, W, ω, ζ) .

Crawley-Boevey then defines a map $\nu : C \rightarrow L^*$ by $\langle \nu(c), l \rangle = \omega(c, lx) + \omega(c, lc) + \omega(x, lc)$. The map is defined so that

$$\langle \mu(x+c), a+l \rangle = \zeta(a+l) + \langle \mu_x(c), a \rangle + \langle \nu(c), l \rangle$$

for $c \in C$, $a \in A_x$ and $l \in L$. There is an important lemma:

Lemma 8.2.5 ([CrB1], Lemma 4.4). *The assignment $c \mapsto x+c$ induces a G_x -equivariant map $\mu_x^{-1}(0) \cap \nu^{-1}(0) \rightarrow \mu^{-1}(\zeta)$, and the induced map $(\mu_x^{-1}(0) \cap \nu^{-1}(0))//G_x \rightarrow \mu^{-1}(\zeta)//G$ is étale at 0.*

Proof. The formula above shows that $x+c \in \mu^{-1}(\zeta)$ if and only if $c \in \mu_x^{-1}(0) \cap \nu^{-1}(0)$. The action of G on M gives a map $\phi : G \times_{G_x} C \rightarrow M$ which is étale at $(1, 0)$. Then Luna's Fundamental Lemma ([Lu], p.94) applies to ϕ : there exists $U \subset X = G \times_{G_x} C$ which is saturated for $\pi_X : X \rightarrow X//G$, such that $\phi|_U$ is étale, $V = \phi(U)$ is an affine open subset of M , saturated for $\pi_M : M \rightarrow M//G$, the corresponding morphism $\phi/G : U//G \rightarrow V//G$ is étale and induces a G -equivariant isomorphism $U \xrightarrow{\sim} V \times_{V//G} U//G$.

The statement of the lemma is then obtained by the base change to the closed subvariety $V' = V \cap \mu^{-1}(\zeta) \subset V$ which is G -invariant. Namely $U' = V' \times_{V//G} U//G$ is actually $X' \cap U$, where $X' = G \times_{G_x} C'$ and $C' = \mu_x^{-1}(0) \cap \nu^{-1}(0)$. It remains to note that $X//G \cong C//G_x$ and that $X'//G \cong C'//G_x$. $\square$

We would like to lift this étale map to the level of GIT quotients. Let us fix a character θ of G , then its restriction to G_x gives a character θ_x . Let C be a vector space with G_x -action and consider its GIT quotient

$$C//_{\theta_x} G_x = \text{Proj} \bigoplus_{n=0}^{\infty} \Gamma(C, \mathcal{O}_C)^{\theta_x^n},$$

where $\Gamma(C, \mathcal{O}_C)^{\theta_x^n}$ is the subspace of functions which are multiplied by $\theta_x(g)^n$ under the action of $g \in G_x$ for all g . We also have $X = G \times_{G_x} C$ and, since G_x is reductive, X is affine. Consider the corresponding GIT-quotient with respect to θ :

$$X//_{\theta} G = \text{Proj} \bigoplus_{n=0}^{\infty} \Gamma(X, \mathcal{O}_X)^{\theta^n}.$$

Lemma 8.2.6. *These two GIT quotients coincide: $X//_{\theta} G = C//_{\theta_x} G_x$*

Proof. $\Gamma(X, \mathcal{O}_X) = (\Gamma(G, \mathcal{O}_G) \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C))^{G_x}$. For each character θ of G the subspace $\Gamma(G, \mathcal{O}_G)^{\theta}$ is one-dimensional and is spanned by the character itself, considered as a function $\theta : G \rightarrow \mathbb{A}^1$. We have a G -equivariant embedding $\Gamma(X, \mathcal{O}_X) \hookrightarrow \Gamma(G, \mathcal{O}_G) \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C)$, corresponding to the G -equivariant map $G \times C \rightarrow X$. In the tensor product G acts only

on $\Gamma(G, \mathcal{O}_G)$ and so $\Gamma(X, \mathcal{O}_X)^\theta$ is actually a subspace of $\mathbb{K} \cdot \theta \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C)$. Moreover, since $\Gamma(X, \mathcal{O}_X) = (\Gamma(G, \mathcal{O}_G) \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C))^{G_x}$, it is equal to its G_x -invariants $(\mathbb{K} \cdot \theta \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C))^{G_x}$. Action of G_x on $G \times C$ is given by $g_x \circ (g, c) = (g \cdot g_x, g_x^{-1} \circ c)$. Since $g_x \circ \theta = \theta_x(g_x) \cdot \theta$, we have $g_x \circ (\theta \otimes f) = \theta_x(g_x) \cdot (\theta \otimes (g_x^{-1} \circ f))$, and so for $(\theta \otimes f)$ to be G_x -invariant we need f to be rescaled by $\theta_x(g_x)$ under the action of g_x . In other words, we get that $\Gamma(X, \mathcal{O}_X) = \mathbb{K} \cdot \theta \otimes_{\mathbb{K}} \Gamma(C, \mathcal{O}_C)^{\theta_x}$. This way we obtain an isomorphism

$$\bigoplus_{n=0}^{\infty} \Gamma(C, \mathcal{O}_C)^{\theta_x^n} \xrightarrow{\sim} \bigoplus_{n=0}^{\infty} \Gamma(X, \mathcal{O}_X)^{\theta^n} \text{ where } f \mapsto \theta^n \otimes f, \text{ for } f \in \Gamma(C, \mathcal{O}_C)^{\theta_x^n},$$

and so $X //_{\theta} G = C //_{\theta_x} G_x$. $\square$

Note that the isomorphism is constructed in a way such that the square

$$\begin{array}{ccc} X //_{\theta} G & \longrightarrow & C //_{\theta_x} G_x \\ \downarrow & & \downarrow \\ X // G & \longrightarrow & C // G_x \end{array}$$

is commutative. So Lemma 8.2.6 shows that in the proof of Lemma 8.2.5 we can replace the categorical quotient with a GIT-quotient with respect to some character θ (or θ_x correspondingly) and extend the étale map to the map of resolutions. Being even more precise, following the terminology of Sect. 5.2, we obtain an étale equivalence of pointed resolutions

$$(\pi, C' //_{\theta_x} G_x, C' // G_x, 0) \sim_{et} (\pi_{\lambda}^{\theta}, N_Q(\lambda, \alpha)^{\theta}, N_Q(\lambda, \alpha), x),$$

where $C' = \mu_x^{-1}(0) \cap \nu^{-1}(0)$.

Lemma 8.2.7 ([CrB1], Lemma 4.8). *There is a morphism $(\mu_x^{-1}(0) \cup \nu^{-1}(0)) // G_x \rightarrow \hat{\mu}^{-1}(0) // G_x$, sending 0 to 0, which is étale at 0.*

Proof. (Sketch) One first proves that $C = C^{\perp} \oplus W$ as A_x - A_x -bimodules and that the restriction of the corresponding projection $p : C \rightarrow W$ to $\nu^{-1}(0) \subset C$ is étale at 0. Then again, using Luna's Fundamental Lemma, there is an open subvariety U' of $\nu^{-1}(0) // G_x$ containing 0, such that the restriction of $p/G_x : \nu^{-1}(0) // G_x \rightarrow W // G_x$ to U' is étale and the image $V' = (p/G_x)(U')$ is an open affine subset.

The étale map from the statement of the lemma is then obtained by the base change to $\hat{\mu}^{-1}(0) \subset W$. Its preimage in $\nu^{-1}(0)$ is exactly $\mu_x^{-1}(0) \cup \nu^{-1}(0)$. $\square$

Note that, following the proof, the desired map was induced by an étale map $U' \rightarrow V' \times_{W // G_x} \nu^{-1}(0) // G_x$ which also gives an étale map between the open subsets of θ_x -semistable points. This way it extends to an étale map between corresponding resolutions. We obtain an étale equivalence

$$(\pi, C' //_{\theta_x} G_x, C' // G_x, 0) \sim_{et} (\hat{\pi}, \hat{\mu}^{-1}(0) //_{\theta_x} G_x, \hat{\mu}^{-1}(0) // G_x, 0).$$

Composing it with the previous one, we get the following:

Proposition 8.2.2. *There exists an étale equivalence of pointed resolutions*

$$(\pi_{\lambda}^{\theta}, N_Q(\lambda, \alpha)^{\theta}, N_Q(\lambda, \alpha), x) \sim_{et} (\hat{\pi}, \hat{\mu}^{-1}(0) //_{\theta_x} G_x, \hat{\mu}^{-1}(0) // G_x, 0).$$

Recall that by Lemma 8.2.2 every quadruple (A, M, ω, ζ) is equivalent to a one coming from the quiver. Let Q_x be the quiver associated to the quadruple $(A_x, W, \omega, 0)$. We also have the corresponding dimension vector $\alpha_x \in \mathbb{N}^{I_x}$, where I_x is the set of vertices of Q_x . We can then reformulate the proposition above in the following way:

Proposition 8.2.3. *Let Q be a quiver with a vertex set I , $\alpha \in \mathbb{N}^I$ be a dimension vector and λ a vector in $\mathbb{K}^I$. Then there exist a quiver Q_x with a vertex set I_x and a dimension vector α_x , such that*

$$(\pi_\lambda^\theta, N_Q(\lambda, \alpha)^\theta, N_Q(\lambda, \alpha), x) \sim_{et} (\pi_0^{\theta_x}, N_{Q_x}(0, \alpha_x)^{\theta_x}, N_{Q_x}(0, \alpha_x), 0)$$

Proposition says that the resolution of singularities at any point of a Marsden-Weinstein reduction for representations of quivers étale locally looks like the resolution of singularities at the central point of some other Marsden-Weinstein reduction.

8.2.4. $\mathbb{G}_m$ -actions. Let $\lambda = 0$ and let's consider some $N_Q(0, \alpha)$. Recall that it was obtained as a Hamiltonian reduction from a symplectic vector space $\text{Rep}(\overline{Q}, \alpha)$. Let's now consider a $\mathbb{G}_m$ -action on $\text{Rep}(\overline{Q}, \alpha)$ which rescales all the operators in the representation, namely $t \circ x_a = tx_a$ for any arrow $a \in \overline{Q}$. This action commutes with the action of $\text{GL}(\alpha)$ and rescales ω : we have $t \circ \omega = t^2 \omega$. Though this action does not preserve the fibers of μ , it preserves $\mu^{-1}(0)$ and so defines an action on it. Moreover this action is contracting: the whole $\text{Rep}(\overline{Q}, \alpha)$ and $\mu^{-1}(0)$ in particular are contracted to a single point 0. Since this action commutes with the action of $\text{GL}(\alpha)$ it descends to a contracting action on $N_Q(0, \alpha)$. Moreover it also defines an action on $N_Q(0, \alpha)^\theta$ and turns $\pi_0 : N_Q(0, \alpha)^\theta \rightarrow N_Q(0, \alpha)$ into a conical resolution.

Combining this with the Proposition 8.2.3 we get that any point $x \in N_Q(\lambda, \alpha)$ has a conical étale neighbourhood (see Definition 5.4.2). In other words:

Theorem 8.2.4. *Resolution $\pi_0 : N_Q(0, \alpha)^\theta \rightarrow N_Q(0, \alpha)$ of singularities of Marsden-Weinstein reduction is a resolution with conical slices.*

8.2.5. *Example: Nakajima quiver varieties.* The main example of Marsden-Weinstein reductions for representations of quivers are Nakajima quiver varieties. Here we follow the *Remarks added after 2000* in [CrB2].

Let Q_0 be a quiver with a vertex set I . For $\underline{v}, \underline{w} \in \mathbb{N}^I$, let $M(\underline{v}, \underline{w})$ be the space

$$\text{Rep}(\overline{Q}_0, \underline{v}) \oplus \bigoplus_{k \in I} \text{Mat}(v_k \times w_k, \mathbb{K}) \oplus \bigoplus_{k \in I} \text{Mat}(w_k \times v_k, \mathbb{K}).$$

We denote the coordinates on this space by (B, i, j) . The natural action of $G_{\underline{v}} = \prod_{k \in I} \text{GL}(v_k)$ on $M(\underline{v}, \underline{w})$ has a moment $\mu : M(\underline{v}, \underline{w}) \rightarrow \bigoplus_{k \in I} \text{Mat}(v_k, \mathbb{K})$, whose k -th component is given by the map sending (B, i, j) to

$$\sum_{\substack{a \in Q_0 \\ h(a)=k}} B_a B_{a^*} - \sum_{\substack{a \in Q_0 \\ t(a)=k}} B_{a^*} B_a + \sum_{k \in I} i_k j_k.$$

Affine Nakajima quiver variety $\mathfrak{M}_0(\underline{v}, \underline{w})$ is defined as $\mu^{-1}(0) // G_{\underline{v}}$. More generally, Nakajima quiver varieties $\mathfrak{M}_\lambda^\theta(\underline{v}, \underline{w})$ are defined as $\mu^{-1}(\lambda) //_\theta G_{\underline{v}}$.

Let's now consider a quiver Q , obtained from Q_0 by adjoining a new vertex ∞ and w_k arrows from ∞ to k for each $k \in I$. Let α be the dimension vector for Q , whose restriction to I is equal to $\underline{v}$ and such that $\alpha_\infty = 1$. Dividing the matrices in $\text{Mat}(v_k \times w_k, \mathbb{K})$ into their columns and the matrices in $\text{Mat}(w_k \times v_k, \mathbb{K})$ into their rows, we get an identification $M(\underline{v}, \underline{w}) \cong \text{Rep}(\overline{Q}, \alpha)$. Since $\alpha_\infty = 1$, we also get that $\mathbb{P}G(\alpha) = (G_{\underline{v}} \times \mathbb{G}_m) / \mathbb{G}_m \cong G_{\underline{v}}$. It is easy to see that moment maps coincide too and that

$$\mathfrak{M}_\lambda^\theta(\underline{v}, \underline{w}) \cong N_Q(\lambda, \alpha)^\theta$$

for all λ and θ . Assume that the map $\pi_0^\theta : \mathfrak{M}_0^\theta(\underline{v}, \underline{w}) \rightarrow \mathfrak{M}_0(\underline{v}, \underline{w})$ is a resolution of singularities (e.g. when the conditions of Proposition 8.2.1 are satisfied for α , λ and Q . Note that α is indivisible, so it is enough to check that $\alpha \in \Sigma_\lambda$). We obtain the following theorem:

Theorem 8.2.5. *If the map $\pi_0^\theta : \mathfrak{M}_0^\theta(\underline{v}, \underline{w}) \rightarrow \mathfrak{M}_0(\underline{v}, \underline{w})$ is a resolution of singularities, it is a resolution with conical slices.*

8.3. Hamiltonian reductions of vector spaces. Marsden-Weinstein reduction for representations of a quiver is a particular case of an algebraic Hamiltonian reduction of a vector space (which can be called Marsden-Weinstein reduction as well). Namely let (V, ω) be a vector space with a symplectic form and let $G \rightarrow \mathrm{Sp}(V)$ be a symplectic representation of a reductive group G over $\mathbb{K}$. We will assume that the corresponding action is effective, namely that the map $G \rightarrow \mathrm{Sp}(V)$ is an embedding. Note that the action of G on V also induces an action of its Lie algebra $\mathfrak{g}$.

Action of G on V defines an action on the corresponding affine space, which is naturally a symplectic manifold. The symplectic G -action on V always has an algebraic moment map $\mu : V \rightarrow \mathfrak{g}^*$. It is given by the following formula:

$$\langle \mu(v), a \rangle = \frac{1}{2} \omega(v, a \circ v)$$

for any $v \in V$ and $a \in \mathfrak{g}$.

Let $Z(\mathfrak{g}^*) = (\mathfrak{g}^*)^G \subset \mathfrak{g}^*$ denote the subspace of the invariants of the coadjoint action. For $\lambda \in Z(\mathfrak{g}^*)$, the subvariety $\mu^{-1}(\lambda)$ is stable under the G -action. Given a character $\theta : G \rightarrow \mathbb{G}_m$ we can consider the GIT quotient with respect to θ :

$$\mathfrak{M}(G, V)_\lambda^\theta = \mu^{-1}(\lambda) //_\theta G.$$

We will omit the superscript θ in the case when $\theta = 0$. For any θ we have a projective morphism $\pi_\lambda^\theta : \mathfrak{M}(G, V)_\lambda^\theta \rightarrow \mathfrak{M}(G, V)_\lambda$.

8.3.1. Darboux-type theorems. Unfortunately the analogue of Theorem 8.2.1 is not true in this context. The important difference here is that an irreducible subrepresentation of V does not need to be isotropic. In particular maximal isotropic G -subrepresentation does not need to be maximal isotropic as a vector space.

Nevertheless the weaker form of Darboux theorem is still true:

Proposition 8.3.1. *Let (V, ω) be a symplectic representation of a reductive group G . Let $S \subset V$ be a maximal isotropic G -submodule, which is Lagrangian: $S = S^\perp$. Then S has an isotropic G -equivariant complement which is also Lagrangian. Moreover V is isomorphic to $S \oplus S^*$ as a symplectic representation of G .*

Proof. Since any representation of G is completely reducible, S has a G -equivariant complement C . S is Lagrangian, so $\dim V = 2 \dim S$ and $\dim C = \dim S$. Since G preserves the symplectic form on V , the natural map $C \rightarrow S^*$ induced by ω is G -equivariant. Moreover as S is Lagrangian and ω is non-degenerate, this map is an isomorphism.

Let $f : S \rightarrow C^*$ be the dual map, which is an isomorphism as well. The restriction of ω to C induces a G -equivariant map $\tilde{\omega}|_C : C \rightarrow C^*$. Let $\theta : C \rightarrow S$ be equal to $(f)^{-1} \circ \tilde{\omega}|_C$. We have $\tilde{\omega}|_C = f \circ \theta$ and so $\omega(\theta c, c') = \omega(c, c')$ for all $c, c' \in C$.

Let $D = \{c - \frac{1}{2}\theta(c) \mid c \in C\}$. Since θ was G -equivariant this is a G -invariant subspace. Clearly $D \cap S = 0$ and $V = S \oplus D$. Now

$$\begin{aligned} \omega(c - \frac{1}{2}\theta(c), c' - \frac{1}{2}\theta(c')) &= \omega(c, c') - \omega(c, \frac{1}{2}\theta(c')) - \omega(\frac{1}{2}\theta(c), c') + \omega(\frac{1}{2}\theta(c), \frac{1}{2}\theta(c')) = \\ &= \omega(c, c') - \frac{1}{2}\omega(c, c') - \frac{1}{2}\omega(c, c') + 0 = 0. \end{aligned}$$

So D is isotropic and has dimension $\frac{1}{2} \dim V$, so is Lagrangian. The map $\tilde{\omega}$ induces a G -equivariant isomorphism $D \xrightarrow{\sim} S^*$ and so $V = S \oplus S^*$.

□

Remark 8.3.1. Let $T \subset V$ be an irreducible G -submodule such that $\omega|_T \neq 0$. Then, by irreducibility, T is symplectic and $\tilde{\omega}_T : T \xrightarrow{\sim} T^*$. For any self-dual irreducible representation T one can compute its Frobenius-Schur indicator $\text{FS}(T)$ [Bou]. Since ω is skew-symmetric and G -invariant, we get that $\text{FS}(T) = -1$ or in other words that T is *quaternionic*. From this it is easy to show that if all irreducible subrepresentations are *ortogonal* or *isotropic* (meaning $\text{FS}(T) = 1$ or $\text{FS}(T) = 0$) then any maximal isotropic subrepresentation is Lagrangian and the full statement of Theorem 8.2.1 holds. This way ([CrB1], Lemma 2.1) can be considered as a statement about Frobenius-Schur indicators of self-dual representations coming from quivers.

With this weaker version of Darboux theorem we still are able to prove the analogue of Corollary 8.2.1. We will first need to prove a simple lemma:

Lemma 8.3.1. *Let $S \subset V$ be an isotropic subrepresentation. Let C be a G -equivariant complement and let W be the intersection $S^\perp \cap C$. Then the restriction of ω to W is non-degenerate and W is symplectic.*

Proof. S is isotropic, so $S \subset S^\perp$. Since C is a complement to S , $S^\perp \cong S \oplus W$. Moreover ω is non-degenerate on $V \oplus V^*$, so $(S^\perp)^\perp = S$. It follows that S is the kernel of the restriction of ω to $S^\perp$ and so the restriction of ω to W should be non-degenerate. □

Note that as ω is preserved by G , $S^\perp$ is G -invariant and so W is a subrepresentation of G .

Proposition 8.3.2. *Let $S \subset V$ be an isotropic subrepresentation. Then there exists a G -equivariant complement C which is coisotropic: $C^\perp \subset C$. In this case $C = W \oplus \text{Ker } \omega|_C$.*

Proof. Let's start with any G -equivariant complement C and consider $W = C \cap S^\perp$. By the lemma above W is symplectic. It follows that $W^\perp$ is symplectic too, and $W \cap W^\perp = 0$.

Note that $\dim W = \dim V - 2 \dim S$. Indeed, by definition of C , $\dim C = \dim V - \dim S$. We have $S \subset S^\perp$ and so $S^\perp + C = V$. Since $W = C \cap S^\perp$ and $\dim S^\perp = \dim V - \dim S$, we get that:

$$\dim W = \dim C + \dim S^\perp - \dim V = \dim V - \dim S + \dim V - \dim S - \dim V = \dim V - 2 \dim S.$$

From this we obtain that $\dim W^\perp = 2 \dim S$. It follows that $S \subset W^\perp$ is a Lagrangian subrepresentation. Applying Theorem 8.3.1 to $S \subset W^\perp$ we know that there exists a G_x -equivariant Lagrangian complement $D \cong S^*$: $W^\perp = S \oplus D$.

Now let's take $C' = W \oplus D$. Clearly C' is a G_x -equivariant complement to S . Also, $(C')^\perp = W^\perp \cap D^\perp = D \subset C'$, so C' is coisotropic. Note that D is exactly $\text{Ker } \omega|_{C'}$.

Finally if C is any coisotropic G -equivariant complement, by the same argument we get that $W^\perp \cap C$ is equal to $\text{Ker } \omega|_C$ and so $C = W \oplus \text{Ker } \omega|_C$. □

8.3.2. Étale local structure. We will generalise Crawley-Boevey's result to the case of the Hamiltonian reduction of a cotangent bundle of a vector space.

For any vector space V , the direct sum $V \oplus V^*$ has a natural symplectic form: $\omega((z_1, w_1), (z_2, w_2)) = \langle z_1, w_2 \rangle - \langle z_2, w_1 \rangle$. Action of $\text{GL}(V)$ on V preserves the pairing $\langle \cdot, \cdot \rangle$, so given any linear representation $G \rightarrow \text{GL}(V)$, the induced representation $G \rightarrow \text{GL}(V \oplus V^*)$ is symplectic. The space $V \oplus V^*$ is identified with the total space T^*V of the cotangent bundle on V .

Let $x \in \mu^{-1}(\lambda)$ be a point with a closed G -orbit. Then, by the theorem of Matsushima [Ma], the stabiliser G_x is a reductive group. Let $\mathfrak{g}_x$ be the Lie algebra of G_x ; group G_x acts by conjugation on $\mathfrak{g}$, preserving $\mathfrak{g}_x \subset \mathfrak{g}$. Let L be a G_x -equivariant complement to $\mathfrak{g}_x$ inside $\mathfrak{g}$: $\mathfrak{g} \cong \mathfrak{g}_x \oplus L$ as representations of G_x . Such L exists because G_x is reductive. Dually we have

a G_x -equivariant decomposition $\mathfrak{g}^* \cong \mathfrak{g}_x^* \oplus L^*$. We have a natural restriction map $r : \mathfrak{g}^* \rightarrow \mathfrak{g}_x^*$ and it is easy to see that $r(Z(\mathfrak{g}^*))$ is a subspace of $Z(\mathfrak{g}_x^*)$.

G_x acts on V , preserving x , moreover it preserves the subspace $\mathfrak{g} \circ x \cong \mathfrak{g}/\mathfrak{g}_x$, which is identified with a tangent space to the G -orbit of x .

Lemma 8.3.2. $\mathfrak{g}/\mathfrak{g}_x \subset V \oplus V^*$ is an isotropic subspace: $\mathfrak{g}/\mathfrak{g}_x \subset (\mathfrak{g}/\mathfrak{g}_x)^\perp$

Proof. The pairing $\langle \cdot, \cdot \rangle$ between V and V^* is being preserved by G , so $\langle a \circ z, w \rangle + \langle z, a \circ w \rangle = 0$ for any $a \in \mathfrak{g}$, $z \in V$, $w \in V^*$. Now

$$\begin{aligned} \omega(a \circ (z, w), b \circ (z, w)) &= \langle a \circ z, b \circ w \rangle - \langle b \circ z, a \circ w \rangle = \\ &= \langle b \circ (a \circ z), w \rangle - \langle a \circ (b \circ z), w \rangle = \mu(z, w)([b, a]) = \lambda([b, a]) = 0 \end{aligned}$$

as $\lambda : \mathfrak{g}_x \rightarrow \mathbb{K}$ is Ad-invariant. $\square$

Remark 8.3.2. This lemma is the main reason why we consider only symplectic representations of the form $V \oplus V^*$. In general setting, if the subspace $\mathfrak{g}/\mathfrak{g}_x$ is isotropic all arguments below are still valid and one can obtain the similar results about the étale local structure.

Let C be a G_x -equivariant coisotropic complement to $\mathfrak{g} \circ x$ inside $V \oplus V^*$. By Proposition 8.3.2 such C exists. We have a G_x -equivariant decomposition $V \oplus V^* \cong \mathfrak{g}/\mathfrak{g}_x \oplus C$. Let W be the intersection $C \cap (\mathfrak{g}/\mathfrak{g}_x)^\perp$. By Lemma 8.3.1 the restriction of ω to W is non-degenerate and so W is a symplectic representation of G_x .

Recall that we have the moment map $\mu : T^*V \rightarrow \mathfrak{g}^*$. Let μ_x be its composition with the restriction $r : \mathfrak{g}^* \rightarrow \mathfrak{g}_x^*$ and let $\hat{\mu}$ be the restriction of μ_x to W .

We define the map $\nu : C \rightarrow L^*$ by $\langle \nu(c), l \rangle = \frac{1}{2}(\omega(c, l \circ x) + \omega(c, l \circ c) + \omega(x, l \circ c))$. It is easy to check that for any $a \in \mathfrak{g}_x$ we have

$$\langle \mu(x + c), a + l \rangle = \lambda(a + l) + \langle \mu_x(c), a \rangle + \langle \nu(c), l \rangle.$$

Recall that $\mathfrak{g}^* \cong \mathfrak{g}_x^* \oplus L^*$. Formula above can be viewed as the fact that the moment map $\mu : T^*V \rightarrow \mathfrak{g}^*$ restricted to $x + C$ decomposes as a sum of its $\mathfrak{g}_x^*$ -component μ_x and the L^* -component $\nu : C \rightarrow L^*$.

Lemma 8.3.3. The assignment $c \mapsto x + c$ induces a G_x -equivariant map $\mu_x^{-1}(0) \cap \nu^{-1}(0) \rightarrow \mu^{-1}(\lambda)$ and the induced map $(\mu_x^{-1}(0) \cap \nu^{-1}(0))/G_x \rightarrow \mu^{-1}(\lambda)/G$ is étale.

Proof. Exactly the same argument as in Lemma 8.2.5 works here as well. $\square$

Lemma 8.3.4. The map $\nu : C \rightarrow L^*$ is smooth at 0, so $\nu^{-1}(0)$ is smooth at 0. Its tangent space is identified with W .

Proof. Identifying the tangent spaces at 0 with C and L^* correspondingly, by the definition of ν the differential of ν at 0 is given by

$$\langle d\nu_0(c), l \rangle = \frac{1}{2}(\omega(c, l \circ x) + \omega(x, l \circ c)).$$

As ω is preserved by G , $\omega(x, l \circ c) = -\omega(l \circ x, c) = \omega(c, l \circ x)$. From this

$$\langle d\nu_0(c), l \rangle = \omega(c, l \circ x),$$

and so $\text{Ker } d\nu_0 = C \cap (\mathfrak{g}/\mathfrak{g}_x)^\perp = W$. As $\dim W + \dim L = \dim C$ we get that $d\nu_0$ is surjective and so ν is smooth at 0. $\square$

As C is isotropic $W = C \cap (\mathfrak{g}/\mathfrak{g}_x)^\perp$ is the symplectic part of C and that $C = W \oplus \text{Ker } \omega|_C$. This composition is G_x -equivariant.

Lemma 8.3.5. *There is a morphism $(\mu_x^{-1}(0) \cap \nu^{-1}(0))//G_x \rightarrow \hat{\mu}^{-1}(0)//G_x$ sending 0 to 0 and which is étale at 0.*

Proof. Let $p : C \rightarrow W$ be the projection along $\text{Ker } \omega|_C$. Let $c = w + c' \in W \oplus \text{Ker } \omega|_C$ and $a \in \mathfrak{g}_x$. By the formula for μ_x :

$$\langle \mu_x(c), a \rangle = \frac{1}{2} \omega(w + c', a \circ (w + c')) = \frac{1}{2} \omega(w, a \circ w) = \langle \hat{\mu}(w), a \rangle,$$

so $\mu_x = \hat{\mu} \circ p$. The restriction $\phi : \nu^{-1}(0) \rightarrow W$ of p is étale at 0 by Lemma 8.3.4 and is G_x -equivariant.

Now Luna's Fundamental Lemma can be applied to ϕ : there exists an affine open subvariety U' of $\nu^{-1}(0)//G_x$ containing 0 and such that the restriction of the map

$$\phi//G_x : \nu^{-1}(0)//G_x \rightarrow W//G_x$$

to U' is étale.

As $\mu_x = \hat{\mu} \circ p$, we have $\phi^{-1}(\hat{\mu}^{-1}(0)) = \mu_x^{-1}(0) \cap \nu^{-1}(0)$. So taking the pull-back of $\phi//G_x$ along the closed embedding $\hat{\mu}^{-1}(0)//G_x \hookrightarrow W//G_x$ we get a morphism $(\mu_x^{-1}(0) \cap \nu^{-1}(0))//G_x \rightarrow W//G_x$, which is étale at 0. $\square$

8.3.3. Étale equivalences. Let's assume that a representation $G \rightarrow \text{GL}(V)$, a vector $\lambda \in Z(\mathfrak{g}^*)^G$ and a character $\theta : G \rightarrow \mathbb{G}_m$ are chosen such that the map $\pi_\lambda^\theta : \mathfrak{M}(G, V \oplus V^*)_\lambda^\theta \rightarrow \mathfrak{M}(G, V \oplus V^*)_\lambda$ is a resolution of singularities.

Using Lemma 8.2.6 we can lift the étale maps from Lemma 8.3.3 and Lemma 8.3.5 to the level of the corresponding resolutions. We obtain the following proposition:

Proposition 8.3.3. *Let V be a representation of a reductive group G over $\mathbb{K}$. Let $\lambda \in Z(\mathfrak{g})^G$ and $\theta : G \rightarrow \mathbb{G}_m$ be such that $\pi_\lambda^\theta : \mathfrak{M}(G, V \oplus V^*)_\lambda^\theta \rightarrow \mathfrak{M}(G, V \oplus V^*)_\lambda$ is a resolution of singularities. Then for any point $x \in \mathfrak{M}(G, V \oplus V^*)_\lambda$ there exists a symplectic subspace $W \subset V \oplus V^*$, invariant under the action of stabiliser G_x , such that there is an étale equivalence*

$$(\pi_\lambda^\theta, \mathfrak{M}(G, V \oplus V^*)_\lambda^\theta, \mathfrak{M}(G, V \oplus V^*)_\lambda, x) \sim_{et} (\pi_0^{\theta_x}, \mathfrak{M}(G_x, W)_{0^x}^{\theta_x}, \mathfrak{M}(G_x, W)_{0,0}) .$$

The contracting $\mathbb{G}_m$ -action on W , which is given by $t \circ w = tw$, commutes with the action of G_x and descends to both $\mathfrak{M}(G_x, W)$ and $\mathfrak{M}(G_x, W)^{\theta_x}$, turning $\pi : \mathfrak{M}(G_x, W)^{\theta_x} \rightarrow \mathfrak{M}(G_x, W)$ into a conical resolution of singularities. So Proposition 8.3.3 implies the following theorem:

Theorem 8.3.1. *Let V be a representation of a reductive group G over $\mathbb{K}$. Let $\theta : G \rightarrow \mathbb{G}_m$ be a character such that $\pi_0^\theta : \mathfrak{M}(G, V \oplus V^*)_0^\theta \rightarrow \mathfrak{M}(G, V \oplus V^*)_0$ is a resolution of singularities and assume in addition that $\mathfrak{M}(G, V \oplus V^*)_0$ is normal. Then π_0^θ is a resolution with conical slices.*

Remark 8.3.3. When the representation V is such that the action of G for a general point p on $V \oplus V^*$ is locally trivial (meaning that the stabiliser G_p is finite), it is not hard to see that $p \in \mu^{-1}(\lambda)$ is a smooth point of $\mu^{-1}(\lambda)$ if and only if the stabiliser is finite. Under this assumptions the moment map μ is surjective, and p is smooth if and only if $d\mu_p$ is surjective too. On the other hand, $\langle d\mu_p(v), a \rangle = \omega(v, a_p)$, so this is equivalent to $a_p \neq 0$ for any $a \in \mathfrak{g}$. In other words the map $a \mapsto a_p$ from $\mathfrak{g}$ to $T_p(T^*V)$ should be an embedding and consequently the stabiliser G_p should be finite. If now $p \in \mu^{-1}(\lambda)^{\theta-ss}$, then this condition is equivalent to p being not only θ -semistable, but θ -stable as well. We get that $\mathfrak{M}(G, V \oplus V^*)_\lambda^\theta$ is smooth if and only if all θ -semistable points are θ -stable. Note that this is not enough though to guarantee that the map π_λ^θ is a resolution of singularities: we need also to know that it is birational. This seems to be hard to show in general.

The first potential issue is that $\mathfrak{M}(G, V \oplus V^*)_\lambda^\theta$ can appear to be empty (when the θ -semistable locus $\mu^{-1}(\lambda)^{\theta-ss} \subset \mu^{-1}(\lambda)$ is empty). A possible way to check that it is not, is to use hyperkähler geometry, namely a consequence of Kempf-Ness theorem (for Kempf-Ness theorem see [KN], for the consequence see e.g. Lemma 3 in [CrB4]) which states that over $\mathbb{C}$ varieties $\mu^{-1}(\theta)//G$ and $\mu^{-1}(0)//_\theta G$ are diffeomorphic (here some additional conditions on G , V , θ and λ should be imposed). In particular one is empty if and only if the other one is.

Even bigger potential issue is that the map π_λ^θ can appear to be not generically 1-to-1 or surjective. This is equivalent to the set of points with closed orbits being dense in $\mu^{-1}(\lambda)$ together with the existence of a θ -semistable point in each of the irreducible components of $\mu^{-1}(\lambda)$.

Finally, one also needs to check that the affine variety $\mathfrak{M}(G, V \oplus V^*)_\lambda$ is normal. It is also not clear how to check this in general.

Remark 8.3.4. It can easily happen that for a fixed λ there does not exist θ , such that $\mathfrak{M}(G, V \oplus V^*)_\lambda^\theta$ is smooth. For example, if G is semisimple then θ automatically equals to 0 and $\mathfrak{M}(G, V \oplus V^*)_\lambda^\theta = \mathfrak{M}(G, V \oplus V^*)_\lambda$, which is often very singular (especially in the case $\lambda = 0$).

8.3.4. Example: Marsden-Weinstein reductions for representations of quivers. Marsden-Weinstein reduction $N_Q(\lambda, \alpha)$ is a particular example of $\mathfrak{M}(G, V \oplus V^*)_\lambda$. Indeed a choice of an orientation on Q provides an identification of $\text{Rep}(\overline{Q}, \alpha)$ with $\text{Rep}(Q, \alpha) \oplus \text{Rep}(Q, \alpha)^*$ as a representation of $\mathbb{P}G(\alpha)$. It is easy to see that after that the two constructions literally coincide. So $N_Q^\theta(\lambda, \alpha) = \mathfrak{M}(\mathbb{P}G(\alpha), \text{Rep}(Q, \alpha) \oplus \text{Rep}(Q, \alpha)^*)_\lambda^\theta$.

In this case, Proposition 8.2.1 provides the sufficient conditions on Q , λ and α that guarantee that the map π_λ^θ is a resolution of singularities.

8.3.5. Example: hypertoric varieties. Hypertoric varieties are a particular case of Hamiltonian reduction, namely in the case when the reductive group G is an algebraic torus. Hypertoric varieties (over $\mathbb{C}$) are hyperkähler analogues of toric varieties. Their geometry is related to complicated combinatorics of polytops, such as zonotopal tilings [AP] and Stanley-Reisner rings of simplicial complexes ([HS], [PW]).

Let $V = \mathbb{K}^n$ be a representation of the k -dimensional algebraic torus $T^k \cong \mathbb{G}_m^k$. After a change of coordinates we can assume that the action of T^k is given by an embedding $\iota : T^k \rightarrow T^n$, where T^n is a standard n -dimensional torus, acting on V . We assume that the representation is faithful or in other words that ι is an embedding.

The quotient T^n/T^k is also an algebraic torus, namely $T^n/T^k \cong T^d$, where $d = n - k$. Let $X_*(T^n) = \text{Hom}_{\mathbb{K}}(\mathbb{G}_m, T^n)$ be the cocharacter lattice. We have a non-canonical isomorphism $X_*(T^n) \cong \mathbb{Z}^n$. A short exact sequence of tori

$$1 \rightarrow T^k \rightarrow T^n \rightarrow T^d \rightarrow 1$$

induces a short exact sequence of cocharacters:

$$0 \rightarrow X_*(T^k) \rightarrow X_*(T^n) \rightarrow X_*(T^d) \rightarrow 0.$$

For each T^n we also have its character lattice $X^*(T) = \text{Hom}_{\mathbb{K}}(T^n, \mathbb{G}_m)$. The lattice $X^*(T)$ is canonically dual to $X_*(T)$ under the pairing given by the composition of a cocharacter with a character (this composition lies in $\text{Hom}_{\mathbb{K}}(\mathbb{G}_m, \mathbb{G}_m) \cong \mathbb{Z}$). Let $\{x_1, \dots, x_n\}$ be a basis of $X_*(T^n)$ and let $\{e_1, \dots, e_n\}$ be the dual basis of $X^*(T^n)$. Let $\{a_1, \dots, a_n\} \in X_*(T^d)$ be the images of $\{x_1, \dots, x_n\}$. The map $X_*(T) \rightarrow X_*(T)$ is surjective, so $\{a_1, \dots, a_n\}$ span $X_*(T^d)$. The set $\{a_1, \dots, a_n\} \in X_*(T^d)$ uniquely defines the map $\iota : T^k \rightarrow T^n$.

The hypertoric variety $\mathfrak{H}\mathfrak{T}^\alpha$ is defined as $\mathfrak{M}(T^k, V \oplus V^*)_0^\alpha$ in the notations of Sect. 8.3. Note that α here is a character of T^k and so is an element of the lattice $X^*(T^k)$. Let $r = (r_1, \dots, r_n) \in X_*(T^n)$ be any lift of α with respect to $\iota^* : X^*(T^n) \rightarrow X^*(T^k)$.

The data which we used to construct $\mathfrak{H}\mathfrak{T}_\lambda^\alpha$ can be encoded in terms of an arrangement of affine hyperplanes in $X^*(T^d)$ endowed with some additional structure. Namely a *weighted, cooriented* hyperplane $H \subseteq X^*(T^d)$ is an affine hyperplane along with a choice of a nonzero $a \in X_*(T^d) \cong (X^*(T^d))^*$, such that $a|_H = 0$. For each $i = 1, \dots, n$ we have a weighted cooriented hyperplane

$$H_i = \{v \in X^*(T^d) \mid \langle v, a_i \rangle + r_i = 0\}$$

with the weight and coorientation given by $a_i \in X_*(T^d)$. We will call this hyperplane arrangement by $\mathcal{A}$. A choice of a different lift of α , say r' , corresponds to a simultaneous translation of all hyperplanes in $\mathcal{A}$ by the vector $r' - r \in X_*(T^d)$. This arrangement uniquely defines the map $\iota : T^k \rightarrow T^n$ along with the choice of α and we call by $\mathfrak{H}\mathfrak{T}(\mathcal{A})$ the corresponding hypertoric variety.

An arrangement $\mathcal{A}$ is called *simple* if for every subset of m hyperplanes in $\mathcal{A}$ their intersection has codimension m .

Remark 8.3.5. Here we use the convention that $\dim \emptyset = -\infty$ and so the codimension of any empty intersection is $+\infty$.

An arrangement $\mathcal{A}$ is called *unimodular* if any collection of d linearly independent weight-orientation vectors $\{a_{i_1}, \dots, a_{i_d}\}$ spans $X_*(T^d)$ over $\mathbb{Z}$. Note that the condition of being unimodular does not depend on the choice of character α . If the arrangement is simple and unimodular, it is called *smooth*. The following theorem provides a reason for that:

Theorem 8.3.2 ([BD], Theorem 3.2). *The variety $\mathfrak{M}(\mathcal{A})$ is smooth if and only if $\mathcal{A}$ is smooth.*

Remark 8.3.6. In the cited paper this fact is proven for the hyperkähler reduction and the statement we need follows by the hyperkähler trick which establishes the diffeomorphism of $\mathfrak{M}(\mathcal{A})_{\mathbb{C}}$ and $\mu^{-1}(\alpha)_{\mathbb{C}} // G$ using Kempf-Ness theorem. Here the character α is also considered as an integral element of $(t^k)^*$.

Now consider the case when the character α is equal to 0. In this case the intersection of all hyperplanes from $\mathcal{A}$ is the point $0 \in X^*T^d$, so $\mathcal{A}$ is not simple unless $n = d$ and $T^k = \{1\}$. Given an arrangement $\mathcal{A} = \{H_1, \dots, H_n\}$ with $r = 0$ (so that $H_i = \{v \in X^*(T^d) \mid \langle v, a_i \rangle = 0\}$ for some $a_i \in X_*(T^d)$), it is not hard to see that for general $r = (r_1, \dots, r_n) \in X_*(T^n)$, the arrangement $\tilde{\mathcal{A}} = \{\tilde{H}_1, \dots, \tilde{H}_n\}$, where $\tilde{H}_i = \{v \in X^*(T^d) \mid \langle v, a_i \rangle + r_i = 0\}$, is simple. In this case the arrangement $\tilde{\mathcal{A}}$ is called a *simplification* of $\mathcal{A}$. Note that the set of weight-orientation vectors for $\tilde{\mathcal{A}}$ and $\mathcal{A}$ is the same. In particular $\tilde{\mathcal{A}}$ is unimodular if and only if $\mathcal{A}$ is unimodular. Let $\alpha \in X^*(T^k)$ be the image of r under the map $\iota^* : X_*(T^n) \rightarrow X_*(T^d)$.

We have a projective $\pi_{\mathcal{A}} : \mathfrak{M}(\tilde{\mathcal{A}}) \rightarrow \mathfrak{M}(\mathcal{A})$ (given by the map $\pi_0^\alpha : \mathfrak{M}(T^k, V \oplus V^*)_0^\alpha \rightarrow \mathfrak{M}(T^k, V \oplus V^*)_0^0$ in the terminology of Sect. 8.3) which is known to be surjective and generically 1-to-1. If the arrangement $\mathcal{A}$ is unimodular this is a resolution of singularities and the base is normal ([BeKu], Section 4). From Theorem 8.3.1 we obtain the following:

Theorem 8.3.3. *The resolution of singularities $\pi_{\mathcal{A}} : \mathfrak{M}(\tilde{\mathcal{A}}) \rightarrow \mathfrak{M}(\mathcal{A})$ (in the case when $\mathcal{A}$ is unimodular) is a resolution with conical slices.*

Remark 8.3.7. The stabiliser of any non-zero point $x \in T^*V$ in a hypertoric variety is a torus T^s of smaller dimension. Moreover there are no self-dual representations of T^s except that of the trivial one. The vector subspace W from Proposition 8.3.3 can be T^s -equivariantly decomposed as $W = \mathbb{K}^m \oplus W'$, where $\mathbb{K}^m$ is the subspace of invariants and W' is the T^s -equivariant complement. It is easy to see from the definitions that $\mathfrak{M}(T^s, W)_0^\alpha$ can be decomposed as $\mathbb{A}^m \times \mathfrak{M}(T^s, W')_0^\alpha$. Since W' does not have any self-dual subrepresentations, by Remark 8.3.1 it can be T^s -equivariantly decomposed as $V_x \oplus V_x^*$. This way $\mathfrak{M}(T^s, W')_0^\alpha$ is also a hypertoric

variety (associated with the representation V_x). Denoting by $\mathcal{A}_x$ the corresponding hyperplane arrangement, we obtain an étale equivalence

$$(\pi_{\mathcal{A}}, \mathfrak{M}(\tilde{\mathcal{A}}), \mathfrak{M}(\mathcal{A}), x) \sim_{et} \mathbb{A}^m \times (\pi_{\mathcal{A}_x}, \mathfrak{M}(\tilde{\mathcal{A}}_x), \mathfrak{M}(\mathcal{A}_x), 0).$$

We point out that both in Crawley-Boevey's context and the context of hypertoric varieties the varieties which appear in the étale equivalences provided by Proposition 8.3.3 are from the same class.

8.4. Slodowy slices. Let $\mathfrak{g}$ be any split semisimple Lie algebra over $\mathbb{K}$, let G be the adjoint group of $\mathfrak{g}$ and let $\mathcal{N} \subset \mathfrak{g}$ be the nilpotent cone. Let $\mathbb{O} \subset \mathcal{N}$ be a nonzero nilpotent adjoint orbit and let $e \in \mathbb{O}$ be a nilpotent element. By the Jacobson-Morozov theorem we can find an $\mathfrak{sl}_2$ triple (e, h, f) , $h, f \in \mathfrak{g}$, associated to e , meaning that $[h, e] = 2e$, $[e, f] = h$, $[h, f] = -2f$. We fix such a triple. Let $Z(f) \subset \mathfrak{g}$ be the centraliser of f inside $\mathfrak{g}$. Then Slodowy slice $\mathcal{S}_e$ to $\mathbb{O}$ at point e is defined using the affine subspace $e + Z(f) \subset \mathfrak{g}$, namely $\mathcal{S}_e = \{e + Z(f)\} \cap \mathcal{N}$ (for more details see [Sl]). Resolution of singularities of $\mathcal{S}_e$ is given by the restriction of the Springer resolution $\pi : T^*\mathcal{B} \rightarrow \mathcal{N}$, where $\mathcal{B}$ is the flag variety of G . The map π is $G \times \mathbb{G}_m$ -equivariant, where $G \times \mathbb{G}_m$ -action on $T^*\mathcal{B}$ is given by the natural action of G on $\mathcal{B}$ and $\mathbb{G}_m$ -action along the fibers of $T^*\mathcal{B}$, while the action on $\mathcal{N}$ is given by the adjoint action of G and the natural action of $\mathbb{G}_m$ by dilation, which contracts $\mathcal{N}$ to 0. Let's denote the resolution by $\pi_e : \tilde{\mathcal{S}}_e \rightarrow \mathcal{S}_e$.

Construction of a contracting $\mathbb{G}_m$ -action on $\mathcal{S}_e$ is not very straightforward. The Lie algebra homomorphism $\mathfrak{sl}_2 \rightarrow \mathfrak{g}$ given by $\mathfrak{sl}_2$ -triple (e, h, f) extends to a Lie group homomorphism $\tilde{\gamma} : \mathrm{SL}_2 \rightarrow G$. We put

$$\gamma : \mathbb{G}_m \rightarrow G, \quad \gamma(t) = \tilde{\gamma} \begin{pmatrix} t & 0 \\ 0 & t^{-1} \end{pmatrix}.$$

Composition with the adjoint action gives a $\mathbb{G}_m$ -action on $\mathcal{N}$. The only problem is that it does not preserve e : $(\mathrm{Ad}\gamma(t))(e) = t^2e$. The action on the slice is produced from this one by normalisation: $\rho(t)(x) = t^2\mathrm{Ad}\gamma(t^{-1})(x)$. $\rho(t)$ preserves $Z(f)$ and contracts it to 0. So, since $\rho(t)(e + x) = e + \rho(t)x$, we get a $\mathbb{G}_m$ -action on $\mathcal{S}_e$, contracting it to e . Since the map π is $G \times \mathbb{G}_m$ -equivariant, this action extends to the resolution π_e turning it into a conical resolution of singularities. Moreover it is known to be an example of a symplectic resolution (with the symplectic form given by the restriction of the canonical symplectic form on $T^*\mathcal{B}$).

To prove that every point admits a conical neighbourhood we note that Slodowy slice is a slice to the orbit, so in our terms gives an étale equivalence

$$(\pi, T^*\mathcal{B}, \mathcal{N}, e) \sim_{et} (\pi_e, \tilde{\mathcal{S}}_e, \mathcal{S}_e, e) \times \mathbb{A}^{\dim \mathbb{O}}.$$

Moreover for every point $e' \in \mathcal{S}_e$ it gives an equivalence

$$(\pi, T^*\mathcal{B}, \mathcal{N}, e') \sim_{et} (\pi_e, \tilde{\mathcal{S}}_e, \mathcal{S}_e, e') \times \mathbb{A}^{\dim \mathbb{O}}.$$

Let $\mathbb{O}' \subset \mathcal{N}$ be the orbit of e' . Then $\dim \mathbb{O}' > \dim \mathbb{O}$ and from the analogous decomposition we get

$$(\pi_e, \tilde{\mathcal{S}}_e, \mathcal{S}_e, e') \times \mathbb{A}^{\dim \mathbb{O}} \sim_{et} (\pi, T^*\mathcal{B}, \mathcal{N}, e') \sim_{et} (\pi_{e'}, \tilde{\mathcal{S}}_{e'}, \mathcal{S}_{e'}, e') \times \mathbb{A}^{\dim \mathbb{O}'}.$$

Finally Lemma 5.5.1 provides from this an equivalence

$$(\pi_e, \tilde{\mathcal{S}}_e, \mathcal{S}_e, e') \sim_{et} (\pi_{e'}, \tilde{\mathcal{S}}_{e'}, \mathcal{S}_{e'}, e') \times \mathbb{A}^{\dim \mathbb{O}' - \dim \mathbb{O}}.$$

Right hand side is a conical resolution of singularities, so we obtain the following

Theorem 8.4.1. *Resolution $\pi_e : \tilde{\mathcal{S}}_e \rightarrow \mathcal{S}_e$ of singularities of the Slodowy slice is a resolution with conical slices.*

REFERENCES

- [ABM] R. Anno, R. Bezrukavnikov, I. Mirković, *Stability conditions for Slodowy slices and real variations of stability*, Mosc. Math. J., **15:2** (2015), 187-203
- [AP] M. Arbo, N. Proudfoot, *Hypertoric varieties and zonotopal tilings*, [arXiv:1511.09138](#)
- [BeKu] G. Bellamy, T. Kuwabara, *On deformation quantisations of hypertoric varieties*, Pacific J. Math. **260** (2012), no. 1, 89-127
- [BK1] R. Bezrukavnikov, D. Kaledin, *Fedosov quantization in positive characteristic*, J. Amer. Math. Soc. **21** (2008), 409-438
- [BK2] R. Bezrukavnikov, D. Kaledin, *McKay equivalence for symplectic resolutions of singularities*, [arXiv:0401002](#)
- [BL] R. Bezrukavnikov, I. Losev, *Etingof conjecture for quantized quiver varieties*, [arXiv:1309.1716v2](#)
- [BMR] R. Bezrukavnikov, I. Mirković, D. Rumynin, *Localization of modules for a semi-simple Lie algebra in prime characteristic*, Ann. of Math. (2) **167** (2008), **3**, pp. 945–991; [arXiv:math.RT/0205144](#)
- [BMS] B. Bhatt, M. Morrow, and P. Scholze, *Integral p-adic Hodge theory*, [arXiv:1602.0314](#)
- [BD] R. Bielawski, A. Dancer, *The geometry and topology of toric hyperkähler manifolds*, Comm. Anal. Geom. **8** (2000), 727-760.
- [BO] A. Bondal, D. Orlov, *Derived categories of coherent sheaves*, Proceedings of the ICM, Vol. II (Beijing, 2002), 47-56
- [Bou] N. Bourbaki, *Lie Groups and Lie Algebras. Chapters 7-9*, Springer-Verlag Berlin Heidelberg, 2005
- [CrB1] W. Crawley-Boevey, *Normality of Marsden-Weinstein reductions for representations of quivers*, Mathematische Annalen, January 2003, **325**, Issue 1, 55-79
- [CrB2] W. Crawley-Boevey, *Geometry of the Moment Map for Representations of Quivers*, Compositio Mathematica (2001) **126**: 257-293
- [CrB3] W. Crawley-Boevey, M. Holland, *Preprojective algebras, differential operators and a Conze embedding for deformations of Kleinian singularities*, Comment. Math. Helv., **74** (1999), 548-574
- [CrB4] W. Crawley-Boevey, *On the exceptional fibres of Kleinian singularities*, Amer. J. Math., **122** (2000), 1027-1037
- [CH] W. Crawley-Boevey, M. Holland, *Noncommutative deformations of Kleinian singularities*, Duke Math. J., **92**, Number 3 (1998), 605-635
- [SGA] P. Deligne, *La formule de dualité globale, exposé XVIII, SGA 4 tome 3*, Lect. Notes Math., **305**, Springer Berlin Heidelberg, 1973, 481-587
- [EGA] J. Dieudonné, A. Grothendieck, *Éléments de géométrie algébrique*, Publ. Math. IHES, **4**, **8**, **11**, **17**, **20**, **24**, **28**, **32**, (1960-7)
- [Dr] V. Drinfeld, *On algebraic spaces with an action of G_m* , [arXiv:1308.2604](#)
- [FK] B. Farbés, M. Kisin, *Rigidity, Locally Symmetric Varieties and the Grothendieck-Katz Conjecture*, International Mathematics Research Notices **2009** **22**
- [GM] S. Gelfand, Y. Manin, *Methods of Homological Algebra*, Springer Monographs in Mathematics, Springer-Verlag Berlin Heidelberg, 2003
- [HS] T. Hausel, B. Sturmfels, *Toric hyperkähler varieties.*, Doc. Math **7** (2002), 495-534.
- [Kac] V.G. Kac, *Root systems, representations of quivers and invariant theory*, Lec. Notes in Math. , **996** (Springer, 1983), 74-108.
- [Kal1] D. Kaledin, *Geometry and topology of symplectic resolutions*, Proceedings of Symposia in Pure Mathematics, 2009, Vol 80, 2, 595-628
- [Kal2] D. Kaledin, *Symplectic singularities from the Poisson point of view*, Journal für die reine und angewandte Mathematik (Crelles Journal), 2006, Issue 600, 135-156
- [Kal3] D. Kaledin, *Derived equivalences by quantization*, [arXiv:math/0504584](#)
- [Katz1] N. Katz, *A conjecture in the arithmetic theory of differential equations*, Bulletin de la Société Mathématique de France **110** (1982): 203-239.
- [Katz2] N. Katz, *Nilpotent connections and the monodromy theorem: applications of a result of Turrittin*, Publications Mathématiques de l’IHÉS, **39** (1970), 175-232.
- [Kaw] Y. Kawamata, *D-equivalence and K-equivalence*, J. Differential Geom. **61** (2002), 147-171
- [KN] G. Kempf, L. Ness, *The length of vectors in representation spaces*, Lecture Notes in Math., **732**, Berlin, New York: Springer-Verlag, 233-243
- [Ki] A.D. King, *Moduli of representations of finite-dimensional algebras*, Quart. J. Math. Oxford, **45** (1994), 515-530
- [La] R.K. Lazarsfeld, *Positivity in Algebraic Geometry I*, Springer-Verlag Berlin Heidelberg, 2004
- [Lu] D. Luna, *Slices étales*, Bull. Soc. Math. France, Mémoire, **33** (1973), 81-105.

- [Ma] Y. Matsushima, *Espaces homogènes de Stein des groupes des Lie complexes*, Nagoya Math. J., **16** (1960), 205-216
- [Mi] J. Milne, *Étale Cohomology*, Princeton University Press, Princeton, New Jersey, 1980
- [Mo] I. Moergijk, *Introduction to the language of stacks and gerbes*, [arXiv:0212266](#) no. 3, 515–560.
- [Na1] Y. Namikawa, *Poisson deformations and birational geometry*, [arXiv:1305.1698](#)
- [Na2] Y. Namikawa, *A finiteness theorem on symplectic singularities*, [arXiv:1411.5585](#)
- [OV] A. Ogus, V. Vologodsky, *Nonabelian Hodge theory in characteristic p* , V. Publ. math. IHES **106** (2007) pp. 1-138
- [P1] D. Popescu, *General Néron desingularization*, Nagoya Math Journal, **100** (1985), pp. 97-126.
- [P2] D. Popescu, *General Néron desingularization and approximation*, Nagoya Math Journal, **104** (1986), pp. 85-115.
- [P3] D. Popescu, *Letter to the editor: General Néron desingularization and approximation*, Nagoya Math Journal, **118** (1990), pp. 45-53.
- [PW] N. Proudfoot, B. Webster, *Arithmetic and topology of hypertoric varieties*, Journal of Algebraic Geometry, **16** (2007), 39-63.
- [Sl] P. Slodowy, *Simple singularities and simple algebraic groups*, Lecture Notes in Math., **815**, Springer Verlag, 1980.

(Dmitry Kubrak) MASSACHUSETTS INSTITUTE OF TECHNOLOGY, DEPARTMENT OF MATHEMATICS, 77 MASSACHUSETTS AVE, CAMBRIDGE, MA 02139 USA

E-mail address: dmkubrak@gmail.com

(Roman Travkin), INSTITUTE FOR ADVANCED STUDY, SCHOOL OF MATHEMATICS, 1 EINSTEIN DRIVE, PRINCETON, NJ 08540 USA

E-mail address: travkin@alum.mit.edu